



This document is a postprint version of an article published in Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology © Elsevier after peer review. To access the final edited and published work see <https://doi.org/10.1016/j.cbpa.2018.11.005>

1 **The effect of algal turbidity on larval performance and the ontogeny of digestive enzymes in the**
2 **grey mullet (*Mugil cephalus*)**

3

4 William Koven^{1*}, Enric Gisbert², Oriya Nixon¹, Mikhail M. Solovyev^{3,4}, Aviad Gaon¹, Guy Allon¹,
5 Iris Meiri-Ashkenazi¹, Amos Tandler¹, Hanna Rosenfeld¹

6

7 Corresponding author: William Koven

8 Israel Oceanographic and Limnological Research, the National Center for Mariculture, P.O.B. 1212,
9 Eilat 88112, Israel

10 BMKoven@gmail.com

11 Tel: +972 8 6361 443, Cell: +972 523 467 822,

12 Fax: +972 8 6375 761

13

14 ¹*Israel Oceanographic and Limnological Research, the National Center for Mariculture (NCM),*
15 *P.O.B. 1212, Eilat 88112, Israel. Email: BMKoven@gmail.com*

16 ²*IRTA, Centre de Sant Carles de la Ràpita (IRTA-SCR), Programa d'Aqüicultura, Crta. del Poble*
17 *Nou Km 5.5, 43540 Sant Carles de la Ràpita, Spain. Email: Enric.Gisbert@irta.cat*

18 ³*Institute of Systematics and Ecology of Animals, Siberian Branch of Russian Academy of Sciences,*
19 *11 Frunze St., Novosibirsk 630091, Russia. Email: Yarmak85@mail.ru*

20 ⁴*Tomsk State University, Tomsk 634050, Russia. Email: Yarmak85@mail.ru*

21 Key words: grey mullet; digestive tract; larvae; turbidity; algae; enzymes; gut maturation

22

23 **Abstract**

24 A study comprised of two trials was carried out to determine and compare the effects of water
25 turbidity produced by live microalgae and inert clay particles on the larval rearing of grey mullet
26 (*Mugil cephalus*). Trial 1 evaluated the effect water turbidity generated by microalgae on grey mullet
27 larval performance and digestive enzyme activities along ontogeny. In particular, two microalgae
28 (*Nannochloropsis oculata* and *Isochrysis. galbana*) were used, whereas water turbidity levels (0.76
29 and 1.20 NTU) and a non-microalgae control (0.26 NTU) were tested from 2 to 23 dph grey mullet
30 larvae (5 treatments). The higher turbidity (1.2 NTU) larvae (5 dph) consumed markedly ($P < 0.05$)
31 more rotifers than other treatment fish, independently of the microalgae type. Alkaline phosphatase
32 activity was *ca.* 8 times higher and α -amylase activity increased 5.3 times in 79 dph fish compared to
33 40 dph individuals. The ratio of alkaline phosphatase and leucine-alanine aminopeptidase indicated
34 gut maturation occurred around 61 dph, as well as a transition from carnivorous to omnivorous
35 feeding habits. Trial 2 compared the most effective *N.occulata* produced turbidity level (1.2 NTU)
36 with the identical water turbidity produced by inert clay on larval performance. *M. cephalus* larvae
37 exposed to high algal turbidity demonstrated superior performance ($P < 0.05$), in terms of rotifer
38 ingestion, dry weight gain and survival, in comparison to cohorts reared under the clay treatment as
39 well as a lower microalgae produced turbidity. These findings suggested that water algal turbidity is
40 not the dominant factor determining improved grey mullet larval performance.

41

42 **1. Introduction**

43 In the commercial rearing of marine fish larvae, tanks are frequently “greened” with microalgae such
44 as *Nannochloropsis oculata* or *Isochrysis galbana*. It is widely believed and demonstrated that the
45 provision of these algae into the tanks significantly improves larval performance and has become an
46 inseparable part of commercial rearing protocols in fish farms around the Mediterranean basin

47 (Papandroulakis et al., 2002; van der Meeren et al., 2007; Bentzon-Tilia et al., 2016). On the other
48 hand, it remains speculative how algal supplementation contributes to larval growth and survival or
49 whether this benefit is species-specific. The biochemical composition of algal species (e.g. fatty
50 acids) varies considerably and it is entirely possible that particular compounds secreted from the algal
51 cell (e.g. polysaccharides) and/or are released during digestion might stimulate the immune system
52 or enhance the digestive process in larvae (Hemaiswarya et al., 2011). In addition, water turbidity
53 from specific algal concentrations may modify the light milieu for larvae, providing optimal
54 backlighting for larvae to facilitate live prey identification (e.g. rotifers), foraging behavior and
55 thereby enhancing hunting success (Rocha et al., 2008).

56 The grey mullet (*Mugil cephalus*) is an economically important euryhaline and eurythermal species
57 contributing to sizable fisheries of estuarine and coastal regions in a variety of countries. It has been
58 traditionally farmed extensively in ponds and enclosures in the Mediterranean region, South East
59 Asia, Korea, Taiwan, China, Japan and Hawaii (FAO, 2018) and has been recently recognized as a
60 highly valued candidate for more intensive aquaculture (Whitfield et al., 2012). Nevertheless, the
61 rearing of the early developmental stages of grey mullet and the mass production of robust, fast
62 growing juveniles remains an obstacle to the successful domestication of this species.

63 A study comprised of two trials was carried out to determine and compare the effects of water
64 turbidity produced by live microalgae and inert clay particles on the larval rearing of grey mullet.
65 More specifically, the aims of the first trial were to (1) investigate the effect of microalgae produced
66 turbidity in the rearing tank on larval performance, in terms of prey capture efficiency, growth and
67 survival, as well as digestive tract enzyme ontogeny. (2) Determine whether turbidity effect varies
68 with microalgae type (*Nannochloropsis oculata* vs. *Isochrysis galbana*) and concentration (cells ml⁻¹).
69 The aim of the second trial was to elucidate whether water turbidity or algal biochemical
70 composition were the dominant factor promoting improved larval performance in this species.

71

72 **2. Material and methods**

73 *2.1 Trial 1: Evaluation of the effect of water turbidity from two species of live microalgae on larval* 74 *and juvenile grey mullet performance and ontogeny of the digestive tract functionality*

75 Grey mullet eggs (gastrula stage) were stocked in fifteen 1.5 m³ V-tanks (100 eggs l⁻¹) in a flow
76 through system where filtered (10 µm), UV-treated, 40 ‰ seawater (25 °C) entered from the tank
77 bottom at a rate of two tank exchanges per day. Two experimental water turbidity levels A (0.76
78 NTU) and B (1.20 NTU) were tested using two microalgae species (*N. oculata* and *I. galbana*) and
79 compared to non-microalgae supplementation C (control; 0.26 NTU) in 2 to 23 days post hatch (dph)
80 grey mullet larvae. This meant that each of the 5 treatments; C (Control), Nanno A (*N. oculata*-
81 turbidity A), Iso A (*I. galbana*-turbidity A), Nanno B (*N. oculata*-turbidity B), Iso B (*I. galbana*-
82 turbidity B) were investigated with three replicate tanks per treatment. The different microalgae
83 turbidity treatments and their concentrations are listed in Table 1.

84 The hatching rate (%) of stocked grey mullet eggs and survival of the pre-larva at the end of the day
85 of hatching (0 dph) were calculated by placing a fertilized egg, at the gastrula stage, in each of 12
86 wells (5 mL) in each of three plastic well plates. The plates were covered and placed in a temperature
87 controlled incubator until hatching where the emerging larvae and surviving newly hatched larvae at
88 the end of 0 dph were counted. After hatching in the tanks, water salinity was progressively lowered
89 to 25 ‰ and the flow rate reduced to one tank water exchange per day. Lighting over the tanks
90 provided 500 lux (14 h light day⁻¹) at the water surface. At 2 dph, once the yolk sac was depleted, the
91 eyes pigmented and the mouth and anus opened. From this point until 16 dph, grey mullet larvae were
92 offered rotifers (*Brachionus rotundiformis*) that were previously enriched with taurine (600 mg L⁻¹)
93 and essential fatty acids (Red PepperTM, Bernaqua, Belgium) for 12 and 8 h, respectively. After

94 feeding rotifers exclusively to 16 dph, larvae were co-fed with enriched rotifers and *Artemia*
95 *metanauplii* (Red Pepper™) until 24 dph. From 25 to 57 dph, fish were offered a 1:1 (w/w) mixture
96 of the weaning diet Caviar™ (Bernaqua, Belgium) and dried and powdered *Ulva lactuca*, which was
97 produced at the IOLR (Eilat, Israel). After 57 dph, fish were fed only the starter feed Ranaan Dry feed
98 (RDF, Israel) until the end of the trial at 79 dph (Table 2).

99

100 2.2 Trial 2: Comparison of the effect of water turbidity from live microalgae and white clay particles 101 on larval and juvenile grey mullet performance

102 In order to test whether the advantage of water turbidity on larval performance is independent of
103 turbidity source (live algae or clay), trial 2 compared the most effective turbidity level produced by
104 live algae and white clay with a lower algae produced turbidity level as a control. The experimental
105 treatments consisted of: (1) low water turbidity (0.8 ± 0.04 NTU; 0.25×10^6 cells of *N. oculata* mL⁻¹)
106 ¹), (2) high water turbidity (1.2 ± 0.04 NTU; 0.50×10^6 cells of *N. oculata* mL⁻¹) and (3) high water
107 turbidity (1.2 ± 0.02 NTU) produced by white clay, which was purchased from DAS Terracotta.
108 F.I.L.O. (Pero, Italy). In this trial, grey mullet eggs, at gastrula stage, were stocked in twelve 1.5 m³
109 V-tanks (100 eggs L⁻¹). The mullet larvae were exposed to the treatments from 2-30 dph while length
110 was measured daily to 29 dph and survival at 50 dph. Larval rearing and husbandry conditions, as
111 well as feeding protocols, were similar to those previously mentioned in trial 1.

112

113 2.3 Water turbidity assessment and control

114 All turbidity values in these trials were determined on triplicate water samples from each tank,
115 including the control no algae treatment, which were first filtered (40 µm mesh size) before being
116 read with a Turbidometer (Lovibond Turbi-check, Amesbury, England). The turbidity value for

117 Nanno B (1.20 NTU) was based on the current concentration of *N. oculata* (0.5×10^6 cells mL⁻¹) used
118 at the IOLR for larval rearing. Previous studies carried out at the IOLR (unpublished data)
119 demonstrated that levels above this concentration significantly reduced rotifer consumption in
120 gilthead sea bream (*Sparus aurata*) larvae. *I. galbana* has a cell size of *ca.* 5 µm, while *N. oculata* is
121 *ca.* 1.5 µm. The concentration of the larger *I. galbana* needed for achieving a turbidity value of *ca.*
122 1.20 for Iso B was empirically determined. The turbidity values for Iso and Nanno A were based on
123 half the *I. galbana* or *N. oculata* concentrations used to achieve the B turbidity values and were
124 empirically determined. Microalgae and the clay were added twice daily during the morning (08:30)
125 and afternoon (14:30) in the larval rearing tanks, before larval feeding, in order to maintain stable
126 water turbidity levels. Algae in the rearing tanks were diluted due to constant water exchange.
127 Consequently, the afternoon algal addition was based on measured turbidity levels, which were
128 increased to designated values.

129

130 *2.4 Determination of rotifer consumption, growth and survival in grey mullet larvae*

131 In order to determine rotifer consumption, five larvae were sampled from each tank in the
132 experimental system 90 min after feeding them with enriched rotifers. Larvae were then sacrificed,
133 according to ethical standards, with an overdose of the anesthetic tricaine methanesulfonate (MS-222,
134 Sigma-Aldrich, Rehovot, Israel) and fixed in 10% buffered formalin and stored at 4 °C until counting
135 the mastaxes (indigestible part of the rotifer's feeding apparatus) in the fish's digestive tract.
136 Anecdotal observations determined that mastaxes can be expelled 2-3 h after feeding suggesting that
137 90 min would allow a good approximation of rotifer consumption.

138 Fish growth was measured as dry weight (DW) when the larvae were 15, 18 and 25 dph at the end of
139 the turbidity Experiment 1 and at 29 dph at the end of Trials a and b in Experiment 2. This was

140 carried out by collecting *ca.* 130 larvae per tank, sacrificing them with an excess of MS-222 and then
141 washing them with distilled water. Fish samples were then dried at 70 °C for 24 h followed by their
142 weighing (A&D HD-120 analytical balance, Japan).

143 Due the minute size and rapid deterioration of dead larvae, it was not possible to accurately follow
144 mortality in large tanks during the course of the experiment. Consequently survival values were
145 determined only at the end of both trials and were expressed as a percentage of the surviving fish,
146 when the fish were harvested, over the number of the fertilized eggs initially stocked (adjusted for
147 hatching rate and survival after 24 h) in tanks and taking into account the number of sampled fish for
148 analytical purposes. Samples for measuring the digestive tract enzyme activities in the turbidity trial
149 were sampled at hatching, 18, 25, 40, 61 and 79 dph.

150

151 *2.4 Digestive enzyme activities*

152 In trial 1, it was not possible to separate tail and trunk musculature from the abdominal region of
153 whole larvae younger than 60 dph, while older fish (60 and 79 dph) were dissected to separate the
154 pancreatic and intestinal segments. Digestive enzyme activities were determined using the following
155 numbers of fish (3 replicates per experimental condition): 350-400 fish at 18 dph, 92-172 fish at 25
156 dph, 12 fish at 41 dph, 3-6 fish at 61 dph, and 4-6 fish at 79 dph. Dissection was conducted under a
157 dissecting microscope on a pre-chilled glass plate maintained at 0 °C and lyophilized (FD-80,
158 Boyikang, China) and shipped to IRTA's facilities for their analysis. For quantifying the activity of
159 enzymes, lyophilized samples were homogenized (Ultra-Turrax T25 basic, IKA[®]-Werke, Germany)
160 in 30 volumes (v/w) of mannitol (50 mM mannitol, 2 mM Tris-HCl buffer; pH 7.0), centrifuged and
161 the supernatant removed for enzyme quantification. Then, 1 mL of the supernatant was stored at -20
162 °C for leucine–alanine peptidase quantification and the rest of the homogenate was used for brush

163 border purification (Crane et al., 1979; Gisbert et al., 2018). Enzyme activities for pancreatic and
164 intestinal enzymes (U mg protein^{-1}) were determined as described in Gisbert et al. (2009) and
165 processed within 15 days to keep their activities intact (Solovyev and Gisbert, 2016). Trypsin and
166 chymotrypsin activities were assayed using BAPNA (N- α -benzoyl-DL-arginine p-nitroanilide) and
167 BTEE (benzoyl tyrosine ethyl ester) as substrates (Holm et al., 1988; Worthington, 1991).
168 Chymotrypsin activity was only assayed in the 79 dph group, whereas for younger groups the activity
169 was below the detection limit levels of the assay. Alpha-amylase was measured using starch as
170 substrate (Métais and Bieth, 1968); bile salt-activated lipase activity was assayed using p-nitrophenyl
171 myristate as substrate (Iijima et al., 1998). Alkaline phosphatase activity was measured using 4-
172 nitrophenyl phosphate (Bessey et al., 1946); the assay of the cytosolic peptidase, leucine-alanine
173 aminopeptidase was performed using leucine-alanine as substrate (Nicholson and Kim, 1975).
174 Soluble protein of extracts was quantified by means of the Bradford's method (Bradford, 1976). All
175 the assays were made in triplicate from each pool of larvae (biological replicate) and absorbance read
176 using a spectrophotometer (Tecan™ Infinite M200, Switzerland).

177

178 2.5 Statistics

179 Statistical analyses were carried out using GraphPad Prism version 5.00 for Windows (GraphPad
180 Software, San Diego California USA, www.graphpad.com). All data are presented as mean \pm SEM.
181 Outliers were identified by calculation of the Z value using the Grubbs test (Rousseeuw and Leroy
182 2003) and removed if calculated Z value was higher than the tabulated value. Every fish sampled for
183 mastax measurement was considered a treatment replicate (15-20 larvae from each age of 2-5 dph
184 were sampled per treatment). Percentage data values were first arcsine transformed, and then analyzed
185 by one-way ANOVA and Barlett's test for equal variances. If significance ($P < 0.05$) was found after
186 ANOVA analysis while Barlett's test was not significant ($P > 0.05$); then, testing differences between

187 groups was carried out by Newman-Keuls Multiple Comparison test. In cases where ANOVA and
188 Barlett's test were both significant ($P < 0.05$); then, the non-parametric Kruskal Wallis Test was
189 applied followed by Dunn's multiple Comparison test to determine significant ($P < 0.05$) differences
190 among treatments. Regression data sets employed Akaike's Information Criteria (AIC) to compare
191 linear, second order polynomial and other models to determine which most likely generated the data.
192 The effect of water turbidity on the overall activity of pancreatic and intestinal enzymes was evaluated
193 by Principal Component Analysis (PCA) in 79 dph fish (Statistica 7.0, StatSoft, Inc.).

194 *2.6 Ethics statement*

195 All animal experimental procedures were conducted in compliance with the Guidelines of the
196 European Union Council (86/609/EU) for the use of laboratory animals.

197

198

199 **3. Results**

200 *3.1 Trial 1: the effect of water turbidity from two species of live microalgae on larval and juvenile* 201 *grey mullet performance and ontogeny of the digestive tract functionality*

202 Figure 1a shows that the water turbidity (NTU) values for different experimental conditions,
203 Turbidity values of treatments using microalgae regardless of the microalgae and cell density used
204 were significantly different from the control group (no algae) from 1 to 23 dph ($P < 0.05$) In Figure
205 1b the turbidity (NTU) values in trial 2 are shown comparing the Clay B treatment (1.2 NTU) with
206 the low (0.75 NTU) Nanno A and high Nanno B (1.2 NTU) microalgae treatments. The turbidity
207 values of the Nanno B and Clay B were very similar while both of these microalgae treatments were
208 significantly ($P < 0.05$) different from those of Nanno A on each day from 2 to 24 dph.

209 *Trial 1: rotifer consumption, larval survival and growth under the different turbidity conditions*

210 Figure 2a demonstrated a significant ($P < 0.05$) water turbidity effect on rotifer consumption in grey
211 mullet larvae aged 5 dph that was independent of the type of microalgae used for increasing water
212 turbidity. Larvae in the Iso B and Nanno B treatments consumed significantly ($P < 0.05$) more rotifers
213 than in the Nanno A and Control groups. The same trend was observed for the Iso A treatment, it but
214 was not significant ($P > 0.05$). Interestingly, this pattern of rotifer consumption was very similar to
215 that of fish survival much later on (Fig. 2b). Fish exposed to high microalgae water turbidity levels
216 (Iso B, Nanno B) from 2 to 23 dph survived significantly ($P < 0.05$) better at 51 dph, which was about
217 4 weeks after the treatments had been discontinued, than fish feeding at the lower water turbidity
218 values (Iso A, Nanno A) or in clear water (control). However, despite the effect of water turbidity on
219 prey consumption and larval survival, there was no significant ($P > 0.05$) treatment effect on their
220 growth performance (Fig. 2c).

221

222 *Trial 1: digestive enzyme specific activity as a function of fish age, microalgae water turbidity levels*
223 *and diet*

224 Trial 1 results indicated that water turbidity treatments tested from 2 to 23 dph larvae had no obvious
225 effects on the specific activities of digestive enzymes when they were measured between 18 and 79
226 dph (Figs. 3-5). On the other hand, the results from PCA (Fig. 6) suggested that lower water turbidity,
227 regardless of the microalgae used, was correlated with higher pancreatic and intestinal enzyme
228 activity in 79 dph grey mullet and much less so to the control and the high turbidity treatments. The
229 exceptions of Nanno A and Nanno B samples showed intermediate values in comparison to the rest
230 of the other groups.

231 Diet composition during the period of enzyme sampling did significantly ($P < 0.05$) affect the specific
232 activity of bile-salt activated lipase and alkaline protease. Bile salt-activated lipase decreased from
233 18 dph (Fig. 3a), when larvae were feeding on high lipid containing rotifers and *Artemia metanauplii*
234 (Table 2), to the lower dietary lipid levels of the co-feeding period based on Caviar: *U. lactuca* (1:1
235 w/w) diet from 25-57 dph (Table 2). However, after switching to the RDF diet at 57 dph, the increased
236 lipid of this feed (14%) resulted in a marked increase of the bile salt-activated lipase activity ($P <$
237 0.05). Similarly, dietary protein decreased from the high levels found in rotifers and *Artemia*
238 *metanauplii* (2 to 25 dph) (Table 2), which corresponded to high alkaline protease activity, to the
239 reduced protein levels of the Caviar: *U. lactuca* (1:1) diet fed between 25 and 57 dph with the
240 subsequent decrease in alkaline protease activity (Fig. 3b). However, the activity of these enzymes
241 tended to increase when fish were ingesting the higher protein levels of the RDF diet from 58 to 79
242 dph (Fig. 3a, b).

243 In contrast, trypsin activity was not affected by the diet throughout the entire sampling period (Fig.
244 3c; $P > 0.05$). The cytosolic enzyme leucine-alanine aminopeptidase (LAP) significantly ($P < 0.05$)
245 decreased in all microalgae treatments from 18 to 61 dph regardless of the live prey provided, but
246 then increased ($P < 0.05$) in 79 dph fish when the fish were fed the RDF diet (Fig. 3d). The activity
247 of the digestive tract marker for brush border membrane (BBM) development, alkaline phosphatase
248 (AP), significantly increased 8.0 times in fish from 40 to 79 dph (Fig. 4a; $P < 0.05$). Consequently,
249 the AP/LAP ratio, an indicator of gut maturation, in almost every treatment peaked at 61 dph, but
250 then declined in 79 dph fish (Fig. 4b). During the period between 40 to 79 dph, α -amylase specific
251 activity (Fig. 4c) increased 5.3 times independently of diet. The alkaline protease/lipase ratio values
252 showed higher protease activity over bile salt-activated lipase levels at 18 dph when larvae were
253 feeding on highly digestible rotifer and *Artemia* protein sources (Fig. 4d). However, total alkaline
254 protease activity dropped compared to bile salt-activated lipase activity at 25 dph, when the fish began

255 to feed on the Caviar: *U. lactuca* diet. On the other hand, the ratio increased significantly ($P < 0.05$)
256 from 25 to 61 dph. Moreover, the amylase/trypsin and amylase/protease ratios supported the
257 increasing capability of amylase production with age (Fig. 5a, b).

258

259 *3.2 Trial 2: larval rotifer consumption, growth and survival*

260 The effect of Nanno A, Nanno B and Clay B treatments on rotifer (mastax) consumption, DW in 30
261 dph larvae and survival in 50 dph fish and are shown, respectively, in Fig. 7a, b, c. The results from
262 trial 2 were in agreement with those of trial 1 and demonstrated that grey mullet larvae exposed to
263 the higher microalgae produced turbidity outperformed ($P < 0.05$), in terms of food ingestion, growth
264 and survival, their cohorts reared under lower microalgae produced turbidity as well as larvae
265 exposed to the identical high turbidity level (1.2 NTU) produced from clay.

266

267 **4. Discussion**

268 The beneficial effects of the presence of microalgae in the rearing tanks of the larvae of many farmed
269 species have long been recognized (Naas et al., 1992; Reitan et al., 1997; Cahu et al., 1998; Lazo et
270 al., 2000; Skiftesvik et al., 2003; Faulk and Holt, 2005). In support of this, different authors have
271 reported the importance of microalgae addition in larval rearing tanks for several mullet species such
272 as the stripped grey mullet (Tamaru et al, 1994) and the thick lipped grey mullet, *Chelon labrosus*
273 (Ben Khemis et al., 2006). Various hypotheses have been postulated for explaining how microalgae
274 might benefit larvae, including (1) providing a direct supply of micronutrients (Van Der Meeren,
275 1991) that trigger key physiological processes (Hjelmeland et al., 1988), (2) releasing of appetite
276 stimulating components (Stottrup et al., 1995), and (3) influencing the bacterial composition of the
277 rearing water and consequently, the larval gut microbial flora (Skjermo and Vadstein, 1993; Bentzon-

278 Tilia et al., 2016). Apart from these potential advantages, a number of authors have also suggested
279 that microalgae turbidity creates a backlighting effect that would contrast the zooplankton prey
280 against their background facilitating larval foraging behavior and in particular, prey detection and
281 hunting success by the larvae (van der Meeren, 1991, Utne-Palm, 2002). In the present study, the
282 increased consumption of rotifers at the higher water turbidity levels (*ca* 1.20 NTU), independently
283 of the microalgae species used, appeared to suggest that the turbidity and the subsequent backlighting
284 effect of the microalgae was the main factor influencing rotifer ingestion and larval performance. In
285 Atlantic halibut, *Hippoglossus hippoglossus* (Naas et al., 1992) and striped grey mullet (Tamaru et
286 al., 1994), it was suggested that the microalgae suspension enhanced visual contrast, allowing larvae
287 to better detect their prey. On the other hand, Rocha et al. (2008) argued that the larval prey capture
288 was influenced by both the effect of algae on water light conditions, as well as by the substances
289 provided by their presence, although the extent of this effect on fish larvae might be species-specific.
290 In fact, an equally plausible interpretation of the results from the first trial was that there are
291 microalgae unidentified factors that are common to both *I. galbana* and *N. oculata*, triggering key
292 physiological processes that modulate prey consumption and larval performance. This line of
293 reasoning is clearly reinforced by the result showing that clay, added to the tank at the same turbidity
294 as the most effective microalgae concentration (Nanno B), performed markedly less well, in terms of
295 prey ingestion, growth and survival. Overall, the present study suggests that in grey mullet larvae,
296 water turbidity played a more subordinate role compared to the likely contribution of microalgae
297 compounds that were released into the water and/or absorbed by the larval digestive tract. On the
298 other hand, the benefit of clay may be species specific and/or vary depending on the rearing
299 conditions. In Atlantic halibut larval rearing, the addition of inorganic clay during the first feeding
300 phase reduced opportunistic pathogenic bacteria such as *Vibrio* spp. in the tank water compared to
301 microalgae supplementation (Bjornsdottir et al., 2011). Similarly, clay addition reduced bacterial load
302 in the larval rearing tanks of cod (*Gadus morhua*) (Attramadal et al., 2012). In fact, the replacement

303 of microalgae in order to achieve so-called “environmental shading” with inorganic clay has become
304 the industrial standard in a number of halibut hatcheries in Norway. However, the negative effects
305 of very high turbidity can outstrip the benefit of microalgae or clay addition when fish are visual
306 feeders (Confer et al., 1978; Gregory and Northcote, 1993).

307 The addition of microalgae to larval rearing tanks has been shown to improve survival in a number
308 of species such as halibut (Naas et al., 1992), cod (van der Meeren et al., 2007), European sea bass,
309 *Dicentrarchus labrax* (Cahu et al., 1998) and cobia, *Rachycentron canadum* (Faulk and Holt, 2005).

310 Interestingly, the significant effect of live microalgae addition and water turbidity level on rotifer
311 consumption in 5 dph larvae was in agreement with the larval survival results measured in 51 dph
312 fish (28 days after the microalgae treatments had discontinued). This suggests that larval survival, to
313 a large extent, is determined early on during rotifer feeding. In fact, the onset of exogenous feeding
314 after the reabsorption of the yolk sac is a critical time during larval development that is frequently
315 characterized by massive mortality (Yufera and Darias, 2007) if young fish are not consuming
316 sufficient levels of prey such as rotifers. Normal neural development, gut maturation, muscle function
317 and growth can be compromised leading to mortality if there is any delay or reduction in first feeding
318 (Gisbert et al. 2004).

319 It is conceivable that larvae reared under the higher microalgae produced water turbidity were
320 surviving better as they were consuming more rotifers and digesting them more efficiently as gut
321 maturation would be accelerated. In support of this, the influence of green water on enhanced
322 pancreatic enzyme production and accelerated brush border membrane development and its
323 subsequent enzyme activity improved survival in European sea bass larvae (Cahu et al., 1998;
324 Zambonino-Infante and Cahu, 1994). However, in the present study no significant differences were
325 found in enzyme activity among larvae from the different turbidity treatments from 18 dph onwards.
326 This may have been due to sampling larvae after the beneficial effect of green water on the maturation
327 of the digestive system was no longer detectable. Cahu et al. (1998) described in European sea bass

328 that the effect of microalgae addition on the activity of digestive enzymes was only observed until 16
329 dph, but not in older fish.

330

331 In contrast, the PCA results of our study implied that 1 to 23 dph larvae exposed to a lower water
332 turbidity of 0.76 NTU, independent of the microalgae species producing it, appeared to be associated
333 with higher digestive enzyme activities in 79 dph juveniles than their cohorts in the control (0.26
334 NTU) and high turbidity (*ca.* 1.20 NTU) treatments. This suggests that relatively low levels of
335 microalgae addition is necessary to stimulate digestive enzyme activity, whereas a higher microalgae
336 concentration would be less stimulatory but potentially more effective to enhance other physiological
337 pathways modulating survival and growth (e.g. stimulating the immune system). Taken altogether,
338 the benefits of “greening” the larval rearing tanks on fish performance later on in development, likely
339 includes a range of factors that are microalgae concentration and species specific dependent.

340 In order to analyze the ontogeny of digestive enzymes in 60 and 79 dph juveniles in this study, it was
341 necessary to separate the tail and trunk musculature from the abdominal region in order to prevent
342 the inclusion of excess protein in the sample homogenate. This would have led to underestimating
343 digestive tract enzyme activities. On the other hand, dissecting out intestinal and pancreatic tissues in
344 less than 60 dph larvae was technically not possible. Consequently, enzyme analyses on whole body
345 larvae was done in all fish younger than 60 dph larvae. Nevertheless, this was considered acceptable
346 as an estimation of intestinal and pancreatic enzyme activity as larval trunk and tail musculature
347 contain considerably less protein than these tissues in juvenile fish.

348

349 The composition of the diets that grey mullet were consuming during development appeared to have
350 influenced the specific activities of bile-salt activated lipase and total alkaline proteases. It should be
351 noted that no acid protease activity was found in experimental samples (data not shown), regardless
352 of the presence of a stomach (Oren, 1981). This suggested that protein digestion in grey mullet larvae

353 is mainly accomplished by alkaline proteases as found in other marine fish larvae and early juveniles
354 (Gisbert et al., 2009). *Ulva* spp. are a relatively rich source of starch (Korzen et al., 2016) where fish
355 α -amylase can hydrolyze the α -1, 4 glycoside bonds of this dietary component. This argues that the
356 increase in α -amylase activity from 25-61 dph, which was likely genetically directed, resulted in an
357 increasing ability to digest *U. lactuca* carbohydrate and potentially exposing more *Ulva* protein for
358 protease digestion. The maltose resulting from amylase digestion is absorbed as glucose after maltase
359 brush border activity. Conceivably, the catabolism of glucose as an energy source might be protein
360 sparing and therefore promote growth. The suggestion that the ontogeny of this carbohydrase is
361 genetically programmed is supported by similar high α -amylase activities found in grey mullet fry
362 that were weaned onto starch poor diets that were rich in fish meal or with a high level of fish meal
363 substitution by plant proteins (Zoutien et al., 2008; Gisbert et al., 2016).

364 In fact, the ontogeny of digestive enzyme activities in grey mullet larvae and juveniles, in general,
365 appeared to be more a function of age and genetic programming than dietary modulation. A case in
366 point are the two enzymes indicative of enterocyte development; (1) the brush border membrane
367 (BBM) alkaline phosphatase, which is a marker of nutrient absorption, and (2) the cytosol based
368 leucine-alanine peptidase (LAP), which is involved in protein intracellular digestion (pinocytosis) in
369 the cytosol of enterocytes. As the BBM develops together with increasing enzyme activity, there is a
370 parallel decrease of and dependence on intracellular digestion activity, resulting in an increase of the
371 AP/LAP ratio (Ma et al., 2005) until reaching the adult mode of digestion. The AP/LAP ratio abruptly
372 increased in *C. labrosus* at 8 dph and then significantly decreased to 36 dph. This prompted the
373 authors to conclude that gut maturation took place rapidly and early in this species. In contrast, the
374 present study on grey mullet showed that AP activity dramatically increased an average of *ca.* 8 fold
375 between 40 and 79 dph, where the peak AP/LAP ratio was detected at 61 dph. In fact, there was a
376 marked drop in the ratio at 79 dph caused by LAP levels abruptly increasing. Zouiten et al. (2008)
377 studying *C. labrosus* found a similar, but much earlier AP/LAP ratio pattern. These results argue for

378 a late maturation of the BBM and/or the transition from a carnivorous to an omnivorous/herbivorous
379 mode of feeding as grey mullet juveniles swim to estuarine environments (Oren, 1981). In addition,
380 the late increase in LAP levels may be due to the lack of acid proteases in grey mullet requiring the
381 combination of both extracellular (intestinal lumen) and intracellular (enterocyte) digestion in order
382 to process proteins more effectively.

383 The suggested late age of the carnivorous-herbivorous shift in grey mullet is supported by the steadily
384 increasing activity of α -amylase from 25-79 dph, particularly between 40 to 79 dph. Alpha-amylase
385 activity is much higher in herbivorous and omnivorous fish compared to carnivores (Solovyev et al.,
386 2015). Overall, these results suggested that 61-79 dph grey mullet juveniles, which approximate the
387 age of this species moving to estuaries (Gisbert et al., 2016), have the capacity to digest both protein
388 and starch. This allows for the exploitation of the relatively starch rich microalgae (Zemke-White and
389 Clements, 1999) and macroalgae (Horn et al., 1989) as well as benthic protein rich organisms
390 characterizing these lower salinity estuarine waters (Oren, 1981). Taken one step further, the results
391 broadly suggest that aquaculture feeds at this developmental stage should include not only
392 considerable protein but also higher levels of starch or other low cost amylolytic energetic compounds
393 compared to starter feeds fed to younger grey mullet or the juvenile stages of carnivorous species.

394

395 **5. Conclusions**

396 The results suggest that in grey mullet larvae, water turbidity played a more subordinate role
397 improving fish performance than the contribution of microalgae chemical composition. On the other
398 hand, microalgae supplementation during larval rearing did not have a conclusive effect on digestive
399 tract enzyme ontogeny. Amylase activity steadily increased between 40 to 79 dph while the fish
400 retained considerable protein digestion capability. This coupled with the late maturation of the gut at
401 61 dph argue for the capability of both carbohydrate and protein digestion as grey mullet juveniles
402 transit from carnivory to omnivory while swimming to estuarine environments. This would allow for

403 the exploitation of the relatively starch rich microalgae and macroalgae as well as benthic, protein
404 rich organisms characterizing these waters.

405

406 **Acknowledgements**

407 This study was funded by the 7th Framework Program “Diversify- Exploring the biological and socio-
408 economic potential of new/emerging candidate fish species for the expansion of the European
409 aquaculture industry (project no. 603121)

410

411

412 **References**

413 Attramadal, K. J. K., Tøndel, B., Salvesen, I., Øie, G., Vadstein, O., Olsen, Y., 2012. Ceramic clay
414 reduces the load of organic matter and bacteria in marine fish larval culture tanks. *Aquaculture*
415 *Engineering* 49, 23-34.

416 Ben Khemis, I., Zouiten, D., Besbes, R., Kamoun, F., 2006. Larval rearing and weaning of thick
417 lipped grey mullet (*Chelon labrosus*) in mesocosm with semi-extensive technology. *Aquaculture* 259,
418 190-201.

419 Bentzon-Tilia, M., Sonnenschein, E. C., Gram, L., 2016. Monitoring and managing microbes in
420 aquaculture—Towards a sustainable industry. *Microbial Biotechnology* 9, 576-584.

421 Bessey, O.A., Lowry, O.H., Brock, M.J., 1946. Rapid coloric method for determination of alkaline
422 phosphatase in five cubic millimeters of serum. *Journal of Biological Chemistry* 164, 321–329.

423 Bjornsdottir, R., Smaradottir, H., Sigurgisladottir, S., Gudmundsdottir, B.K., 2011. Effects of
424 different environmental shading on the cultivable bacterial community and survival of first feeding
425 Atlantic halibut larvae. *Icelandic Agricultural Sciences* 24, 33–41.

426 Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of
427 protein utilizing the principle of protein–dye binding. *Analytical Biochemistry* 72, 248–254.

428 Cahu, C.L., Zambonino Infante, J.L., 1995. Maturation of the pancreatic and intestinal digestive
429 function in sea bass (*Dicentrarchus labrax*): effect of weaning with different protein sources. *Fish*
430 *Physiology and Biochemistry* 14, 431–437.

431 Cahu, C.L., Zambonino Infante, J.L., Péres, A., Quazuguel, P., Le Gall, M.M., 1998. Algal addition
432 in sea bass (*Dicentrarchus labrax*) larvae rearing: Effect on digestive enzymes. *Aquaculture* 161,
433 479–489.

434 Confer, J.L., Howick, G.L., Corzett, M.H., Kramer, S.L., Fitzgibbon, S., Landesberg, R., 1978. Visual
435 predation by planktivorous. *Oikos* 31, 27–37.

436 Crane, R. K., Boge, G., Rigal, A., 1979. Isolation of brush border membranes in vesicular form from
437 the intestinal spiral valve of the small dogfish *Scyliorhinus canicula*. *Biochimica et Biophysica Acta*
438 554, 264–267.

439 Der Meeren, T., Mangor-Jensen, A., Pickova, J., 2007. The effect of green water and light intensity
440 on survival, growth and lipid composition in Atlantic cod (*Gadus morhua*) during intensive larval
441 rearing. *Aquaculture* 265, 206-217.

442 FAO. 2018. Cultured Aquatic Species Information Programme. *Mugil cephalus*. In: FAO Fisheries
443 and Aquaculture Department, Rome. http://www.fao.org/fishery/culturedspecies/Mugil_cephalus/en

444 Faulk, C.K., Holt, G.J., 2005. Advances in rearing cobia *Rachycentron canadum* larvae in
445 recirculating aquaculture systems: Live prey enrichment and green water culture. *Aquaculture* 249,
446 231–243.

447 Folch, J., Lees, M., Sloane Stanley, G.H., 1957. A simple method for the isolation and purification of
448 total lipids from animal tissues. *Journal of Biological Chemistry* 226, 497–509.

449 Gisbert, E., Giménez, G., Fernandez, I., Kotzamanis, Y., Estévez, A., 2009. Development of digestive
450 enzymes in common dentex, *Dentex dentex*, during early ontogeny. *Aquaculture* 287, 381–387.

451 Gisbert, E., Mozanzadeh, M.T., Kotzamanis, Y., Estévez, A., 2016. Weaning wild flathead grey
452 mullet (*Mugil cephalus*) fry with diets with different levels of fish meal substitution. *Aquaculture*
453 462, 92–100.

454 Gisbert, E., Nolasco, H., Solovyev, M., 2018. Towards the standardization of brush border
455 purification and intestinal alkaline phosphatase quantification in fish with notes on other digestive
456 enzymes. *Aquaculture* 487, 102-108.

457 Gisbert, E., Conklin, D.B., Piedrahita, R.H., 2004. Effects of delayed first feeding on the nutritional
458 condition and mortality of California halibut larvae. *Journal of Fish Biology* 64, 116–132.

459 Gregory, R.S., Northcote, T.G., 1993. Surface, planktonic, and benthic foraging by juvenile chinook
460 salmon (*Oncorhynchus tshawytscha*) in turbid laboratory conditions. *Canadian Journal of Fisheries
461 and Aquatic Sciences* 50, 233– 240.

462 Hemaiswarya, S., Raja, R., Ravi Kumar, R., Ganesan, V., Anbazhagan, C., 2011. Microalgae: a
463 sustainable feed source for aquaculture. *World Journal of Microbiology and Biotechnology* 27, 1737-
464 1746.

465 Henning, S.J., 1987. Functional development of the gastrointestinal tract. In: Johnson, L.R. (Ed.),
466 *Physiology of the Gastrointestinal Tract*. 2nd edition, Raven Press, New York, pp. 285–300.

467 Hjelmeland, K., Pedersen, B.H., Nilssen, E.M., 1988. Trypsin content in intestines of herring larvae,
468 *Clupea harengus*, ingesting inert polystyrene spheres or live crustacea prey. *Marine Biology* 98, 331-
469 335.

470 Holm, H., Hanssen, L.E., Krogdahl, A., Florholmen, J., 1988. High and low inhibitor soybean meals
471 affect human duodenal proteinase activity differently: in vivo comparison with bovine serum
472 albumin. *Journal of Nutrition* 118, 515–520.

473 Horn, M.H., 1989. Biology of marine herbivorous fishes. *Oceanography and Marine Biology. Annual*
474 *Review* 27, 167–272.

475 Iijima, N., Tanaka, S., Ota, Y., 1998. Purification and characterization of bile salt activated lipase
476 from the hepatopancreas of red sea bream, *Pagrus major*. *Fish Physiology and Biochemistry* 18, 59–
477 69.

478 Korzen, L., Abelson, A., Israel, A., 2016. Growth, protein and carbohydrate contents in *Ulva rigida*
479 and *Gracilaria bursa-pastoris* integrated with an offshore fish farm. *Journal of Applied Ecology* 28,
480 1835–1845.

481 Lazo, J., Holt, G., Arnold, C., 2000. Ontogeny of pancreatic enzymes in larval red drum *Sciaenops*
482 *ocellatus*. *Aquaculture Nutrition* 6, 183-192.

483 Métais, P., Bieth, J., 1968. Détermination de l' α -amylase par une microtechnique. *Annales De*
484 *Biologie Clinique* 26, 133-142.

485 Ma, H., Cahu, C., Zambonino, J., Yu, H., Duan, Q., Le Gall, M.M., Mai, K., 2005. Activities of
486 selected digestive enzymes during larval development of red drum (*Sciaenops ocellatus*).
487 *Aquaculture* 245, 239-248.

488 Naas, K.E., Naess, T., Harboe, T., 1992. Enhanced first-feeding of halibut larvae (*Hippoglossus*
489 *hippoglossus* L.) in green water. *Aquaculture* 105, 143-156.

490 Nicholson, J.A., Kim, Y.S., 1975. A one-step l-amino acid oxidase assay for intestinal peptide
491 hydrolase activity. *Analytical Biochemistry* 63, 110–117.

492 Oren, O.H., 1981. *Aquaculture of Grey Mulletts*. Cambridge University Press, Cambridge
493 507 pp.

494 Papandroulakis, N., Divanach, P., Kentouri, M., 2002. Enhanced biological performance of intensive
495 sea bream (*Sparus aurata*) larviculture in the presence of phytoplankton with long photophase.
496 *Aquaculture* 204, 45– 63.

497 Reitan, K.I., Rainuzzo, J.R., Oie, G., Olsen, Y., 1997. A review of the nutritional effects of algae in
498 marine fish larvae. *Aquaculture* 155, 207–221.

499 Rocha, R.J., Ribeiro, L., Costa, R., Dinis, M.T., 2008. Does the presence of microalgae influence fish
500 larvae prey capture? *Aquatic Research* 39, 362–369.

501 Rousseeuw, P.J., Leroy, A.M., 2003. *Robust regression and outlier detection*. Wiley Hoboken, p. 195.

502 Skiftesvik, A.B., Browman, H.I., St-Pierre, J.-F., 2003. Life in green water: the effect of microalgae
503 on the behavior of Atlantic cod (*Gadus morhua*) larvae. In: Browman, H.I., Skiftesvik, A.B. (Eds.),
504 *The Big Fish Bang*. Proceedings of the 26th Annual Larval Fish Conference, pp. 97-103.

505 Skjermo, J., Vadstein, O., 1993. The effect of microalgae on skin and gut bacterial flora of halibut
506 larvae. In: Reinertsen, H., Dahle, L.A., Jørgensen, L., Tvinnerein, K. Eds., *Proceedings from*
507 *International Conference on Fish Farming Technology*, Trondheim, Norway, August 1993, pp. 61–
508 67.

509 Solovyev, M.M., Kashinskaya, E.N., Izvekova, G.I., Gisbert, E., Glupov, V.V., 2015. Feeding habits
510 and ontogenic changes in digestive enzyme patterns in five freshwater teleosts. *Journal of Fish*
511 *Biology* 85, 1395-1412.

512 Solovyev, M., Gisbert, E., 2016. Influence of time, storage temperature and freeze/thaw cycles on the
513 activity of digestive enzymes from gilthead sea bream (*Sparus aurata*). *Fish Physiology and*
514 *Biochemistry* 42, 1383–1394.

515 Solovyev, M.M., Campoverde, C., Öztürk, S., Moreira, C., Diaz, M., Moyano, F.J., Estévez, A.,
516 Gisbert, E., 2016. Morphological and functional description of the development of the digestive
517 system in meagre (*Argyrosomus regius*): An integrative approach. *Aquaculture* 464, 381-391.

518 Stottrup, J.G., Gravningen, K., Norsker, N.H., 1995. The role of different algae in the growth and
519 survival of turbot larvae (*Scophthalmus maximus* L.) in intensive rearing systems. *ICES Marine*
520 *Sciences Symposium* 201, 173-186.

521 Tamaru, C.S., Murashige, R., Lee, C.S., 1994. The paradox of using background phytoplankton
522 during the larval culture of striped mullet, *Mugil cephalus* L. *Aquaculture* 119, 167–174.

523 Utne-Palm, A. C., 2002. Visual feeding of fish in a turbid environment: Physical and behavioural
524 aspects. *Marine Freshwater Behaviour and Physiology* 35, 111–128.

525 Van der Meeren, T., 1991. Algae as first food for cod larvae, *Gadus morhua* L.: filter feeding or
526 ingestion by accident? *Journal of Fish Biology*, 39, 225–237.

527 Van der Meeren, T., Mangor-Jensen, A., Pickova, J., 2007. The effect of green water and light
528 intensity on survival, growth and lipid composition in Atlantic cod (*Gadus morhua*) during intensive
529 larval rearing. *Aquaculture* 265, 206-217.

530 Whitfield, A.K., Panfili, J., Durand, J.D., 2012. A global review of the cosmopolitan flathead mullet
531 *Mugil cephalus* Linnaeus 1758 (Teleostei: Mugilidae), with emphasis on the biology, genetics,
532 ecology and fisheries aspects of this apparent species complex. *Reviews in Fish Biology and Fisheries*
533 22, 641–681.

534 Worthington, C.C. (1991). *Worthington Enzyme Related Biochemicals Manual*, 3rd ed., Freehold,
535 New Jersey, USA.

536 Yúfera, M., Darias, M.J., 2007. The onset of exogenous feeding in marine fish larvae. *Aquaculture*
537 268, 53-63.

538 Zambonino-Infante, J.L., Cahu, C., 1994. Development and response to a diet change of some
539 digestive enzymes in sea bass (*Dicentrarchus labrax*) larvae. *Fish Physiology and Biochemistry* 12,
540 399–408.

541 Zemke-White, W.L., Clements, K.D., 1999. Chlorophyte and rhodophyte starches as factors in diet
542 choice by marine herbivorous fish. *Journal of Experimental Marine Biology and Ecology* 240, 137–
543 149.

544 Zouiten, D., Khemis, I. Ben, Besbes, R., Cahu, C., 2008. Ontogeny of the digestive tract of thick
545 lipped grey mullet (*Chelon labrosus*) larvae reared in “mesocosms.” *Aquaculture* 279, 166–172.

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563 **Figure legends**

564 **Figure 1** Daily turbidity measurements (shortly before larval feeding) of the microalgae treatments
565 in (a) Experiment 1 and the (b) red and (c) white clay trials in Experiment 2. Different turbidity levels
566 (NTU) were significantly ($P<0.05$) different from each other in each experiment.

567 **Figure 2** The effect of microalgae turbidity treatments on larval (a) mastaxes consumed, (b) survival
568 at 51 dph and (c) dry weight (DW) at 15, 18 and 25 dph. Mastax consumption (5 dph) and percent
569 (%) survival (51 dph) values having different letter(s) were significantly ($P<0.05$) different.

570 **Figure 3** The effect of the control (C), Iso A (IA), Iso B (IB), Nanno A (NA), Nanno B (NB)
571 microalgae turbidity treatments on 25 dph larval (a) docosahexaenoic acid (DHA), eicosapentaenoic
572 acid (EPA) and arachidonic acid (ArA) levels (mg/g DW) as well as (b) saturated (SAT),
573 monounsaturated (MONO) and polyunsaturated fatty acid levels (mg/g DW). Values within an
574 essential fatty acid or fatty acid group having different letter(s) were significantly ($P<0.05$) different.

575 **Figure 4** The effect of the control (c), Iso A, Iso B, Nanno A, Nanno B microalgae turbidity treatments
576 on 23 dph larval (a) lipase and (b) alkaline proteases (c) trypsin and (d) leu-ala peptidase levels (U/
577 mg protein). Values within a turbidity treatment having different letter(s) were significantly ($P<0.05$)
578 different.

579 **Figure 5** The effect of the control, Iso A, Iso B, Nanno A, Nanno B microalgae turbidity treatments
580 on 23 dph larval (a) alkaline phosphatase (U/ mg protein), (b) alkaline phosphatase (P)/leu-ala ratio
581 (c) amylase (U/ mg protein) and (b) alkaline proteases/lipase ratio. Values within a turbidity treatment
582 having different letter(s) were significantly ($P<0.05$) different.

583 **Figure 6** The effect of the control, Iso A, Iso B, Nanno A, Nanno B microalgae turbidity treatments
584 on 23 dph larval (a) amylase/trypsin and (b) amylase/alkaline proteases ratio levels. Values within a
585 turbidity treatment having different letter(s) were significantly ($P<0.05$) different.

586 **Figure 7** The effect of the turbidity treatments; no algae (0), low turbidity *Isochrysis galbana* (ISO
587 A), high turbidity *Isochrysis galbana* spp. (ISO B), low *Nannochloropsis oculata* (NANNO A) and
588 high *Nannochloropsis oculata* (NANNO B) added to rearing tanks growing 2-23 dph grey mullet
589 larvae. PCA is based on the matrix of covariations that, in turn, was calculated on the specific activity
590 of all the studied (pancreatic and intestinal) digestive enzymes in 79 dph specimens.

591 **Figure 8** The effect of experiment 2 clay trials a (Nanno A, Nanno B and Clay B) and b (Nanno A2,
592 Nanno B2, Clay B2) on (a, b) mastax consumption, (c) larval survival (%) and (d) dry weight (DW).

593 Regression analysis of mastax consumption curves (selected from AIC analyses) demonstrated that
 594 the curve of the Nanno B treatment was significantly different ($P < 0.05$) than the curves of Nanno
 595 A(2) and Clay B(2) in trials a and b. Survival (after arcsine transformation) and dry weight (DW)
 596 values within a trial having different numbers of asterisks (*) or letters were significantly ($P < 0.05$)
 597 different.

598

599

600

601

602

603

604

605

606

607

608

609 **Tables**

610 Table 1 The microalgae treatments with their concentrations (cell mL^{-1}), designations and turbidities
 611 (NTU)

Treatments	Designation	Turbidity (NTU)
Control (no microalgae)	Control	0.26 ± 0.01^a
<i>Isochrysis galbana</i> A ($0.0144 \times 10^6 \text{ cell mL}^{-1}$)	Iso A	0.77 ± 0.01^b
<i>Nannochloropsis oculata</i> A ($0.2 \times 10^6 \text{ cells mL}^{-1}$)	Nanno A	0.75 ± 0.01^b
<i>Isochrysis galbana</i> A ($0.0288 \times 10^6 \text{ cell mL}^{-1}$)	Iso B	1.18 ± 0.02^c
<i>Nannochloropsis oculata</i> A ($0.4 \times 10^6 \text{ cells mL}^{-1}$)	Nanno B	1.20 ± 0.02^c

612

613

614 Table 2 Composition (% DW) of food used to feed the grey mullet at different stages of
615 development

Feed	Rotifers¹	Unenriched <i>Artemia</i>	Enriched <i>Artemia</i>	Caviar™ (Bernaqua, Belgium)**	Ranaan Dry Feed (RDF, Israel)*	<i>Ulva lactuca</i> (IOLR,Israel)
Days fed (dph)	1-23	15	16-24	25-50	50-79	25-50
Protein (%)	57.3	49.1	53.1	55	56	34
Lipid (%)	27.3	12.8	28.7	15	14	7.4
Carbohydrate (%)	12.5	15	10	8	1	56
Ash	1.5	5	2	12	14.8	2.6

616 *800 µm

617 ** Fed together with *U. lactuca* at a ratio of 1:1 (w/w)

618

619

620

621

622

623


624

625

626 **Figures**

627

628 (a)





629

630

(b)

(c)



631

632 **Figure 1**

633

634


635

636

637

638


639 (a)



640

641


(b)



642

643

(c)




644

645 **Figure 2**

646

647



648

649 **Figure 3**

650

651

652

653

654

655

656

657

658



659

660

661



662

663



664

665



666

667 **Figure 4**

668

669

670


671

672


673

674

675




(b)




676

677



(d)



678

679

Figure 5

680

681

682


683

684


685

686

(a)



(b)



687

688

689 **Figure 6**

690

691

692

693

694

695

696

697

698


699

700

701

702

703



704

705 **Figure 7**

706

707

708

709

710

711

712

713

714

715


716

717


718

719

720




721



722


723



724

(b)

(d)



725 **Figure 8**