



This document is a postprint version of an article published in *Aquaculture*© Elsevier after peer review. To access the final edited and published work see <https://doi.org/10.1016/j.aquaculture.2019.734854>

Document downloaded from:



1 **Effects of rearing density on growth, digestive conditions, welfare indicators and gut**
2 **bacterial community of gilthead sea bream (*Sparus aurata*, L. 1758) fed different**
3 **fishmeal and fish oil dietary levels**

4
5 Luca Parma^{a*}, Nicole Francesca Pelusio^a, Enric Gisbert^b, Maria Angeles Esteban^c,
6 Federica D'Amico^d, Matteo Soverini^d, Marco Candela^d, Francesco Dondi^a, Pier Paolo
7 Gatta^a, Alessio Bonaldo^a

8
9 ^aDepartment of Veterinary Medical Sciences, University of Bologna, Via Tolara di Sopra
10 50, 40064 Ozzano Emilia, Italy

11 ^bIRTA – Sant Carles de la Ràpita, Programa d'Aqüicultura, Crta. del Poble Nou km 5.5,
12 43540 Sant Carles de la Ràpita, Spain

13 ^cDepartment of Cell Biology and Histology, Faculty of Biology, Campus Regional de
14 Excelencia Internacional “Campus Mare Nostrum”, University of Murcia, 30100 Murcia,
15 Spain

16 ^dUnit of Microbial Ecology of Health, Department of Pharmacy and Biotechnology,
17 University of Bologna, Via Belmeloro 6, 40126 Bologna, Italy

18
19 **Corresponding author:* Luca Parma, Department of Veterinary Medical Sciences,
20 University of Bologna, Viale Vespucci 2, 47042 Cesenatico, FC, Italy. *Tel.:* +39 0547
21 338931; *Fax:* +39 0547 338941

22 E-mail address: luca.parma@unibo.it (L. Parma)

23

24

25 **Abstract**

26

27 In Mediterranean aquaculture, significant advances have been made towards a
28 reduction of marine-derived ingredients in aquafeed formulation, as well as in defining
29 the effect on how environmental factors such as rearing density interact with fish health.
30 Little research, however, has examined the interaction between rearing density and
31 dietary composition on main key performance indicators, physiological processes and gut
32 bacterial community. A study was undertaken, therefore to assess growth response,
33 digestive enzyme activity, humoral immunity on skin mucus, plasma biochemistry and
34 gut microbiota of gilthead sea bream (*Sparus aurata*, L. 1758) reared at high (HD, 36-44
35 kg m⁻³) and low (LD, 12-15 kg m⁻³) final stocking densities and fed high (FM30/FO15,
36 30% fishmeal FM, 15% fish oil, FO) and low (FM10/FO3; 10% FM and 3% FO) FM and
37 FO levels. Isonitrogenous and isolipidic extruded diets were fed to triplicate fish groups
38 (initial weight: 96.2 g) to overfeeding over 98 days. The densities tested had no major
39 effects on overall growth and feed efficiency of sea bream reared at high or low FM and
40 FO dietary level. However, HD seems to reduce feed intake compared to LD mainly in
41 fish fed FM30/FO15. Results of digestive enzyme activity indicated a comparable
42 digestive efficiency among rearing densities and within each dietary treatment even if
43 intestinal brush border enzymes appeared to be more influenced by stocking density
44 compared to gastric and pancreatic enzymes. Plasma parameters related to nutritional and
45 physiological conditions were not affected by rearing densities under both nutritional
46 conditions a similar observation was also achieved through the study of lysozyme,
47 protease, antiprotease and total protein determination in skin mucus, however; in this case
48 lysozyme was slightly reduced at HD. For the first time on this species, the effect of

49 rearing density on gut bacterial community was studied. Different response in relation to
50 dietary treatment under HD and LD were detected. Low FM-FO diet maintained steady
51 the biodiversity of the gut bacterial community between LD and HD conditions while fish
52 fed high FM-FO level showed a reduced biodiversity at HD. According to the results, it
53 seems feasible to rear gilthead sea bream at the on-growing phase at a density up to 36-
54 44 kg m⁻³ with low or high FM-FO diet without negatively affecting growth, feed
55 efficiency, welfare condition and gut bacterial community.

56

57 **Keywords**

58

59 Gilthead sea bream, rearing density, fishmeal and fish oil replacement, digestive
60 enzyme, humoral immunity on skin mucus, gut bacterial community.

61

62 **Introduction**

63

64 Despite the considerable advances addressing the study of nutritional requirements and
65 sustainable feed ingredients in fish, which have resulted in a deep knowledge about the
66 optimal composition of aquafeeds for Mediterranean fish species, technical performance
67 indicators such as growth, feed utilization and survival in Mediterranean aquaculture have
68 not improved over the last decade. The intensification of production systems and their
69 possible effects on stress and welfare or the less explored interaction between nutrition,
70 feeding management and suboptimal environmental conditions may have contributed to
71 this stagnation. Among stress factors, inadequate rearing density has been recognized as
72 a source of chronic stress in fish species which could affect physiological processes such

73 as osmoregulation or immune competence, mobilization of energy sources and alterations
74 in behaviour, which are generally translated into a decreased feed intake, reduced feed
75 efficiency and decreased growth performance (Ellis et al., 2002; Tort et al., 2011). In
76 gilthead sea bream (*Sparus aurata*), several studies have evaluated the effects of stocking
77 density on growth and fish health. In juveniles, Canario et al. (1998) found that growth
78 was negatively correlated to stocking density when fish were reared at a final stocking
79 density of 16.8 kg m⁻³ compared to 2.4 kg m⁻³, while Montero et al. (1999) did not find
80 an effect on growth and feed intake when specimens (22-85 g) were reared up to 40.8 kg
81 m⁻³, even if a negative effect on plasma and serum parameters were detected. More
82 recently high stocking density (final density 57 kg m⁻³) decreased growth performance,
83 feed intake and feed efficiency of gilthead sea bream (12-58 g) in comparison to lower
84 density 5-26 kg m⁻³ (Diógenes et al., 2019). In addition, in adult fish (272-425g) rearing
85 density was increased up to 20 kg m⁻³ without affecting physiological parameters and
86 growth, when oxygen level was maintained above 70% of the saturation level (Araujo-
87 Luna et al., 2018). Concerning the effect of rearing density on welfare in this species,
88 several studies have elucidated the effect on different physiological parameters, including
89 plasma parameters, neuroendocrine factors, skin mucus biomarkers, liver proteome,
90 carbohydrate metabolism of several tissues and behavioural studies (Montero et al., 1999;
91 Sangiao-Alvarellos et al., 2005; Mancera et al., 2008; Alves et al., 2010; Sánchez-Muros
92 et al., 2017; Guardiola et al., 2018; Skrzynska et al., 2018; Diógenes et al., 2019). Most
93 of those studies were conducted using standard diets and whether these density-associated
94 changes in performance and welfare are consistent when fish are fed current low fishmeal
95 (FM) and fish oil (FO) diets remains little investigated (Wong et al., 2013). In addition,
96 only a few studies in fish species have evaluated whether the interaction between stocking

97 density and diet composition may affect gut microbiota and none of these have been
98 evaluated in gilthead sea bream. The exposure to stress factors can impact the gut
99 microbiome community profile by altering the relative proportions of the main microbiota
100 phyla (Galley et al., 2014), while a recent study on blunt snout bream (*Megalobrama*
101 *amblycephala*) provided new evidence that the gut microbiome might be involved in the
102 response to crowding and consequently to the adaptation of fish to environmental
103 stressors (Du et al., 2019). The aim of the present study was to explore the effect of high
104 and low rearing density on growth, digestive enzyme activity, plasma biochemistry,
105 humoral immunity of skin mucus and gut microbiome structure during the on-growing of
106 gilthead sea bream fed low and high FM and FO dietary levels.

107

108 **Materials and methods**

109

110 *2.1 Experimental diets*

111

112 Ingredients and proximate composition of the experimental diets are presented in
113 Table 1. Two isonitrogenous (46% protein) and isolipidic (17% lipid) diets were
114 formulated to contain high and low FM and FO dietary levels (FM30/FO15 and
115 FM10/FO3; 30% FM, 15% FO and 10% FM and 3% FO, respectively). Diets were
116 formulated with FM and with a mixture of vegetable ingredients currently used for sea
117 bream in aquafeed (Parma et al., 2016). The diets were produced via extrusion (pellet size
118 = 4.0 mm) by SPAROS Lda (Portugal).

119

120 *2.2 Fish density and rearing*

121

122 The experiment was carried out at the Laboratory of Aquaculture, Department of
123 Veterinary Medical Sciences of the University of Bologna (Cesenatico, Italy). Gilthead
124 sea bream were obtained from the fish farm Cosa s.r.l (Orbello, GR) and adapted to the
125 laboratory facilities for 10 days before the beginning of the trial. Afterwards, two rearing
126 densities (low density and high density, LD and HD, respectively) were established by
127 randomly distributing 40 and 120 fish per tank ($96.2 \pm 2.1\text{g}$) in six 800L tanks
128 corresponding to an initial density of 4.8 and 14.4 kg m^{-3} , respectively (Table 2).

129 Each diet was administered to triplicate tanks at both rearing densities over 98 days.
130 Tanks were provided with natural seawater and connected to a closed recirculation system
131 (overall water volume: 15 m^{-3}). The rearing system consisted of a mechanical sand filter
132 (PTK 1200, Astralpool, Barcelona, Spain), ultraviolet lights (25 mJ cm^{-2} : $32 \text{ m}^{-3} \text{ h}^{-1}$,
133 Blaufish, Barcelona, Spain) and a biofilter (PTK 1200, Astralpool, Barcelona, Spain).
134 The water exchange rate within each tank was 100% every hour, while the overall water
135 renewal amount in the system was 5% daily. During the trial, the temperature was kept at
136 $24 \pm 1.0 \text{ }^\circ\text{C}$ and the photoperiod was maintained at 12 h light and 12 h dark by means of
137 artificial light. The oxygen level was kept constant ($8.0 \pm 1.0 \text{ mg L}^{-1}$) through a liquid
138 oxygen system regulated by a software programme (B&G Sinergia snc, Chioggia, Italy).
139 Ammonia (total ammonia nitrogen $\leq 0.1 \text{ mg L}^{-1}$) and nitrite ($\leq 0.2 \text{ mg L}^{-1}$) were daily
140 monitored spectrophotometrically (Spectroquant Nova 60, Merck, Lab business,
141 Darmstadt, Germany) while salinity (30 g L^{-1}) was measured by a salt refractometer (106
142 ATC). Sodium bicarbonate was added on a daily basis to keep pH constant at 7.8–8.0.
143 Fish were fed *ad libitum* twice a day (8:30, 16:30) for six days a week (one meal on
144 Sundays) via automatic feeders using an overfeeding approach with a daily feeding ration

145 10% higher than the daily ingested ration of the previous days as reported by Bonvini et
146 al. (2018a). Each meal lasted 1 h, after which the uneaten pellets of each tank were
147 collected, dried overnight at 105°C, and weighted for overall calculation.

148

149 *2.3 Sampling*

150

151 At the beginning and at the end of the experiment, all the fish in each tank were
152 anaesthetised by 2-phenoxyethanol at 300 mg L⁻¹ and individually weighed. The
153 proximate composition of the carcasses was determined at the beginning of the trial on a
154 pooled sample of 10 fish and on a pooled sample of 5 fish per tank at the end of the trial.

155 At the end of the trial, for the assessment of the specific activity of gastric (pepsin) and
156 pancreatic (trypsin, chymotrypsin, total alkaline proteases, α -amylase and bile salt-
157 activated lipase) digestive enzymes, 3 fish per tank (n = 9 fish per diet treatment) at 5
158 hours post meal (hpm) were randomly sampled, euthanized with overdose anaesthetic and
159 immediately eviscerated. The alimentary tract was dissected, adherent adipose and
160 connective tissues carefully removed and the gastrointestinal tract was stored at -80 °C
161 until their analysis. For the analysis of intestinal enzymes (alkaline phosphatase, maltase,
162 aminopeptidase-N and leucine-alanine peptidase), 3 fish per tank were sampled at 8 hpm,
163 at the same time, after fish dissection, anterior and posterior intestines were dissected and
164 stored at -80 °C until their analysis. Sampling times were selected in order to maximize
165 pancreatic enzyme levels in the stomach and anterior region of the intestine coinciding
166 with their maximal secretion into the gut from the exocrine pancreas due to the presence
167 of feed in the gut, while the activity of intestinal enzymes was measured at the end of the
168 digestion process (Deguara et al., 2013).The measurements of digestive enzymes was

169 then obtained by pooling the 3 fish sampled per tank during the analyses, as the tank was
170 considered as the experimental unit and not the organism. At the same time, digesta
171 content from posterior intestine (n = 15 fish per diet treatment, n = 5 fish per replicate)
172 was also individually sampled and immediately stored at -80°C for gut microbiota
173 analysis according to Parma et al. (2016).

174 For the assessment of plasma biochemistry, blood from 5 fish per tank (n=15 fish per
175 diet treatment) was collected from the caudal vein. Samples were then centrifuged (3000
176 x g, 10 min, 4°C) and plasma aliquots were stored at -80°C until analysis (Bonvini et al.,
177 2018b). Skin mucus samples were collected from 8 fish per tank according to the method
178 of Guardiola et al. (2014). Briefly, skin mucus was collected by gently scraping the
179 dorsolateral surface of specimens using a cell scraper, taking care to avoid contamination
180 with urino-genital and intestinal excretions. Collected mucus samples were then stored at
181 -80°C until analyses.

182 All experimental procedures were evaluated and approved by the Ethical-Scientific
183 Committee for Animal Experimentation of the University of Bologna, in accordance with
184 European directive 2010/63/UE on the protection of animals used for scientific purposes.

185

186 *2.4 Calculations*

187

188 The following formulae were used to calculate different performance parameters:
189 specific growth rate (SGR) ($\% \text{ day}^{-1}$) = $100 * (\ln \text{FBW} - \ln \text{IBW}) / \text{days}$ (where FBW and
190 IBW represent the final and the initial body weights, respectively). Feed Intake (FI) (g kg
191 $\text{ABW}^{-1} \text{ day}^{-1}$) = $((1000 * \text{total ingestion}) / (\text{ABW} * \text{days}))$ (where average body weight,
192 $\text{ABW} = (\text{IBW} + \text{FBW}) / 2$. Feed conversion ratio (FCR) = feed intake / weight gain. Protein

193 efficiency rate (PER) = (FBW – IBW) / protein intake. Gross protein efficiency (GPE)
194 (%) = 100 * [(% final body protein * FBW) - (% initial body protein * IBW)] / total
195 protein intake fish. Gross lipid efficiency (GLE) = 100 * [(final body lipid (%) * FBW) -
196 (initial body lipid (%) * IBW)] / total lipid intake fish. Lipid efficiency ratio (LER) =
197 [(FBW-IBW)/lipid intake].

198

199 *2.5 Proximate composition analysis*

200

201 Diets and whole body of sampled fish were analysed for proximate composition.
202 Moisture content was obtained by weight loss after drying samples in a stove at 105 °C
203 until a constant weight was achieved. Crude protein was determined as total nitrogen (N)
204 by using the Kjeldahl method and multiplying N by 6.25. Total lipids were determined
205 according to Bligh and Dyer's (1959) extraction method. Ash content was estimated by
206 incineration to a constant weight in a muffle oven at 450 °C. Gross energy was determined
207 by a calorimetric bomb (Adiabatic Calorimetric Bomb Parr 1261; PARR Instrument, IL,
208 U.S.A).

209

210 *2.6 Digestive enzyme activity*

211

212 Determination of pancreatic (α -amylase, bile salt-activated lipase, total alkaline
213 proteases), gastric (pepsin) and intestinal (alkaline phosphatase, aminopeptidase-N,
214 maltase and leucine-alanine peptidase) digestive enzymes were based on methods
215 previously described by Gisbert et al. (2009). In addition, spectrophotometric analyses
216 were performed as recommended by Solovyev and Gisbert (2016) in order to prevent

217 sample deterioration. In brief, the stomach and pyloric caeca samples (including 1 cm of
218 anterior intestine) were homogenized in 5 volumes (ww/v) of distilled water at 4 °C for
219 1 min followed by a sonication process of 30 sec. After a centrifugation (9,000 x g for 10
220 min at 4 °C), the supernatant was collected, aliquoted and stored at -20°C for the
221 quantification of gastric and pancreatic digestive enzymes.

222 Regarding intestinal enzymes, the anterior and posterior intestine samples were
223 homogenized in 30 volumes (w/v) of ice-cold Mannitol (50 mM), Tris-HCl buffer (2 mM)
224 pH 7.0, at a maximum speed for 30 s (IKA, Ultra-turrax®, USA), then 100 µL of 0.1M
225 CaCl₂ was added to the homogenate, stirred and centrifuged (9,000 x g for 10 min at 4
226 °C). A fraction of the supernatant was collected and stored at -20 °C for the leucine-
227 alanine peptidase (LAP) activity quantification. After a second centrifugation (3,400 x g
228 for 20 min at 4 °C), the supernatant was discarded, and the pellet containing the intestinal
229 brush border enzymes (alkaline phosphatase, aminopeptidase-N and maltase) dissolved
230 in 1 mL of Tris-Mannitol.

231 Total alkaline protease activity was measured using azocasein (0.5%) as substrate in
232 Tris-HCl 50 nmol L⁻¹ (pH = 9). One unit (U) of activity was defined as the nmoles of azo
233 dye released per minute and per mL of tissue homogenate, and the absorbance read at λ
234 = 366 nm. Trypsin activity was assayed using BAPNA (N-α-benzoyl-DL-arginine p-
235 nitroanilide) as substrate. One unit of trypsin per mL (U) was defined as 1 µmol BAPNA
236 hydrolyzed min⁻¹ mL⁻¹ of enzyme extract at λ = 407 nm (Holm et al., 1988).
237 Chymotrypsin activity was quantified using BTEE (benzoyl tyrosine ethyl ester) as
238 substrate and its activity (U) corresponded to the µmol BTEE hydrolyzed min⁻¹ mL⁻¹ of
239 enzyme extract at λ = 256 nm (Worthington, 1991). Alpha-amylase activity was
240 determined using 0.3% soluble starch as substrate (Métais and Bieth, 1968), and its

241 activity (U) was defined as the amount of starch (mg) hydrolysed during 30 min per mL
242 of tissue homogenate at $\lambda = 580$ nm. Bile salt-activated lipase activity was assayed for 30
243 min using p-nitrophenyl myristate as substrate. The reaction was stopped with a mixture
244 of acetone: n-heptane (5:2), the extract centrifuged (2 min at $6,080 \times g$ and 4°C) and the
245 increase in absorbance of the supernatant read at $\lambda = 405$ nm. Lipase activity (U) was
246 defined as the amount (nmol) of substrate hydrolyzed per min per mL of enzyme extract
247 (Iijima et al., 1998). Pepsin activity (U) was defined as the nmol of tyrosine liberated per
248 min per mL of tissue homogenate read at $\lambda = 280$ nm (Worthington, 1991).

249 Regarding intestinal digestive enzymes, alkaline phosphatase was quantified using 4-
250 nitrophenyl phosphate (PNPP) as substrate. One unit (U) was defined as $1 \mu\text{mol}$ of pNP
251 released $\text{min}^{-1} \text{mL}^{-1}$ of brush border homogenate at $\lambda = 407$ nm (Gisbert et al., 2018).
252 Aminopeptidase-N was determined using 80mM sodium phosphate buffer (pH = 7.0) and
253 L-leucine p-nitroanilide as substrate (in 0.1 mM DMSO) (Maroux et al., 1973). One unit
254 of enzyme activity (U) was defined as $1 \mu\text{g}$ nitroanilide released per min per mL of brush
255 border homogenate at $\lambda = 410$ nm. Maltase activity was determined using d(+)-maltose
256 as substrate in 100 mM sodium maleate buffer (pH = 6.0) (Dahkqvist, 1970). One unit of
257 maltase (U) was defined as μmol of glucose liberated per min per mL of homogenate at
258 $\lambda = 420$ nm. The assay of the cytosolic peptidase, LAP was performed on intestinal
259 homogenates applying the method described by Nicholson and Kim (1975) which utilized
260 L-alanine as substrate in 50 mM Tris-HCl buffer (pH = 8.0). One unit of enzyme activity
261 (U) was defined as 1 nmol of the hydrolyzed substrate $\text{min}^{-1} \text{mL}^{-1}$ of tissue homogenate
262 at $\lambda = 530$ nm. Soluble protein of crude enzyme extracts was quantified by means of the
263 Bradford's method (Bradford, 1976) using bovine serum albumin as standard. All
264 enzymatic activities were measured at $25\text{-}26^\circ\text{C}$ and expressed as specific activity defined

265 as units per mg of protein (U mg protein^{-1}). All the assays were made in triplicate
266 (methodological replicates) for each tank and the absorbance was read using a
267 spectrophotometer (TecanTM Infinite M200, Switzerland).

268

269 2.7 Humoral immunity on skin mucus

270 2.7.1. Lysozyme, protease, antiprotease and total protein determination

271

272 Lysozyme activity was measured according to the turbidimetric method described by
273 Swain et al. (2007). Briefly, 20 μL of skin mucus were placed in flat-bottomed 96-well
274 plates. To each well, 180 μL of freeze-dried *Micrococcus lysodeikticus* (0.2 mg mL^{-1} ,
275 Sigma-Aldrich) in 40 mM sodium phosphate (pH 6.2) was added as lysozyme substrate.
276 As blanks of each sample, 20 μL of skin mucus were added to 180 μL of sodium
277 phosphate buffer. The absorbance at $\lambda = 450 \text{ nm}$ was measured after 20 min at 35 °C in a
278 microplate reader (BMG Labtech). The amounts of lysozyme present in the samples were
279 obtained from a standard curve made with hen egg white lysozyme (HEWL, Sigma)
280 through serial dilutions in the above buffer. Skin mucus lysozyme values are expressed
281 as U mL^{-1} equivalent of HEWL activity.

282 Protease activity was quantified using the azocasein hydrolysis assay according to
283 Guardiola et al. (2014). Aliquots of 100 μL of each mucus sample were incubated with
284 100 μL of 100 mM ammonium bicarbonate buffer containing 0.7% azocasein (Sigma-
285 Aldrich) for 19 h at 30 °C. The reaction was stopped by adding 4.6% trichloro acetic acid
286 (TCA) and the mixture centrifuged ($10,000 \times g$, 10 min). The supernatants were
287 transferred to a 96-well plate in triplicate containing 100 $\mu\text{L well}^{-1}$ of 0.5 N NaOH. In
288 both cases, the OD was read at $\lambda = 450 \text{ nm}$ using a plate reader. Skin mucus was replaced

289 by trypsin (5 mg mL⁻¹, Sigma), as positive control (100% of protease activity), or by
290 buffer, as negative controls (0 % of protease activity).

291 Total antiprotease activity was determined in skin mucus by its ability to inhibit trypsin
292 activity (Hanif et al., 2004). Briefly, 10 µL of skin mucus were incubated (10 min, 22 °C)
293 with the same volume of standard trypsin solution (5 mg mL⁻¹) in a 96-well flat-bottomed
294 plate. After adding a volume of 100 µL of 100 mM ammonium bicarbonate buffer and
295 125 µL of buffer containing 2% azocasein (Sigma), samples were incubated (2 h, 30 °C)
296 and, following the addition of 250 µL 10% TCA, were incubated again (30 min, 30 °C).
297 The mixture was then centrifuged (10,000 x g, 10 min) and the supernatant was
298 transferred to a 96-well plate in triplicate, containing 100 µL well⁻¹ of 1 N NaOH before
299 the OD was read at $\lambda = 450$ nm using a plate reader. For a positive control, the reaction
300 buffer replaced mucus and trypsin, and for a negative control, the reaction buffer replaced
301 the mucus. The antiprotease activity was expressed in terms of the percentage of trypsin
302 inhibition according to the formula: % Trypsin inhibition = (Trypsin OD – Sample OD)/
303 Trypsin OD x 100.

304 Skin mucus protein concentration was determined by the dye binding method of
305 Bradford (1976) using bovine serum albumin (BSA, Sigma-Aldrich) as the standard.
306 Briefly, 2 mg mL⁻¹ solution of BSA was prepared and serial dilutions made with
307 phosphate buffer saline (PBS Sigma-Aldrich) as standards. Dilutions of 5 µL of skin
308 mucus and 15 µL of PBS were prepared. Then 250 µL of Bradford reagent (Sigma-
309 Aldrich) was added to BSA and skin mucus dilutions and incubated at room temperature
310 for 10 min. The absorbance of each sample was then read at $\lambda = 595$ nm and the results
311 were taken and plotted onto the standard curve to obtain the total protein content of skin

312 mucus. All spectrophotometry reads were conducted with a Varioskan 2.4.5, (Thermo
313 Scientific, MA, USA).

314

315 *2.8 Gut bacterial community DNA extraction and sequencing*

316

317 Total bacterial DNA was extracted and analysed from individual distal intestine
318 content obtained from 5 fish per tank as previously reported in Parma et al. (2019).
319 Afterwards, the V3–V4 hypervariable region of the 16S rRNA gene was amplified using
320 the 341F and 785R primers (Klindworth et al., 2013) with added Illumina adapter
321 overhang sequences and 2x KAPA HiFi HotStart ReadyMix (KAPA Biosystems).
322 Briefly, the thermal cycle consisted of an initial denaturation at 95 °C for 3 min, 30 cycles
323 of denaturation at 95 °C for 30 s, annealing at 55 °C for 30 s and extension at 72°C for
324 30 s, and a final extension step at 72 °C for 5 min. PCR reactions were cleaned up for
325 sequencing by using Agencourt AMPure XP magnetic beads as recommended in the
326 Illumina protocol “16S Metagenomic Sequencing Library Preparation” for the MiSeq
327 system, and as used in several other publications (Biagi et al., 2018; Soverini et al., 2016).
328 Sequencing was performed on Illumina MiSeq platform using a 2 x 250 bp paired-end
329 protocol according to the manufacturer’s instructions (Illumina, San Diego, CA). The
330 sequencing process resulted in a total of 1,553,593 high quality reads that were processed
331 using the QIIME 2 pipeline (Bolyen et al., 2019). After length (minimum/maximum =
332 250/550 bp) and quality filtering with default parameters, reads were cleaned using
333 DADA2 (Callahan et al., 2016) and clustered into OTUs at a 0.99 similarity threshold
334 using VSEARCH (Rognes et al., 2016). Assignment was carried out by using the RDP
335 classifier against Silva database (Quast et al., 2013).

336

337 *2.9 Metabolic parameters in plasma*

338

339 The levels of glucose (GLU), urea, creatine, uric acid, total bilirubin, bile acid,
340 amylase, lipase, cholesterol (CHOL), triglycerides (TRIG), total protein (TP), albumin
341 (ALB), aspartate aminotransferase (AST), alanine transaminase (ALT), alkaline
342 phosphatase (ALP), gamma-glutamyl transferase (GGT), creatine kinase (CK), lactate
343 dehydrogenase (LDH), calcium (Ca^{+2}), phosphorus (P), potassium (K^+) sodium (Na^+),
344 iron (Fe), chloride (Cl), magnesium (Mg), unsaturated iron binding capacity (UIBC), total
345 iron binding capacity (TIBC) and cortisol were determined in the plasma using samples
346 of 500 μL on an automated analyser (AU 400; Beckman Coulter) according to the
347 manufacturer's instructions. The ALB/globulin (GLOB), Na/K ratio and Ca x P were
348 calculated.

349

350 *2.10 Statistical analysis*

351

352 All data are presented as mean \pm standard deviation (SD). A tank was used as the
353 experimental unit for analysing growth performance and a pool of five and three sampled
354 fish were considered the experimental unit for analysing carcass composition and enzyme
355 activity respectively. Individual fish were used for analysing plasma biochemistry and
356 mucus stress parameters. Data of growth performance, nutritional indices, enzyme
357 activity, plasma and skin mucus parameters were analysed by a two-way analysis of
358 variance (ANOVA) and in case of significance ($p \leq 0.05$) Tukey's post hoc test was
359 performed. The normality and/or homogeneity of variance assumptions were validated

360 for all data preceding ANOVA. The R packages “Stats” and “Vegan” were used to
361 perform gut microbiota statistical analysis. In particular, to compare the microbiota
362 structure among different groups for alpha and beta-diversity, Wilcoxon rank-sum test
363 was used while the PCoA was tested using a permutation test with pseudo-F ratios
364 (function “Adonis” in the “Vegan” package). Alpha diversity of the different ecosystems
365 was computed using Hill numbers (Hill, 1973; Chao et al., 2014). Beta diversity was
366 estimated using both weighted and unweighted UniFrac metrics. Statistical analyses were
367 performed using GraphPad Prism 6.0 for Windows (Graph Pad Software, San Diego, CA,
368 USA) and RStudio interface for R (<https://www.r-project.org>). The differences among
369 treatments were considered significant at $p \leq 0.05$.

370

371 **3. Results**

372

373 *3.1 Growth*

374

375 Results on growth performance parameters are summarised in Table 2. No significant
376 effects on growth (FBW, weight gain and SGR) were detected between LD and HD
377 groups for both dietary treatments ($p > 0.05$). However, fish fed FM30/FO15 displayed
378 higher FBW, weight gain and SGR values compared to the FM10/FO3 group ($p < 0.05$).
379 Values of FI were lower in HD compared to LD (density effect $p = 0.002$) with more
380 marked differences in FM30/FO15 than FM10/FO3, whereas no significant diet effect on
381 FI was detected ($p > 0.05$). No significant effect of density on FCR was observed ($p >$
382 0.05), while the FM10/FO3 group showed higher FCR values, followed by FM30/FO15.
383 Survival rates were lower in the LD group ($p < 0.05$).

384 Data on body composition and nutritional indices are shown in Table 3. Whole body
385 composition values were not significantly influenced by different fish density ($p > 0.05$),
386 while lipid content was lower in fish fed the FM10/FO3 diet compared to the FM30/FO15
387 group ($p < 0.05$); however, ash and moisture levels were higher in FM10/FO3 than
388 FM30/FO15 fish ($p < 0.05$). No significant effects of fish density on PER, GPE, GLE and
389 LER were detected ($p > 0.05$); however, fish fed FM10/FO3 displayed lower PER, GPE,
390 GLE and LER compared to FM30/FO15 ($p < 0.05$).

391

392 *3.2 Digestive enzyme activity*

393

394 Data on specific activity of gastric, pancreatic and intestinal digestive enzymes are
395 shown in Table 4. The activities of both pancreatic (trypsin, chymotrypsin, total alkaline
396 proteases, amylase and bile salt-activated lipase) and gastric (pepsin) enzymes were not
397 significantly affected by the rearing density nor the diet ($p > 0.05$); with the exception of
398 trypsin, which was slightly affected by the diet composition ($p = 0.053$) with lower values
399 recorded in fish fed the FM10/FO3 diet compared to those fed the FM30/FO15 diet.
400 Regarding intestinal brush border enzymes measured in the anterior segment of the
401 intestine, aminopeptidase-N and maltase activities were not significantly affected by the
402 diet nor rearing density ($p > 0.05$), while phosphatase alkaline and LAP were slightly (p
403 < 0.1) lower in FM10/FO3 than FM30/FO15. The activity of LAP was significantly
404 higher at HD compared to LD for both dietary treatments ($p < 0.05$). Concerning the
405 intestinal enzymes measured in the posterior region of the intestine, aminopeptidase and
406 LAP were significantly affected by the rearing density with lower values recorded at HD
407 in comparison to those recorded in fish kept at LD ($p < 0.05$). Diet significantly affected

408 aminopeptidase-N and maltase activities which were significantly lower in sea bream fed
409 the FM10/FO3 diet ($p < 0.05$). No significant effects of both diets and tested densities
410 were detected in the phosphatase alkaline activities in the posterior intestine ($p > 0.05$).

411

412 *3.3 Plasma biochemistry*

413

414 The results of plasma parameters are shown in Table 5. No significant effect ($p > 0.05$)
415 of density on plasma parameters was detected under both feeding regimes. Concerning
416 the effect of diet on plasmatic parameters like urea, lipase, UIBC, A/G, TIBC, Na^+ , K^+ ,
417 Cl^- , these were higher in fish from the FM10/FO3 group compared to those from the
418 FM30/FO15 group ($p < 0.05$), while creatine, Ca^{2+} , Mg, CHOL, TP, ALB and Na^+/K^+
419 were lower in FM10/FO3 compared to FM30/FO15 fish ($p < 0.05$). No significant
420 differences related to density and feeding regimes for GLU, uric acid, creatine, total
421 bilirubin, AST, ALT, ALP, amylase, GGT, CK, LDH, P, TRIG, Bile acid, CaxP, Fe and
422 cortisol were detected among experimental groups ($p > 0.05$).

423

424 *3.4 Skin mucus non-specific immune biomarkers*

425

426 Results of skin mucus lysozyme, protease, antiprotease and total proteins are presented
427 in Figure 1 (A-D). Lysozyme activity was slightly affected by the rearing density (density
428 effect $p = 0.04$) with higher values recorded under LD rearing conditions. Specifically,
429 lysozyme was significantly higher in fish fed FM30/FO15 at LD rearing conditions
430 compared to those fed FM10/FO3 and reared at HD (Fig 1A; $p < 0.05$). Protease was
431 significantly reduced under fish fed FM10/FO3 (diet effect $p = 0.0006$), while no

432 significant effect of rearing density was detected ($p > 0.05$). Specifically, protease activity
433 in skin mucus was significantly higher in fish fed the FM30/FO15 diet at both rearing
434 densities compared to those fed FM10/FO3 and reared at LD (Fig 1B; $p < 0.05$). No
435 significant effect of density or diet were detected in antiprotease activity and total proteins
436 of skin mucus from fish belonging to the different experimental groups (Fig. 1, C-D; $p >$
437 0.05).

438

439 3.5 Gut bacterial community profiles

440

441 Taxonomic characterisation of the gut bacterial community at different phylogenetic
442 levels is represented in Figure 2: phylum in panel (A) and family in panel (B) and in
443 Supplementary Table 1. At phylum level, the most abundant taxa were Firmicutes,
444 Actinobacteria and Proteobacteria. In addition, the families most represented, all
445 belonging to Firmicutes phylum, were *Lactobacillaceae* (FM30/FO15_{HD}: 77.9% \pm
446 16.1%; FM30/FO15_{LD}: 86.5% \pm 4.4%; FM10/FO3_{HD}: 61.3% \pm 12.4%; FM10/FO3_{LD}:
447 67.6% \pm 12.2%), *Streptococcaceae* (FM30/FO15_{HD}: 2.0% \pm 1.5%; FM30/FO15_{LD}: 1.3%
448 \pm 1.4%; FM10/FO3_{HD}: 4.1 % \pm 3.7%; FM10/FO3_{LD}: 3.2% \pm 2.3%) and
449 *Staphylococcaceae* (FM30/FO15_{HD}: 1.4 % \pm 1.0 %; FM30/FO15_{LD}: 0.9 % \pm 0.4 %;
450 FM10/FO3_{HD}: 0.6% \pm 1.3%; FM10/FO3_{LD}: 0.3% \pm 0.5%). No significant differences
451 (Wilcoxon test $p > 0.05$, FDR correction) among groups at phylum level were detected
452 between specimens fed with the same diet but in different rearing density condition. On
453 the other hand, significant differences in several families such as *Staphylococcaceae* were
454 observed, values that were higher in the FM30/FO15_{HD} group than in FM10/FO3_{HD} group
455 ($p < 0.05$, Wilcoxon rank-sum test), and *Streptococcaceae*, higher in FM10/FO3_{HD} group

456 compared to FM30/FO15_{HD} group ($p < 0.05$). Moreover, at LD, both diets determined a
457 significant difference in the abundance of *Lactobacillaceae* and *Staphylococcaceae*, both
458 higher in FM30/FO15_{LD} group compared to FM10/FO3_{LD} ($p < 0.05$, Wilcoxon rank-sum
459 test) (Figure 2 C).

460 The biodiversity among microbiota from fish fed different diets and kept at different
461 stocking densities, expressed using Hill numbers of different magnitudes (from $q = 0$ to
462 $q = 2$), is represented in panel A of Figure 3. For all the q value magnitude, diet FM10/FO3
463 is characterised by a more even distribution of bacterial species characteristic that is
464 strengthened going from order $q = 0$ to order $q = 2$. According to the results, diet FM10/FO3
465 was more effective in the maintenance of a greater biodiversity in the sea bream gut
466 ecosystem. Furthermore, it is interesting to notice that for a $q = 0$, diet FM30/FO15
467 showed a number of species comparable to diet FM10/FO3, shifting to a significantly
468 more uneven ecosystem ($p < 0.05$, t-test) increasing the weight of the microbial core (q
469 values of 1 and 2, respectively). These results also showed that the response to rearing
470 conditions shifted depending on the fishes feeding regimen: diet FM10/FO3 maintained
471 steady the biodiversity of the gut microbiota between HD and LD (p value > 0.05 ; t-test).
472 On the other hand, diet FM30/FO15 was not able to maintain the evenness of the
473 community, as highlighted in the q value of 2, in which the FM30/FO15_{HD} group showed
474 a significantly reduced biodiversity when compared to the other groups (p value < 0.05 ,
475 t-test). To assess whether these different treatments could influence the gut bacterial
476 ecosystem, a multivariate analysis was performed. In both Principal Coordinates Analysis
477 (PCoA) graphs obtained using both weighted UniFrac metric (Figure 3 B) and
478 unweighted UniFrac metric (Figure 3 C) a significant separation was observed between

479 the different groups in the two-dimensional space (Adonis $p < 0.01$), except for
480 FM30/FO15_{HD} vs FM30/FO15_{LD} which did not show a significant ($p > 0.05$) separation.

481

482 **Discussion**

483

484 Several studies have investigated the effect of high rearing density on growth,
485 physiological responses and health in gilthead sea bream; however, studies concerning
486 the possible interaction between rearing density and low FM FO-based diets have been
487 less explored. In the present study, fish reared at high density (14.5-36/44 kg m⁻³, initial
488 and final density, respectively) within each FM and FO dietary levels showed similar
489 performance in terms of growth and feed utilisation in comparison to those reared at low
490 density (4.8-12/15 kg m⁻³). The results of the present study during the on-growing phase
491 (96-318g) go beyond the maximum density tested (20-31 kg m⁻³) by Araújo-Luna et al.
492 (2018) for gilthead sea bream at similar size (268-435 g). The authors did not find any
493 negative effects of high rearing density on SGR even if a significant linear relationship
494 between FCR and increasing stocking densities was observed. Indeed, the results of the
495 present study are consistent with a previous observation reported on juveniles (22-85 g)
496 in which high density up to 40.8 kg m⁻³ did not negatively affect growth (Montero et al.,
497 1999). However, more recently, Diogenes et al. (2019) found that rearing density up to
498 57 kg m⁻³ impaired FI, growth and FCR in sea bream juveniles (12-58g). The authors
499 suggested that 40 kg m⁻³ could be near the maximum tolerable stocking density for
500 gilthead sea bream of the weight range tested. This seems in agreement also for the size
501 tested in the present study; even if high density had no negative effect on the overall
502 growth and feed utilisation, high density significantly ($p = 0.002$) reduced FI.

503 Interestingly, this effect was mainly reported in high FM and FO dietary level and this
504 could be a consequence of the higher final stocking density obtained under this treatment
505 (44 vs 36 kg m⁻³, FM30/FO15, FM10/FM3, respectively) or be due to the fact that density
506 could have increased feeding competition only in a potentially more palatable and
507 digestible diet. The differences observed in growth performance between diets were
508 mainly related to a lower feed utilisation occurring in FM10/FO3; however it should be
509 taken into account that the growth performance achieved in the present trial under both
510 diets is in line with those found in literature for similar dietary formulation and that the
511 sole comparison between the two diets was not the purpose of the present study.

512 Stress conditions can disrupt the endocrine system and affect some physiological
513 functions such as digestive capacity (Trenzado et al., 2018). Few studies have evaluated
514 the effect of stocking density with a dietary interaction on digestive enzyme activity at
515 the on-growing stage in fish species (Wong et al., 2013). In the present study rearing
516 density did not affect pancreatic digestive enzyme specific activities under both dietary
517 treatments. Similarly, protease, lipase and amylase activities were not affected by rearing
518 density in gilthead sea bream fed increasing dietary tryptophan level with alternative
519 vegetable protein sources (Diogenes et al., 2019) or in Nile tilapia (*Oreochromis*
520 *niloticus*) fed dietary live and heat-inactive baker's yeast in vegetable-meal based diet
521 (Ran et al., 2016). Contrarily, Trenzado et al. (2018) studying the interaction between
522 stocking density and dietary lipid content in rainbow trout (*Oncorhynchus mykiss*) found
523 that stocking density inhibited the adaptive response of lipase activity and enhanced the
524 protease activity inhibition due to higher dietary lipid content. Compared to the pancreatic
525 enzyme activity, in the present study, density seemed to slightly affect the proteolytic
526 enzyme activity measured in the intestinal brush border of enterocytes. In particular, LAP

527 activity measured in the brush border of the anterior intestine tended to increase at high
528 density while aminopeptidase and LAP activity in the posterior intestine was slightly
529 reduced at high density in particular in the low FM-FO diet. The alkaline phosphatase of
530 the intestinal brush border is used as a marker of intestinal integrity and among its
531 functions was found to keep gastrointestinal inflammation under control (Lalles et al.,
532 2019, Messina et al., 2019). In addition, Nile tilapia reared at higher density displayed
533 higher alkaline phosphatase activity, possibly in line with higher pathogenic stressors at
534 high rearing density (Ran et al., 2016). In the present study, the absence of differences in
535 the alkaline phosphatase activity suggested no major functional changes in the integrity
536 of the intestine under different rearing density in both dietary treatments. The evaluation
537 of several plasma biochemical parameters is considered a valuable approach for assessing
538 the suitability of feeding practices, metabolic disorders, rearing conditions and presence
539 of acute or chronic stressors (Peres et al., 2013; Guardiola et al., 2018). No significant
540 effect of stocking density on any of the twenty-seven different plasma parameters
541 measured was detected under both dietary treatments. It is commonly accepted that high
542 stocking density generally leads to increased plasma cortisol levels in different fish
543 species, enhancing metabolic rate and compromising energy availability for several
544 physiological processes such as growth (Ashley, 2007). However, an opposite cortisol
545 response to stocking density has been also observed in some fish species suggesting that
546 cortisol response to stocking density is species-dependent and related to the gregarious
547 behaviour of the species at a specific stage of life (De las Heras et al., 2015; Millán-
548 Cubillo et al., 2016). Previous study of juveniles and adult sea bream held at high stocking
549 density, giving rise to chronic stress, showed significantly higher levels of plasma cortisol
550 than those held at low density, suggesting the incapacity of this species to reach adaptation

551 under chronic high rearing density conditions (Montero et al., 1999; Sangia-Alvarellos et
552 al., 2005). In accordance, TP, CHOL, TRIG were also found to be reduced at high
553 stocking density as a consequence of increased energy demand under stressful conditions
554 and possibly mediated by increased plasma cortisol (Diogenes et al., 2019). As also
555 reported for Senegal sole (*Solea senegalensis*) by Azeredo et al. (2019) the fact that fish
556 held at high density did not show higher plasma cortisol than their low-density
557 counterparts might be related to negative feedback mechanisms established in the HPI
558 axis, as a strategy of chronically stressed animals to attenuate an exacerbated stress
559 response (Bonga, 1997; Mommsen et al., 1999). In addition, the absence of effects of
560 rearing density on GLU, CHOL, TP and TRIG, suggests that the differences in rearing
561 density were not able to alter the metabolic processes related to growth and feed
562 utilisation. Non-specific plasma enzymes, such as AST, GGT, ALP, CK and LDH are
563 considered useful indicators of the health status and their elevated plasma level may
564 indicate specific tissue damage of several organs including liver, muscle, spleen and
565 kidney related to pathological processes, toxic chemical exposure, or traumatic conditions
566 or hypoxia, whereas specific references for this species and age are few (Peres et al.,
567 2013; Guardiola et al., 2018). Values of AST, CK, GGT and LDH were found in the lower
568 part of the range proposed by Peres et al. (2013) for healthy juvenile sea bream (70 g) fed
569 FM-based diet at low rearing density (3-5 kg m⁻³) and in line with those found by
570 Guardiola et al. (2018) during a feeding trial in sea bream of similar size. Levels of ALP
571 were higher than values previously found by Peres et al. (2013) and Guardiola et al.
572 (2018), a difference which can be related to FI since this enzyme is involved in the
573 absorption and transport of lipid and carbohydrates from the intestine, and its intestinal
574 activities are positively correlated with food ingestion and growth rate (Lemieux et al.,

1999; Lalles et al., 2019). The values of plasma electrolytes provided in the trial were comparable with the values reported in sea bream (Peres et al., 2013; Guardiola et al., 2018) and sobaity sea bream (*Sparidentex hasta*) (Hekmatpoure et al., 2019). Plasma electrolytes are indicators of the secondary phase of stress response in fish, providing an indirect indication of altered plasma cortisol levels; in particular plasma phosphorus and calcium levels were found to be sensitive to fish stocking density (Hrubec et al., 2000) while potassium levels are accepted as a general indicator of stress in fish (Guardiola et al., 2018).

Evaluation of skin mucosal immunity has been proposed recently as a promising alternative stress assessment in fish species after stressful conditions including crowding or transportation, whereas data of specific mucosal component in response to different stressors are still scarce (Guardiola et al 2016; Sanahuja et al., 2019). Enzymes in the epidermal mucus such as lysozyme, protease and antiprotease play an important role in humoral and skin mucus defence acting directly on a pathogen, or activating and enhancing the production of various immunological components of fish subjected to stressful situations (Esteban, 2012; Guardiola et al., 2016). The present results indicate different effects of treatments on specific skin mucus components, lysozyme being slightly reduced by high rearing density while protease was mainly reduced by low FM-FO diets. Both enzymes have been shown to be modulated either by diet or environmental conditions in sea bream. Most studies have shown the possibility of increasing lysozyme activity of skin mucus by dietary additives, such as selenium nanoparticles, *Moringa oleifera* leaves or probiotics; but crowding conditions at 20 kg m⁻³ for 30 days has also been reported to lead to an increase in lysozyme gene expression in sea bream skin mucus (Cordero et al., 2016; Mansour et al., 2018; Dawood et al., 2019). Concerning protease

599 activity, Guardiola et al. (2016) found a significant increase in this activity after 24 and
600 48 h of acute 50 kg m⁻³ crowding stress. However, in the same study a reduction in the
601 protease activity was also found after 48 h. The effect of protease activity under chronic
602 stressful conditions has been poorly investigated. Easy et al. (2010) studied the skin
603 mucus components following short- and long-term handling stress in Atlantic salmon
604 (*Salmo salar*), and no correspondence between skin mucus component and plasma
605 cortisol level in long-term stress was observed, suggesting that the activation of mucus
606 proteases may have been triggered by short-term elevated cortisol levels or that skin
607 mucus protease activation could result from physical disturbances such as abrasion due
608 to netting or overcrowding. More studies are needed to understand the role played by skin
609 mucus on stress in fishes.

610 Although the study of the gut microbiota by next-generation sequencing (NGS) has
611 already been conducted in this species under different feeding treatment, no information
612 concerning the effects of rearing density on gut microbiota is available. According to our
613 findings, the gut bacterial community is dominated by Firmicutes (69.9-92.2%), followed
614 by Actinobacteria and Proteobacteria. The dominance of Firmicutes we observed is in
615 general agreement with the previous NGS-based survey of the gut bacterial community
616 in sea bream and other marine or freshwater species fed similar aquafeed ingredients
617 employed in the present study (FM, soy-derivates, corn glutens, wheat gluten and wheat
618 meal) (Parma et al., 2016, Rimoldi et al., 2018a, 2018b; Parma et al., 2019). However,
619 our data differ from previous findings concerning the gut bacterial community of gilthead
620 sea bream and other Mediterranean fish species which displayed a dominance of
621 Proteobacteria and detected Firmicutes as the subdominant component (Carda-Diéguéz
622 et al., 2014; Gatesoupe et al., 2016, Piazzon et al., 2017). These works characterised the

623 mucosa-adherent gut microbiota, which could differ from the microbiota of the intestinal
624 lumen (Ringo et al., 2018). In this context, a recent comparison between mucosa-adherent
625 gut microbiota and intestinal lumen gut microbiota in sea bream highlighted the
626 dominance of Proteobacteria in the gut mucosa while Firmicutes dominated the intestinal
627 lumen in the same specimens (unpublished data). In addition, other studies revealed that
628 the differences in abundance between Firmicutes and Proteobacteria could also have been
629 related to the dietary composition. In rainbow trout, the presence of Proteobacteria was
630 favoured by an animal protein-based diet while the inclusion of at least 25% of plant
631 proteins in the diet favoured the presence of Firmicutes (Rimoldi et al., 2018b).

632 At the family level, the gut bacterial community of the present study was widely
633 dominated by *Lactobacillaceae* ranging from 61.3 to 86.5 %. The presence and the role
634 of *Lactobacillaceae* and other lactic acid bacteria (LAB) in fish species is still
635 controversial (Ringo et al., 2018). Several studies have associated a high LAB abundance
636 with a high inclusion level of dietary plant ingredients or functional additives in sea bream
637 (Parma et al., 2016; Rimoldi et al., 2018a) or other marine fish species (Apper et al., 2016;
638 Rimoldi et al., 2018b; Parma et al., 2019). However, some studies found a reduction in
639 LAB relative abundance when high FM replacement was also associated with a decrease
640 in performance (Estruch et al., 2015; Miao et al., 2018), while others found a higher
641 abundance of LAB in relation to vegetable protein associated with impaired gut health
642 (Gajardo et al., 2017). The results of the present study reinforce previous observation that
643 the dominance of *Lactobacillaceae* mainly *Lactobacillus* could be considered a valid
644 indicator of optimal gut health condition in sea bream.

645 No significant differences related to rearing density of any specific component within
646 each diet at phylum level were detected (Wilcoxon ran-sum test, $p > 0.05$, FDR

647 correction). However, different responses of the intestinal gut microbial composition in
648 relation to dietary treatment under high and low rearing density were detected as also
649 highlighted by weighted and unweighted UniFrac PCoA. In particular, no significant
650 separation was found between densities when fish were fed high FM-FO level, while
651 under low FM-FO diet density had a significant effect. Focusing on specific components
652 of the gut bacterial community, the results indicated that under high rearing density high
653 FM-FO level led to a significant increase in *Staphylococcaceae* and a reduction in
654 *Streptococcaceae* abundances compared to low FM-FO diet, while under low rearing
655 density *Lactobacillaceae* were less abundant in low FM-FO diet than high FM-FO diet.
656 Although no significant differences were detected, high rearing density seems to reduce
657 the amount of *Lactobacillaceae* (mainly *Lactobacillus spp*) within each dietary treatment
658 (Supplementary Table 1). No studies are available to compare the effect of rearing density
659 on specific gut microbial components in fish. In the present study, no evident signs of
660 stress induced by high rearing density were detected by results of performance, plasma
661 and skin mucus parameters; however, *Lactobacillaceae* may be highly sensitive in
662 relation to environmental stressors in fish and may deserve further attention for future
663 studies.

664 Analysis of biodiversity of the microbial community has highlighted a different
665 response to the feeding regimes, showing a general higher biodiversity in fish fed diets
666 containing higher vegetable ingredients. This is in general agreement with previous
667 findings detecting feeding habit as a key factor influencing fish gut microbial diversity
668 and observing an increasing trend in diversity following the order of carnivores,
669 omnivores and herbivores (Wang et al., 2018). In addition, a significant increase in α -
670 diversity indices at increasing FM replacement with vegetal ingredients was observed in

671 carnivorous fish species (Desai et al 2012; Miao et al., 2018). Concerning the interaction
672 between diet and rearing density, a low FM-FO diet maintained steady the biodiversity
673 of the ecosystem between low and high-density conditions while fish fed high FM-FO
674 level showed a significantly reduced biodiversity at high rearing density when compared
675 to the other groups. It has been suggested that in fish, reduction in diversity leads to
676 reduced competition for opportunistic or invading pathogens which may enter the
677 gastrointestinal tract of fish via feed or water (Apper et al., 2016). In several fish species,
678 α -diversity was not found to be affected by dietary vegetal ingredients (Apper et al., 2016;
679 Parma et al., 2016; Rimoldi et al., 2018b), by the interaction between diet and rearing
680 density (Wong et al., 2013) or by stocking density (Du et al., 2019). Also in pigs, stocking
681 density did not significantly affect biodiversity indices of gut microbiota (Li et al., 2017).
682 Interestingly, recent findings in the African cichlid *Astatotilapia burtoni* highlighted that
683 fish which experienced stressful conditions induced by subordinate social rank displayed
684 a reduced faecal microbial community α -diversity (Singh et al., 2019). Also in captive
685 mice and in wild red squirrels (*Tamiasciurus hudsonicus*) socially mediated stress
686 affected the intestinal microbiota leading to a reduction in microbial diversity and
687 richness (Bailey et al., 2011; Stothart et al., 2016). The reduction of biodiversity observed
688 in the present study only under the high FM-FO level could be correlated to increased
689 feeding competition only when a potentially more palatable high FM-FO diet is offered.
690 Another explanation may be associated with the lower feed intake observed under high
691 rearing density when fed high FM-FO level, or a combination of both factors: feeding
692 competition and feed intake. Recently, in perch (*Perca fluviatilis*) Zha et al. (2018) found
693 that gut microbial diversity responded to predation stress and food ration with a reduction
694 in diversity due to the presence of a predator and a reduced feed ration. The authors

695 suggested that a high ration of food favours bacteria that are quick colonizers and fast
696 growers while at lower food rations bacteria that are good competitors would be favoured.
697 In addition, the fact that in our study the reduction in gut microbial diversity was not
698 supported by evident altered physiological signs of stress could indicate a high sensitivity
699 of the gut microbial community structure to food competition, or to other social
700 interaction induced by rearing density. Thus, the analysis of gut microbial community
701 diversity could represent a valuable tool to assess social stress conditions for future
702 studies related to feeding behaviour and feeding competition.

703

704 **Conclusion**

705

706 In conclusion, the different rearing densities tested in this trial had no major effects on
707 overall performances and feed efficiency of gilthead sea bream reared at high or low fish
708 meal and fish oil dietary level. However, rearing density reduced feed intake in fish fed
709 high fish meal and fish oil dietary level. Results of digestive enzyme activities indicated
710 a comparable digestive efficiency among rearing densities and within dietary treatment
711 even if intestinal brush boarder enzymes such as LAP and aminopeptidase seems to be
712 more influenced by stocking density compared with other (gastric and pancreatic)
713 enzymes. Plasma parameters related to nutritional and physiological conditions were not
714 affected by rearing densities, indicating that sea bream can well cope with high rearing
715 density up to 36-44 kg m⁻³ and that a high level of vegetable dietary ingredients does not
716 amplify the potential stressful effects of rearing density. A similar observation was
717 achieved through the study of skin mucosal immunity; however in this case lysozyme
718 was slightly reduced at high density. For the first time the effect of rearing density on gut

719 bacterial community of this species was studied. Different responses in relation to dietary
720 treatment under high and low rearing density were detected. Low FM-FO diet maintained
721 steady the biodiversity of gut bacterial community between low and high rearing density
722 while fish fed high FM-FO level showed a significantly reduced biodiversity at high
723 rearing density possibly indicating higher social stress conditions related to feeding
724 competition under this treatment. According to the results, it seems feasible to rear
725 gilthead sea bream at the on-growing phase at a density up to 36-44 kg m⁻³ with low or
726 high FM-FO diet without negatively affecting growth, feed efficiency, welfare condition
727 and gut microbial community.

728

729 **Acknowledgment**

730

731 This research was supported by ERC (European Research Council) in MedAID project
732 (Mediterranean Aquaculture Integrated Development), Call H2020-SFS-2016-2017
733 (Sustainable Food Security – Resilient and resource-efficient value chains), Grant
734 Agreement n. 727315. Analyses of digestive enzymes conducted at IRTA were partially
735 supported by the project ADIPOQUIZ (RTI2018-095653-R-I00) funded by the
736 Ministerio de Ciencia, Innovación y Universidades (Spain). The authors would like to
737 thank Gillian Forlivesi Heywood for English language editing and Stefano Porcelli for
738 the technical contribution in fish rearing and laboratory analysis.

739

740

741

742

743 **References**

744

745 Alves, R.N., Cordeiro, O., Silva, T.S., Richard, N., de Vareilles, M., Marino, G., Di
746 Marco, P., Rodrigues, P.M., Conceição, L.E.C., 2010. Metabolic molecular
747 indicators of chronic stress in gilthead seabream (*Sparus aurata*) using
748 comparative proteomics. *Aquaculture*, 299, 1-4.

749 Apper, E., Weissman, D., Respondek, F., Guyonvarch, A., Baron, F., Boisot, P., Rodiles,
750 A., Merrifield, D.L., 2016. Hydrolysed wheat gluten as part of a diet based on
751 animal and plant proteins supports good growth performance of Asian seabass
752 (*Lates calcarifer*), without impairing intestinal morphology or microbiota.
753 *Aquaculture* 453, 40–48.

754 Araújo-Luna, R., Ribeiro, L., Bergheim, A., Pousão-Ferreira, P., 2018. The impact of
755 different rearing condition on gilthead seabream welfare: Dissolved oxygen levels
756 and stocking densities. *Aquac. Res.* 49 (12).

757 Ashley, P.J., 2007. Fish welfare: current issues in aquaculture. *Appl. Anim. Behav. Sci.*
758 104, 199–235.

759 Azeredo, R., Machado, M., Martos-Sitcha, J.A., Martínez-Rodríguez, G., Moura, J.,
760 Peres, H., Oliva-Teles, A., Afonso, A., Mancera, J.M., Costas, B. 2019. Dietary
761 tryptophan induces opposite health-related responses in the Senegalese sole (*Solea*
762 *senegalensis*) reared at low or high stocking densities with Implications in disease
763 resistance. *Front. Physiol.* art. no. 508.

764 Bailey, M.T., Dowd, S.E., Galley, J.D., Hufnagle, A.R., Allen, R.G., Lyte, M., 2011.
765 Exposure to a social stressor alters the structure of the intestinal microbiota:

766 Implications for stressor-induced immunomodulation. *Brain Behav. Immun.* 25
767 (3), 397-407.

768 Biagi, E., D'Amico, F., Soverini, M., Angelini, V., Barone, M., Turrone, S., Rampelli, S.,
769 Pari, S., Brigidi, P., Candela, M., 2018. Faecal bacterial communities from
770 Mediterranean loggerhead sea turtles (*Caretta caretta*). *Environ Microbiol Rep.*
771 doi: 10.1111/1758-2229.12683.

772 Bligh, E.G., Dyer, W.J., 1959. A rapid method of total lipid extraction and purification.
773 *Can. J. Biochem. Phys.* 37, 911–917.

774 Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C., Al-Ghalith, G.A.,
775 Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., et al., 2019. Reproducible,
776 interactive, scalable and extensible microbiome data science using QIIME 2. *Nat.*
777 *Biotechnol.* 37 (8), 852-857.

778 Bonga, S.E.W., 1997. The stress response in fish. *Physiol. Rev.* 77, 591–625.

779 Bonvini, E., Bonaldo, A., Parma, L., Mandrioli, L., Sirri, R., Grandi, M., Fontanillas, R.,
780 Viroli, C., Gatta, P.P., 2018a. Feeding European sea bass with increasing dietary
781 fibre levels: Impact on growth, blood biochemistry, gut histology, gut evacuation,
782 *Aquaculture*, 494, 1-9.

783 Bonvini, E., Bonaldo, A., Mandrioli, L., Sirri, R., Dondi, F., Bianco, C., Fontanillas, R.,
784 Mongile, F., Gatta, P.P., Parma, L., 2018b. Effects of feeding low fishmeal diets
785 with increasing soybean meal levels on growth, gut histology and plasma
786 biochemistry of sea bass. *Animal*, 12(5), 923-930.

787 Bradford, M., 1976. A rapid and sensitive method for the quantification of microgram
788 quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*
789 72, 248–254.

790 Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P.,
791 2016. DADA2: high-resolution sample inference from Illumina amplicon data.
792 Nat Methods 13, 581–583.

793 Canario, A.V.M., Condeça, J., Power, D.M., Ingleton, P.M., 1998. The effect of
794 stocking density on growth in the gilthead sea-bream, *Sparus aurata* (L.). Aquac.
795 Res. 29, 177–181.

796 Carda-Diéguez, M., Mira, A., Fouz, B., 2014. Pyrosequencing survey of intestinal
797 microbiota diversity in cultured sea bass (*Dicentrarchus labrax*) fed functional
798 diets. Fems Microbiol. Ecol. 87, 451–459.

799 Chao, A., N. J. Gotelli, T. Hsieh, E. L. Sander, K. Ma, R. K. Colwell, and A. M. Ellison.
800 2014. Rarefaction and extrapolation with Hill numbers: a framework for sampling
801 and estimation in species diversity studies. Ecological Monographs 84, 45–67.

802 Cordero, H., Morcillo, P., Cuesta, A., Brinchmann, M.F., Esteban, M.A., 2016.
803 Differential proteome profile of skin mucus of gilthead seabream (*Sparus aurata*)
804 after probiotic intake and/or overcrowding stress. J. Proteomics 132, 41-50.

805 Dahqvist, A., 1970. Assay of intestinal disaccharidase. Enzym. Biol. Clin. 11, 52–66.

806 Dawood, M.A.O., Koshio, S., Zaineldin, A.I., Van Doan, H., Moustafa, E.M., Abdel-
807 Daim, M.M., Angeles Esteban, M., Hassaan, M.S., 2019. Dietary supplementation
808 of selenium nanoparticles modulated systemic and mucosal immune status and
809 stress resistance of red sea bream (*Pagrus major*). Fish Physiol. Biochem., 45 (1),
810 219-230.

811 De las Heras, V., Martos-Sitcha, J.A., Yúfera, M., Mancera, J.M., Martínez-Rodríguez,
812 G., 2015. Influence of stocking density on growth, metabolism and stress of
813 thicklipped grey mullet (*Chelon labrosus*) juveniles. Aquaculture 448, 29–37.

814 Deguara, S., Jauncey, K., Agius, C., 2003. Enzyme activities and pH variations in the
815 digestive tract of gilthead sea bream. *J. Fish Biol.*, 62(5), 1033-1043.

816 Desai, A.R., Links, M.G., Collins, S.A., Mansfield, G.S., Drew, M.D., Van Kessel, A.G.,
817 Hill, J.E., 2012. Effects of plant-based diets on the distal gut microbiome of
818 rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 350, 134-142.

819 Diógenes, A.F., Teixeira, C., Almeida, E., Skrzynska, A., Costas, B., Oliva-Teles, A.,
820 Peres, H., 2019. Effects of dietary tryptophan and chronic stress in gilthead
821 seabream (*Sparus aurata*) juveniles fed corn distillers dried grains with solubles
822 (DDGS) based diets. *Aquaculture* 498, 396-404.

823 Du, F., Li, Y., Tang, Y., Su, S., Yu, J., Yu, F., Li, J., Li, H., Wang, M., Xu, P., 2019.
824 Response of the gut microbiome of *Megalobrama amblycephala* to crowding
825 stress. *Aquaculture* 500, 586-596.

826 Easy, R.H., Ross, N.W., 2010. Changes in Atlantic salmon *Salmo salar* mucus
827 components following short- and long-term handling stress. *J. Fish Biol.*, 77, 1616-
828 1631.

829 Ellis, T., North, B., Scott, A.P., Bromage, N.R., Porter, M., Gadd, D., 2002. Review
830 paper: the relationships between stocking density and welfare in farmed rainbow
831 trout. *J. Fish. Biol.* 61:493–531.

832 Esteban, M. A., 2012. An overview of the immunological defenses in fish skin, *ISRN*
833 *Immunol.* 1–29 <https://doi.org/10.5402/2012/853470>.

834 Estruch, G., Collado, M.C., Peñaranda, D.S., Tomás Vidal, A., Jover Cerdá, M., Pérez
835 Martínez, G., Martínez-Llorens, S., 2015. Impact of fishmeal replacement in diets
836 for gilthead sea bream *Sparus aurata* on the gastrointestinal microbiota
837 determined by pyrosequencing the 16S rRNA gene. *PLoS One* 10, e0136389.

838 Gajardo, K., Jaramillo-Torres, A., Kortner, T.M., Merrifield, D.L., Tinsley, J., Bakke,
839 A.M., Krogdahl, Å., 2017. Alternative protein sources in the diet modulate
840 microbiota and functionality in the distal intestine of Atlantic salmon (*Salmo*
841 *salar*). *Appl. Environ. Microbiol.* 83, e02615.

842 Galley, J.D., Nelson, M.C., Yu, Z., Dowd, S.E., Walter, J., Kumar, P.S., Lyte, M., Bailey,
843 M.T., 2014. Exposure to a social stressor disrupts the community structure of the
844 colonic mucosa-associated microbiota. *BMC Microbiol.* 14, 189.

845 Gatesoupe, F.J., Huelvan, C., Le Bayon, N., Le Delliou, H., Madec, L., Mouchel, O.,
846 Quazuguel, P., Mazurais, D., Zambonino-Infante, J., 2016. The highly variable
847 microbiota associated to intestinal mucosa correlates with growth and hypoxia
848 resistance of sea bass, *Dicentrarchus labrax*, submitted to different nutritional
849 histories. *BMC Microbiol.* 16, 266.

850 Gisbert, E., Giménez, G., Fernández, I., Kotzamanis, Y., Estevez, A., 2009. Development
851 of digestive enzymes in common dentex *Dentex dentex* during early ontogeny.
852 *Aquaculture* 287, 381–387.

853 Gisbert, E., Nolasco, H., Solovyev, M., 2018. Towards the standardization of brush
854 border purification and intestinal alkaline phosphatase quantification in fish with
855 notes on other digestive enzymes. *Aquaculture* 487, 102-108.

856 Guardiola, F.A., Cuesta, A., Arizcun, M., Meseguer, J., Esteban, M.A., 2014.
857 Comparative skin mucus and serum humoral defence mechanisms in the teleost
858 gilthead seabream (*Sparus aurata*). *Fish Shellfish Immunol.* 36, 545–551.

859 Guardiola, F.A., Cuesta, A., Esteban, M.A., 2016. Using skin mucus to evaluate stress in
860 gilthead seabream (*Sparus aurata* L.). *Fish Shellfish Immunol.* 59, 323-330.

861 Guardiola, F.A., Saraiva-Fraga, M., Cuesta, A., Esteban, M.A., 2018. Changes in natural
862 haemolytic complement activity induced by stress in gilthead seabream (*Sparus*
863 *aurata* L.). *Fish Shellfish Immunol.* 78, 317-321.

864 Hanif, A., Bakopoulos, V., Dimitriadis, G.J., 2004. Maternal transfer of humoral specific
865 and non-specific immune parameters to seabream (*Sparus aurata*) larvae. *Fish*
866 *Shellfish Immunol.* 17, 411-435.

867 Hekmatpour, F., Kochanian, P., Marammazi, J.G., Zakeri, M., Mousavi, S.-M., 2019.
868 Changes in serum biochemical parameters and digestive enzyme activity of
869 juvenile sobaity sea bream (*Sparidentex hasta*) in response to partial replacement
870 of dietary fish meal with poultry by-product meal. *Fish Physiol. Biochem.* 45,
871 599-611.

872 Hill, M. 1973. Diversity and evenness: A unifying notation and its consequences. *Ecology*
873 54, 427–432.

874 Holm, H., Hanssen, L.E., Krogdahl, A., Florholmen, J., 1988. High and low inhibitor
875 soybean meals affect human duodenal proteinase activity differently: in vivo
876 comparison with bovine serum albumin. *J. Nutr.* 118, 515–520.

877 Hrubec, T. C., Cardinale, J. L., Smith, S. A., 2000. Hematology and plasma chemistry
878 reference intervals for cultured tilapia (*Oreochromis hybrid*). *Vet. Clin. Pathol.*
879 29, 7–12.

880 Iijima, N., Tanaka, S., Ota, Y., 1998. Purification and characterization of bile-salt
881 activated lipase from the hepatopancreas of red sea bream, *Pagrus major*. *Fish*
882 *Physiol. Biochem.* 18, 59–69.

883 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glöckner, F.O.,
884 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical

885 and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* 41
886 (1), e1, <http://dx.doi.org/10.1093/nar/gks808>.

887 Lallès, J.P., 2019. Intestinal alkaline phosphatase in the gastrointestinal tract of fish:
888 biology, ontogeny, and environmental and nutritional modulation. *Rev. Aquac.*
889 DOI: 10.1111/raq.12340.

890 Li, L., Chen, S., Li, X., Wan, D., Liu, G., Yin, Y. L., 2017. Intestinal microbiota in
891 growing pigs: Effects of stocking density. *Food Agric. Immunol.*
892 doi:10.1080/09540105.2017.1409195

893 Lemieux, H., Blier, P., Dutil, J.D., 1999. Do digestive enzymes set a physiological limit
894 on growth rate and food conversion efficiency in the Atlantic cod (*Gadus*
895 *morhua*)? *Fish Physiol. Biochemist* 20, 293–303.

896 Mancera, J.M., Vargas-Chacoff, L., García-López, A., Kleszczyńska, A., Kalamarz, H.,
897 Martínez-Rodríguez, G., Kulczykowska, E., 2008. High density and food
898 deprivation affect arginine vasotocin, isotocin and melatonin in gilthead sea bream
899 (*Sparus auratus*). *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 149, 92–97.

900 Mansour, A.T., Miao, L., Espinosa, C., García-Beltrán, J.M., Ceballos Francisco, D.C.,
901 Esteban, M.Á., 2018. Effects of dietary inclusion of *Moringa oleifera* leaves on
902 growth and some systemic and mucosal immune parameters of seabream. *Fish*
903 *Physiol. Biochemist.* 44 (4), 1223-1240.

904 Maroux, S., Louvard, D., Baratti, J., 1973. The aminopeptidase from hog-intestinal brush
905 border. *Biochim. Biophys. Acta* 321, 282–295.

906 Messina, M., Bulfon, C., Beraldo, P., Tibaldi, E., Cardinaletti, G., 2019. Intestinal
907 morpho-physiology and innate immune status of European sea bass

908 (*Dicentrarchus labrax*) in response to diets including a blend of two marine
909 microalgae, *Tisochrysis lutea* and *Tetraselmis suecica*. *Aquaculture* 500, 660-669.

910 Métais, P., Bieth, J., 1968. Détermination de l' α -amylase par une microtechnique. *Ann.*
911 *Biol. Clin.* 26, 133–142.

912 Miao, S., Zhao, C., Zhu, J., Hu, J., Dong, X., Sun, L., 2018. Dietary soybean meal affects
913 intestinal homeostasis by altering the microbiota, morphology and inflammatory
914 cytokine gene expression in northern snakehead. *Sci. Rep.* 8, 113.
915 <https://doi.org/10.1038/s41598-017-18430-7>.

916 Millán-Cubillo, A. F., Martos-Sitcha, J. A., Ruiz-Jarabo, I., Cárdenas, S., Mancera, J. M.,
917 2016. Low stocking density negatively affects growth, metabolism and stress
918 pathways in juvenile specimens of meagre (*Argyrosomus regius*, Asso 1801).
919 *Aquaculture* 451, 87–92.

920 Mommsen, T. P., Vijayan, M. M., Moon, T. W., 1999. Cortisol in teleosts: dynamics,
921 mechanisms of action, and metabolic regulation. *Rev. Fish Biol. Fisher* 9, 211–
922 268. doi: 10.1023/A:1008924418720.

923 Montero, D., Izquierdo, M.S., Tort, L., Robaina, L., Vergara, J.M., 1999. High stocking
924 density produces crowding stress altering some physiological and biochemical
925 parameters in gilthead seabream, *Sparus aurata*, juveniles. *Fish Physiol.*
926 *Biochem.* 20, 53–60.

927 Nicholson, J.A., Kim, Y.S., 1975. A one-step L-amino acid oxidase assay for intestinal
928 peptide hydrolase activity. *Anal. Biochem.* 63, 110–117.

929 Parma, L., Candela, M., Soverini, M., Turrone, S., Consolandi, C., Brigidi, P., Mandrioli,
930 L., Sirri, R., Fontanillas, R., Gatta, P.P., Bonaldo, A., 2016. Next-generation
931 sequencing characterization of the gut bacterial community of gilthead sea bream

932 (*Sparus aurata*, L.) fed low fishmeal based diets with increasing soybean meal
933 levels. Anim. Feed Sci. Technol. 222, 204–216.

934 Parma, L., Yúfera, M., Navarro-Guillén, C., Moyano, F.J., Soverini, M., D'Amico, F.,
935 Candela, M., Fontanillas, R., Gatta, P.P., Bonaldo, A., 2019. Effects of calcium
936 carbonate inclusion in low fishmeal diets on growth, gastrointestinal pH, digestive
937 enzyme activity and gut bacterial community of European sea bass (*Dicentrarchus*
938 *labrax* L.) juveniles. Aquaculture, 510, 283-292.

939 Peres, H., Santos, S., Oliva-Teles, A., 2013. Selected plasma biochemistry parameters in
940 gilthead seabream (*Sparus aurata*) juveniles. J. Appl. Ichthyol. 29, 630–636.

941 Piazzon, M.C., Calduch-Giner, J.A., Fouz, B., Estensoro, I., Simó-Mirabet, P., Puyalto,
942 M., Karalazos, V., Palenzuela, O., Sitjà-Bobadilla, A., Pérez-Sánchez, J., 2017.
943 Under control: how a dietary additive can restore the gut microbiome and
944 proteomic profile, and improve disease resilience in a marine teleostean fish fed
945 vegetable diets. Microbiome 5:Article 164 DOI 10.1186/s40168-017-0390-3.

946 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J.,
947 Glöckner, F.O., 2013. The SILVA ribosomal RNA gene database project:
948 improved data processing and web-based tools. Nucleic Acid Res. 41, 590-596.

949 Ran, C., Huang, L., Hu, J., Tacon, P., He, S., Li, Z., Wang, Y., Liu, Z., Xu, L., Yang, Y.,
950 Zhou, Z., 2016. Effects of dietary live and heat-inactive baker's yeast on growth,
951 gut health, and disease resistance of Nile tilapia under high rearing density. Fish
952 Shellfish Immunol. 56, 263-271.

953 Rimoldi, S., Gliozheni, E., Ascione, C., Gini, E., Terova, G., 2018a. Effect of a specific
954 composition of short- and medium-chain fatty acid 1-Monoglycerides on growth

955 performances and gut microbiota of gilthead sea bream (*Sparus aurata*). PeerJ,
956 (7), art. no. e5355.

957 Rimoldi, S., Terova, G., Ascione, C., Giannico, R., Brambilla, F., 2018b. Next generation
958 sequencing for gut microbiome characterization in rainbow trout (*Oncorhynchus*
959 *mykiss*) fed animal by-product meals as an alternative to fish meal protein sources.
960 PLoS One 13, e0193652. <https://doi.org/10.1371/journal.pone.0193652>.

961 Ringo, E., Hossein, S., Ghosh, K., Doan, H.V., Beck, B.R., Song, S., 2018. Lactic acid
962 bacteria in finfish—an update. Front. Microbiol. 9, 1818.

963 Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: A versatile
964 open source tool for metagenomics. PeerJ, 10, art. no. e2584.

965 Sanahuja, I., Fernández-Alacid, L., Ordóñez-Grande, B., Sánchez-Nuño, S., Ramos, A.,
966 Araujo, R.M., Ibarz, A., 2019. Comparison of several non-specific skin mucus
967 immune defences in three piscine species of aquaculture interest. Fish Shellfish
968 Immunol. 89, 428-436.

969 Sánchez-Muros, M.J., Sánchez, B., Barroso, F.G., Toniolo, M., Trenzado, C.E., Sanz Rus,
970 A., 2017. Effects of rearing conditions on behavioural responses, social kinetics
971 and physiological parameters in gilthead sea bream *Sparus aurata*. Appl. Anim.
972 Behav. Sci. 197, 120-128.

973 Sangiao-Alvarellos, S., Guzmán, J.M., Laiz-Carrión, R., Míguez, J.M., Martín del Río,
974 M.P., Mancera, J.M., Soengas, J.L., 2005. Interactive effects of high stocking
975 density and food deprivation on carbohydrate metabolism in several tissues of
976 gilthead sea bream *Sparus auratus*. J. Exp. Zool. A. 303, 761–775.

977 Singh, A., Faber-Hammond, J.J., O'Rourke, C.F., Renn, S.C.P., 2019. Gut microbial
978 diversity increases with social rank in the African cichlid fish, *Astatotilapia*
979 *burtoni*, *Anim Behav* 152, 79-91.

980 Skrzynska, A.K., Martos-Sitcha, J.A., Martínez-Rodríguez, G., Mancera, J.M., 2018.
981 Unraveling vasotocinergic, isotocinergic and stress pathways after food
982 deprivation and high stocking density in the gilthead sea bream. *Comp. Biochem.*
983 *Physiol. A-Mol. Integr. Physiol.* 215, 35-44.

984 Solovyev, M., Gisbert, E., 2016. Influence of time, storage temperature and
985 freeze/thawcycles on the activity of digestive enzymes from gilthead sea bream
986 (*Sparus aurata*). *Fish Physiol. Biochem.* 42, 1383–1394.

987 Soverini, M., Quercia, S., Biancani, B., Furlati, S., Turrone, S., Biagi, E., Consolandi, C.,
988 Peano, C., Severgnini, M., Rampelli, S., Brigidi, P., Candela, M., 2016. The
989 bottlenose dolphin (*Tursiops truncatus*) faecal microbiota. *FEMS Microbiol Ecol.*
990 92, (4).

991 Stothart, M. R., Bobbie, C. B., Schulte-Hostedde, A. I., Boonstra, R., Palme, R.,
992 Mykytczuk, N. C. S., et al., 2016. Stress and the microbiome: Linking
993 glucocorticoids to bacterial community dynamics in wild red squirrels. *Biol. Lett.*
994 12(1), 20150875. <https://doi.org/10.1098/rsbl.2015.0875>.

995 Swain, N.S.P., Dash, S., Sahoo, P.K., Routray, P., Sahoo, S.K., Gupta, S.D., Meher, P.K.,
996 2007. Non-specific immune parameters of brood Indian major carp *Labeo rohita*
997 and their seasonal variations. *Fish Shellfish Immunol.* 22, 38–43.

998 Tort, L., 2011. Stress and immune modulation in fish. *Dev. Comp. Immunol.* 35, 1366–
999 1375.

1000 Trenzado, C.E., Carmona, R., Merino, R., García-Gallego, M., Furné, M. Domezain, A.
1001 Sanz, A., 2018. Effect of dietary lipid content and stocking density on digestive
1002 enzymes profile and intestinal histology of rainbow trout (*Oncorhynchus mykiss*),
1003 Aquaculture, 497, 10-16.

1004 Wang, A.n R., Ran, C., Ringø, E., Zhou, Z., 2018. Progress in fish gastrointestinal
1005 microbiota research. Rev. Aquacult. 10, 626–640.

1006 Wong, S., Waldrop, T., Summerfelt, S., Davidson, J., Barrows, F., Kenney, P.B., Welch,
1007 T., Wiens, G.D., Snekvi, K., Rawls, J., Good, C., 2013. Aquacultured rainbow
1008 trout (*Oncorhynchus mykiss*) possess a large core intestinal microbiota that is
1009 resistant to variation in diet and rearing density. Appl. Environ. Microbiol. 79
1010 (16), 4974-4984.

1011 Worthington Biochemical Corporation, 1991. Worthington Enzyme Manual: Enzymes,
1012 Enzyme Reagents, Related Biochemical. Worthington Biochemical Corp.,
1013 Freehold, New Jersey.

1014 Zha, Y., Eiler, A., Johansson, F., Svanbäck, R., 2018. Effects of predation stress and food
1015 ration on perch gut microbiota. Microbiome, 6 (1), art. no. 28.

1016
1017
1018
1019
1020
1021
1022
1023

Table 1. Ingredients and proximate composition of the experimental diets

	FM30/FO15	FM10/FO3
<i>Ingredients, % of the diet</i>		
Fish meal (LT70)	30.0	10.0
Soybean meal 48	9.0	9.0
Soy protein concentrate	10.0	20.5
Wheat gluten	5.0	10.2
Corn gluten	10.0	15.0
Wheat meal	9.7	7.3
Rapeseed meal	5.0	4.0
Sunflower meal	5.0	4.0
Fish oil	15.0	3.0
Rapeseed oil	0	13.0
Vit/Min premix ¹	1.0	1.0
Antioxidant powder (Paramega)	0.2	0.2
Sodium propionate	0.1	0.1
MCP		2.0
Lysine	-	0.3
Methionine	-	0.1
L-Tryptophan		0.3
<i>Proximate composition, % on a wet weight basis</i>		
Moisture	5.83	4.9
Protein	46.3	44.7
Lipid	17.2	17.8
Ash	8.2	6.4
Gross energy cal g ⁻¹	4945.7	4823.6

¹Vitamins and mineral premix (IU or mg kg⁻¹ diet; Invivo NSA, Portugal); DL-alpha tocopherol acetate, 200 mg; sodium menadione bisulphate, 10 mg; retinyl acetate, 16650 IU; DL-cholecalciferol, 2000 IU; thiamine, 25 mg; riboflavin, 25 mg; pyridoxine, 25 mg; cyanocobalamin, 0.1 mg; niacin, 150 mg; folic acid, 15 mg; L-ascorbic acid monophosphate, 750 mg; inositol, 500 mg; biotin, 0.75 mg; calcium panthotenate, 100 mg; choline chloride, 1000 mg; betaine, 500 mg; copper sulphate heptahydrate, 25 mg; ferric sulphate monohydrate, 100 mg; potassium iodide, 2 mg; manganese sulphate monohydrate, 100 mg; sodium selenite, 0.05 mg; zinc sulphate monohydrate, 200 mg
MCP: monocalcium phosphate

1025

1026

1027

1028

1029

1030

Table 2. Growth performance of gilthead sea bream reared at low and high stocking density and fed the experimental diets over 98 days.

Experimental diets

	FM30/FO15		FM10/FO3		Density	<i>P value</i>	
	LD	HD	LD	HD		Diet	Inter
Initial density kg m ⁻³	4.8±0.1 ^a	14.5±0.6 ^b	4.8±0.1 ^a	14.3±0.1 ^b	<0.0001	0.7078	0.7078
Final density kg m ⁻³	15.2±0.5 ^b	43.6±0.5 ^d	12.1±1.3 ^a	35.9±0.5 ^c	<0.0001	<0.0001	0.0011
IBW(g)	96.1±1.1	96.4±3.7	96.6±2.6	95.5±0.8	0.768	0.878	0.630
FBW(g)	317.8±5.6 ^b	292.5±3.9 ^b	253.1±27.2 ^a	246.2±2.8 ^a	0.084	0.0001	0.292
Weight gain (g)	221.7±5.4 ^b	196.2±0.5 ^b	156.5±25.3 ^a	150.7±3.0 ^a	0.071	0.0001	0.224
SGR	1.22±0.02 ^b	1.13±0.03 ^b	0.98±0.09 ^a	0.97±0.02 ^a	0.127	0.0001	0.248
FI	15.6±0.19 ^b	14.6±0.21 ^a	15.4±0.64 ^{ab}	14.5±0.03 ^a	0.002	0.506	0.818
FCR	1.43±0.02 ^a	1.42±0.01 ^a	1.70±0.21 ^b	1.61±0.02 ^{ab}	0.433	0.005	0.495
Survival %	95.8±1.4 ^a	99.4±0.5 ^b	95.8±1.4 ^a	97.2±0.5 ^{ab}	0.004	0.111	0.111

Data are given as the mean (n=3) ± SD. In each line, different superscript letters indicate significant differences among treatments ($P \leq 0.05$). FM30/FO15 = 300g kg⁻¹ fishmeal (FM), 150 g kg⁻¹ fish oil (FO); FM10/FO3 = 100g kg⁻¹ FM; 30g kg⁻¹ FO. LD, low rearing density; HD, high rearing density.

IBW = Initial body weight.

FBW = Final body weight.

SGR = Specific growth rate (% day⁻¹) = 100 * (ln FBW - ln IBW) / days.

ABW = average body weight = (IBW + FBW)/2.

FI= Feed intake (g kg ABW⁻¹ day⁻¹) = ((1000*total ingestion)/(ABW))/days).

FCR = feed conversion rate = feed intake (g) /weight gain (g)

1031

1032

Table 3. Body composition and nutritional indices of gilthead sea bream reared at low and high stocking density and fed the experimental diets over 98 days.

	Experimental diets				<i>P</i> -value		
	FM30/FO15		FM10/FO3		<i>Density</i>	<i>Diet</i>	<i>Inter.</i>
	LD	HD	LD	HD			
Whole body composition, %							
Protein	17.0 ± 0.5	17.2 ± 0.1	17.0 ± 0.0	16.9 ± 0.1	0.835	0.333	0.358
Lipid	21.4 ± 2.5 ^b	19.5 ± 1.5 ^{ab}	16.6 ± 0.7 ^a	17.0 ± 0.8 ^a	0.451	0.003	0.233
Ash	3.43 ± 0.11	3.57 ± 0.25	3.88 ± 0.08	3.83 ± 0.21	0.662	0.008	0.37
Moisture	58.0 ± 0.49	58.7 ± 0.7	59.5 ± 0.8	60.3 ± 0.9	0.206	0.024	0.949
Nutritional indices							
PER	1.51 ± 0.02	1.52 ± 0.01	1.32 ± 0.16	1.39 ± 0.02	0.443	0.009	0.567
GPE	25.8 ± 0.88	26.4 ± 0.38	22.6 ± 2.74	23.4 ± 0.20	0.455	0.006	0.879
GLE	101 ± 14.8 ^b	91.7 ± 9.0 ^b	60.9 ± 9.4 ^a	66.2 ± 4.6 ^a	0.768	0.000	0.253
LER	4.08 ± 0.05 ^b	4.11 ± 0.03 ^b	3.32 ± 0.40 ^a	3.48 ± 0.04 ^a	0.476	0.000	0.579

Data are given as the mean (n=3) ± SD. In each line, different superscript letters indicate significant differences among treatments ($p \leq 0.05$). FM30/FO15 = 300g kg⁻¹ fishmeal (FM), 150 g kg⁻¹ fish oil (FO); FM10/FO3 = 100g kg⁻¹ FM; 30g kg⁻¹ FO. LD, low rearing density; HD, high rearing density.

PER = Protein efficiency ratio = ((FBW-IBW)/protein intake).

GPE = Gross protein efficiency = 100*[(%final body protein*FBW) - (%initial body protein*IBW)]/total protein intake fish.

GLE = Gross lipid efficiency = 100*[(%final body lipid*FBW) - (%initial body lipid*IBW)]/total lipid intake fish.

LER = Lipid efficiency ratio = ((FBW-IBW)/lipid intake).

1033

1034

Table 4. Specific (U mg protein⁻¹) digestive enzyme activities of pancreatic (stomach and anterior intestine, AI) and intestinal brush border enzymes of gilthead sea bream reared at low (LD) and high (HD) stocking density and fed the experimental diets over 98 days.

	<i>Experimental diets</i>				<i>P-value</i>		
	FM30/FO15		FM10/FO3		<i>Density</i>	<i>Diet</i>	<i>Inter.</i>
	LD	HD	LD	HD			
<i>Pancreatic (Stomach/AI)</i>							
Pepsin	0.33 ± 0.11	0.34 ± 0.10	0.27 ± 0.18	0.55 ± 0.20	0.157	0.414	0.165
Trypsin	0.07 ± 0.03	0.04 ± 0.02	0.02 ± 0.02	0.03 ± 0.01	0.225	0.053	0.225
Chymotrypsin	0.60 ± 0.06	0.31 ± 0.17	0.34 ± 0.41	0.30 ± 0.20	0.276	0.366	0.413
Total alkaline proteases	0.56 ± 0.15	0.33 ± 0.15	0.25 ± 0.28	0.27 ± 0.13	0.333	0.119	0.270
Alpha-amylase	4.49 ± 1.47	3.38 ± 0.82	3.90 ± 3.24	2.37 ± 1.32	0.271	0.496	0.856
Bile salt activated lipase	0.017 ± 0.01	0.017 ± 0.01	0.022 ± 0.02	0.025 ± 0.01	0.784	0.264	0.819
<i>Brush border AI</i>							
Aminopeptidase-N	0.021 ± 0.01	0.022 ± 0.02	0.012 ± 0.01	0.008 ± 0.01	0.816	0.128	0.722
Phosphatase alkaline	1.83 ± 0.91	1.69 ± 0.31	1.10 ± 0.43	0.97 ± 0.09	0.701	0.075	0.981
Maltase	126.4 ± 25.8	124.1 ± 35.9	122.6 ± 36.9	64.9 ± 8.0	0.157	0.140	0.186
LAP	33.0 ± 3.1 ^{ab}	62.3 ± 18.7 ^b	24.7 ± 6.8 ^a	41.3 ± 4.8 ^{ab}	0.011	0.065	0.374
<i>Brush Border PI</i>							
Aminopeptidase	0.043 ± 0.01 ^b	0.026 ± 0.005 ^{ab}	0.0260 ± 0.005 ^{ab}	0.021 ± 0.005 ^a	0.031	0.031	0.169
Phosphatase alkaline	0.49 ± 0.10	0.94 ± 1.13	0.22 ± 0.08	0.13 ± 0.02	0.600	0.137	0.432
Maltase	130.5 ± 70.1	164.7 ± 62.9	64.8 ± 13.2	73.2 ± 26.1	0.524	0.042	0.700
LAP	46.6 ± 8.1 ^{ab}	45.9 ± 1.9 ^{ab}	55.6 ± 5.9 ^b	41.8 ± 0.9 ^a	0.038	0.430	0.058

Data are given as the mean (n = 3) ± SD. In each line, different superscript letters indicate significant differences among treatments ($p \leq 0.05$). FM30/FO15 = 300g kg⁻¹ fishmeal (FM), 150 g kg⁻¹ fish oil (FO); FM10/FO3 = 100g kg⁻¹ FM; 30g kg⁻¹ FO. LD, low rearing density; HD, high rearing density; AI, anterior intestine; PI posterior intestine; LAP, leucine-alanine peptidase.

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

Table 5. Plasma biochemistry values for sea bream kept under high (HD) and low (LD) rearing density and fed the experimental diets.

Parameters	<i>Experimental diets</i>						<i>P - value</i>	<i>Interaction</i>
	FM30/FO15		FM10/FO3		<i>Density</i>	<i>Diet</i>		
	LD	HD	LD	HD				
Glucose (mg dL ⁻¹)	119±26	123±29	117±31	101±24	0.374	0.079	0.145	
Urea (mg dL ⁻¹)	10.7±2.0 ^{ab}	9.25±1.44 ^a	11.6±2.1 ^{bc}	13.5±2.8 ^c	0.760	0.000	0.003	
Creatine (mg dL ⁻¹)	0.37±0.14 ^b	0.30±0.10 ^b	0.22±0.04 ^a	0.21±0.04 ^a	0.169	0.000	0.090	
Uric acid (mg dL ⁻¹)	0.51±0.40	0.39±0.25	0.42±0.42	0.32±0.30	0.206	0.361	0.868	
Tot bil (mg dL ⁻¹)	0.02±0.02	0.03±0.01	0.04±0.03	0.07±0.13	0.368	0.063	0.606	
Bil. Ac. (µmol dL ⁻¹)	69.3±39.7	64.8±41.7	48.9±30.4	61.2±40.8	0.685	0.215	0.381	
Amylase (U L ⁻¹)	2.88±5.35	0.88±0.34	1.25±1.00	1.50±2.12	0.226	0.488	0.121	
Lipase (U L ⁻¹)	2.20±2.43 ^a	1.69±1.74 ^a	4.13±2.92 ^{ab}	5.22±3.62 ^b	0.602	0.000	0.289	
CHOL (mg dL ⁻¹)	311±75 ^b	287±71 ^b	195±27 ^a	171±35 ^a	0.089	0.000	0.987	
TRIG (mg dL ⁻¹)	792±276	793±374	810±241	830±327	0.892	0.720	0.903	
TP (mg dL ⁻¹)	4.26±0.76 ^b	4.10±0.71 ^{ab}	3.78±0.29 ^{ab}	3.59±0.41 ^a	0.213	0.001	0.909	
ALB (g dL ⁻¹)	0.97±0.19 ^b	0.90±0.15 ^{ab}	0.89±0.06 ^{ab}	0.84±0.10 ^a	0.081	0.040	0.724	
AST(U L ⁻¹)	49.2±31.1	43.0±32.4	55.5±40.8	53.3±26.3	0.606	0.310	0.808	
ALT (U L ⁻¹)	1.81±1.76	1.31±0.60	1.19±0.54	1.11±0.32	0.232	0.088	0.378	
ALP (U L ⁻¹)	493±190	555±265	597±259	594±274	0.632	0.251	0.601	
CK (U L ⁻¹)	226±295	118±66	112±91	117±89	0.204	0.155	0.159	
GGT (U L ⁻¹)	0.10±0.00	0.10±0.00	0.10±0.00	0.10±0.0	1.000	1.000	1.000	
LDH (U L ⁻¹)	519±662	406±409	530±646	719±527	0.792	0.259	0.292	
Ca ⁺² (mg dL ⁻¹)	15.0±1.7 ^b	14.7±1.2 ^{ab}	14.3±0.7 ^{ab}	13.8±0.9 ^a	0.142	0.008	0.670	
P (mg dL ⁻¹)	13.3±2.1	12.0±1.8	12.2±1.4	12.3±2.4	0.249	0.381	0.183	
K ⁺ (mEq L ⁻¹)	7.16±2.45 ^b	5.28±1.58 ^a	7.06±1.70 ^{ab}	8.33±2.0 ^b	0.530	0.003	0.002	
Na ⁺ (mEq L ⁻¹)	188±6 ^a	189±5 ^{ab}	194±6 ^b	191±5 ^{ab}	0.566	0.005	0.094	
Fe (µg dL ⁻¹)	135±33	111±28	124±30	127±37	0.206	0.766	0.090	
Cl (mEq L ⁻¹)	148±4 ^a	150±4 ^a	157±5 ^b	156±4 ^b	0.325	0.000	0.131	
Mg (mg dL ⁻¹)	4.97±0.98 ^b	4.30±0.78 ^{ab}	3.86±0.50 ^a	3.86±0.72 ^a	0.078	0.000	0.073	
UIBC (µg dL ⁻¹)	464±78	433±97	502±68	488±96	0.300	0.031	0.695	
TIBC (µg dL ⁻¹)	599±97	544±116	626±74	616±105	0.193	0.049	0.373	
Cortisol (µg dL ⁻¹)	3.11±1.74	3.78±2.87	4.45±3.26	4.25±3.99	0.837	0.244	0.278	
ALB/GLOB	0.30±0.03 ^{ab}	0.28±0.02 ^a	0.31±0.02 ^b	0.31±0.02 ^b	0.174	0.002	0.158	
CaxP	201±50	178±39	175±24	169±36	0.138	0.068	0.366	
Na/K	28.9±8.8 ^a	38.8±10.7 ^b	29.1±7.5 ^a	24.1±6.0 ^a	0.243	0.001	0.001	

Data are given as the mean (n=15) ± SD. Different letters indicate significant difference ($P \leq 0.05$) between treatments. FM30/FO15 = 300g kg⁻¹ fishmeal (FM), 150 g kg⁻¹ fish oil (FO); FM10/FO3 = 100g kg⁻¹ FM; 30g kg⁻¹ FO. LD, low rearing density; HD, high rearing density. Tot Bil, total bilirubin; CHOL, cholesterol; TRIG, triglycerides; TP, total protein; ALB, albumin; AST, aspartate aminotransferase; ALT, alanine transaminase; ALP, alkaline phosphatase; GGT, gamma-glutamyl transferase; CK, creatine kinase; LDH, lactate dehydrogenase, Ca⁺², calcium; P, inorganic phosphorus; K⁺, potassium; Na⁺, sodium; Fe, iron; Cl, chloride; Mg, magnesium; UIBC, unsaturated iron binding capacity; TIBC, total iron binding capacity; GLOB, globuline.

1045
1046
1047
1048

1049

1050 **Key to Figures**


1051 Figure 1. A, Lysozyme (U mL^{-1}); B, protease activity (%); C, antiprotease activity (%);
1052 D, total protein (mg mL^{-1}) in skin mucus of gilthead seabream reared at low (LD, light
1053 grey) and high (HD, dark grey) stocking density and fed the experimental diets over 98
1054 days. FM30/FO15 = 300g kg^{-1} fishmeal (FM), 150 g kg^{-1} fish oil (FO); FM10/FO3 =
1055 100g kg^{-1} FM; 30g kg^{-1} FO. Data represent the mean \pm S.D. (N=24). Different letters
1056 denote significant differences between experimental groups ($p < 0.05$).

1057 Figure 2. Barplots representing the sea bream gut bacterial community at two
1058 phylogenetic levels: A) phylum; B) Family. In panel C) are reported the boxplots with
1059 the families showing a significant difference in relative abundance among groups (p value
1060 < 0.05 , Wilcoxon ran-sum test; FDR correction). FM30/FO15 = 300g kg^{-1} fishmeal (FM),
1061 150 g kg^{-1} fish oil (FO); FM10/FO3 = 100g kg^{-1} FM; 30g kg^{-1} FO. LD, low rearing
1062 density; HD, high rearing density.

1063 Figure 3. Internal biodiversity of sea bream gut microbiota in both feeding regimen and
1064 rearing densities computed using Hill numbers (A) highlighted a significant difference
1065 between diets ($p < 0.05$; Wilcoxon ran-sum test). Principal Coordinates Analysis
1066 (PCoA) plots obtained using weighted (B) and unweighted UniFrac (C) showing a
1067 significant difference among groups ($p < 0.01$; except FM30/FO15_{HD} vs FM30/FO15_{LD},
1068 $p > 0.05$; permutation test with pseudo-F ratios, Adonis). FM30/FO15 = 300g kg^{-1}
1069 fishmeal (FM), 150 g kg^{-1} fish oil (FO); FM10/FO3 = 100g kg^{-1} FM; 30g kg^{-1} FO. LD,
1070 low rearing density; HD, high rearing density.

1071

1072 Figure 1
1073




1074

1075

1076

1077


1078 Figure 2



1079

1080

1081 Figure 3



1082

1083

Supplementary Table 1.

Mean relative abundance (%) \pm SD (n=15) of bacterial phyla, classes, orders, families and genera detected in the distal intestine content of gilthead sea bream fed different diets under high and low rearing density. FM30/FO15 = 300g kg⁻¹ fishmeal (FM), 150 g kg⁻¹ fish oil (FO); FM10/FO3 = 100g kg⁻¹ FM; 30g kg⁻¹ FO. LD, low rearing density; HD, high rearing density. Only taxa with mean relative abundance \geq 0.1% in at least 1 treatment were included.

Diet	FM30/FO15_{HD}		FM30/FO15_{LD}		FM10/FO3_{HD}		FM10/FO3_{LD}	
Phylum	Mean	SD	Mean	SD	Mean	SD	Mean	SD
<i>Actinobacteria</i>	6.7	6.0	5.0	3.0	12.5	7.1	7.8	8.9
<i>Bacteroidetes</i>	1.4	4.3	0.2	0.3	0.5	0.4	0.9	0.7
<i>Chlamydiae</i>	0.1	0.2	0.4	0.4	1.1	1.0	1.6	1.6
<i>Chloroflexi</i>	0.0	0.0	0.0	0.1	0.1	0.2	0.1	0.2
<i>Cyanobacteria</i>	1.5	2.8	0.2	0.5	5.4	4.0	1.9	2.5
<i>Firmicutes</i>	87.3	9.4	92.2	4.3	69.9	13.4	77.9	13.7
<i>Gracilibacteria</i>	0.0	0.0	0.1	0.3	0.2	0.9	0.0	0.0
<i>Lentisphaerae</i>	0.1	0.2	0.0	0.0	0.0	0.0	0.0	0.0
<i>Planctomycetes</i>	0.1	0.3	0.3	0.2	0.8	0.9	1.4	1.7
<i>Proteobacteria</i>	2.5	2.9	1.2	0.9	7.6	6.3	7.1	6.1
<i>Saccharibacteria</i>	0.3	0.5	0.2	0.4	0.6	1.3	0.8	1.1
<i>Spirochaetae</i>	0.1	0.4	0.0	0.0	0.5	1.7	0.0	0.0
<i>TM6 (Dependentiae)</i>	0.0	0.0	0.0	0.0	0.1	0.3	0.1	0.2
<i>Verrucomicrobia</i>	0.0	0.1	0.0	0.2	0.1	0.3	0.0	0.0
<i>WS6</i>	0.0	0.0	0.0	0.1	0.2	0.7	0.0	0.1
<i>Unassigned;Other</i>	0.0	0.1	0.1	0.2	0.2	0.8	0.2	0.4
Class								
<i>Acidimicrobiia</i>	0.0	0.0	0.1	0.1	0.5	0.9	0.9	1.6
<i>Actinobacteria</i>	6.0	5.6	4.9	3.0	11.6	7.1	6.7	7.7
<i>Coriobacteriia</i>	0.4	0.9	0.0	0.1	0.4	0.8	0.1	0.3
<i>Thermoleophilia</i>	0.3	1.0	0.0	0.0	0.1	0.2	0.0	0.0
<i>Bacteroidia</i>	1.3	4.3	0.1	0.2	0.1	0.1	0.4	0.5
<i>Flavobacteriia</i>	0.1	0.2	0.1	0.1	0.3	0.2	0.4	0.5
<i>Sphingobacteriia</i>	0.0	0.0	0.0	0.0	0.2	0.3	0.2	0.2
<i>Chlamydiae</i>	0.1	0.2	0.4	0.4	1.1	1.0	1.6	1.6
<i>Chloroflexi;KD4-96</i>	0.0	0.0	0.0	0.1	0.1	0.2	0.1	0.2
<i>Chloroplast</i>	1.5	2.8	0.2	0.5	5.4	4.0	1.9	2.5
<i>Bacilli</i>	83.6	16.2	91.1	4.4	68.2	13.1	75.5	13.4
<i>Clostridia</i>	3.2	7.0	1.0	0.5	1.5	1.6	2.1	1.2
<i>Erysipelotrichia</i>	0.2	0.6	0.0	0.0	0.0	0.0	0.1	0.1
<i>Negativicutes</i>	0.3	0.8	0.0	0.1	0.2	0.3	0.3	0.4
<i>Gracilibacteria;Other</i>	0.0	0.0	0.1	0.3	0.2	0.9	0.0	0.0
<i>Planctomycetacia</i>	0.1	0.3	0.3	0.2	0.8	0.9	1.4	1.7
<i>Alphaproteobacteria</i>	0.6	1.1	0.3	0.3	2.5	2.3	2.0	2.6

<i>Betaproteobacteria</i>	0.2	0.4	0.1	0.1	0.5	0.7	0.1	0.2
<i>Deltaproteobacteria</i>	0.1	0.4	0.0	0.0	0.0	0.0	0.0	0.0
<i>Epsilonproteobacteria</i>	0.1	0.2	0.0	0.0	0.0	0.0	0.2	0.6
<i>Gammaproteobacteria</i>	1.5	2.4	0.8	0.6	4.5	4.5	4.8	5.8
<i>Saccharibacteria;uncultured bacterium</i>	0.3	0.5	0.2	0.4	0.6	1.3	0.8	1.1
<i>Spirochaetes</i>	0.1	0.4	0.0	0.0	0.5	1.7	0.0	0.0
<i>TM6 (Dependentiae);uncultured bacterium</i>	0.0	0.0	0.0	0.0	0.1	0.3	0.1	0.2
<i>Verrucomicrobiae</i>	0.0	0.0	0.0	0.2	0.1	0.3	0.0	0.0
Order								
<i>Acidimicrobiales</i>	0.0	0.0	0.1	0.1	0.5	0.9	0.9	1.6
<i>Bifidobacteriales</i>	0.2	0.5	0.1	0.1	0.3	0.7	0.3	0.4
<i>Corynebacteriales</i>	3.4	4.8	3.0	3.0	10.0	7.1	5.2	7.8
<i>Micrococcales</i>	2.3	1.4	1.5	0.6	0.8	0.6	0.9	0.4
<i>Propionibacteriales</i>	0.0	0.1	0.2	0.2	0.4	1.1	0.3	0.3
<i>Streptomycetales</i>	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
<i>Coriobacteriales</i>	0.4	0.9	0.0	0.1	0.4	0.8	0.1	0.3
<i>Solirubrobacterales</i>	0.2	0.8	0.0	0.0	0.1	0.2	0.0	0.0
<i>Bacteroidales</i>	1.3	4.3	0.1	0.2	0.1	0.1	0.4	0.5
<i>Flavobacteriales</i>	0.1	0.2	0.1	0.1	0.3	0.2	0.4	0.5
<i>Sphingobacteriales</i>	0.0	0.0	0.0	0.0	0.2	0.3	0.2	0.2
<i>Chlamydiales</i>	0.1	0.2	0.4	0.4	1.1	1.0	1.6	1.6
<i>Chloroflexi;KD4-96;uncultured bacterium</i>	0.0	0.0	0.0	0.1	0.1	0.2	0.1	0.2
<i>Chloroplast;Other</i>	1.5	2.8	0.2	0.5	5.4	4.0	1.9	2.5
<i>Bacillales</i>	2.8	1.1	1.8	0.7	1.7	1.2	1.1	0.7
<i>Lactobacillales</i>	80.8	15.9	89.2	4.2	66.4	12.7	74.4	13.1
<i>Clostridiales</i>	3.2	7.0	1.0	0.5	1.5	1.6	2.1	1.2
<i>Erysipelotrichales</i>	0.2	0.6	0.0	0.0	0.0	0.0	0.1	0.1
<i>Selenomonadales</i>	0.3	0.8	0.0	0.1	0.2	0.3	0.3	0.4
<i>Gracilibacteria;Other</i>	0.0	0.0	0.1	0.3	0.2	0.9	0.0	0.0
<i>Planctomycetales</i>	0.1	0.3	0.3	0.2	0.8	0.9	1.4	1.7
<i>Rhizobiales</i>	0.2	0.8	0.2	0.2	1.5	2.0	1.2	2.5
<i>Rhodobacterales</i>	0.0	0.1	0.0	0.1	0.2	0.3	0.3	0.4
<i>Rhodospirillales</i>	0.0	0.0	0.0	0.0	0.1	0.1	0.3	0.9
<i>Rickettsiales</i>	0.3	0.6	0.1	0.2	0.6	0.8	0.2	0.4
<i>Sphingomonadales</i>	0.0	0.2	0.0	0.0	0.1	0.2	0.1	0.1
<i>Burkholderiales</i>	0.2	0.4	0.1	0.1	0.5	0.7	0.1	0.2
<i>Campylobacterales</i>	0.1	0.2	0.0	0.0	0.0	0.0	0.2	0.6
<i>Aeromonadales</i>	0.5	1.6	0.0	0.0	0.0	0.0	0.0	0.0
<i>Enterobacteriales</i>	0.3	0.7	0.1	0.1	0.2	0.3	0.2	0.3
<i>Gammaproteobacteria;HTA4</i>	0.1	0.2	0.0	0.1	0.4	1.0	0.3	0.6

<i>Legionellales</i>	0.2	0.2	0.3	0.4	1.9	2.7	1.4	1.1
<i>Pseudomonadales</i>	0.1	0.3	0.1	0.1	0.2	0.3	0.2	0.2
<i>Vibrionales</i>	0.3	1.2	0.0	0.1	0.3	0.6	1.6	6.1
<i>Xanthomonadales</i>	0.0	0.0	0.0	0.0	0.2	0.2	0.2	0.3
<i>Gammaproteobacteria; Other</i>	0.0	0.1	0.1	0.2	1.2	1.1	0.8	1.0
<i>Saccharibacteria; uncultured bacterium</i>	0.3	0.5	0.2	0.4	0.6	1.3	0.8	1.1
<i>Spirochaetales</i>	0.1	0.4	0.0	0.0	0.5	1.7	0.0	0.0
<i>TM6 (Dependentiae); uncultured bacterium;</i>	0.0	0.0	0.0	0.0	0.1	0.3	0.1	0.2
<i>Verrucomicrobiales</i>	0.0	0.0	0.0	0.2	0.1	0.3	0.0	0.0
<i>WS6; Other</i>	0.0	0.0	0.0	0.1	0.1	0.6	0.0	0.0
<i>Unassigned; Other</i>	0.0	0.1	0.1	0.2	0.2	0.8	0.2	0.4
Family								
<i>Acidimicrobiales; OM1 clade</i>	0.0	0.0	0.0	0.0	0.2	0.4	0.4	0.8
<i>Acidimicrobiales; uncultured</i>	0.0	0.0	0.0	0.0	0.1	0.2	0.4	1.3
<i>Bifidobacteriaceae</i>	0.2	0.5	0.1	0.1	0.3	0.7	0.3	0.4
<i>Corynebacteriaceae</i>	2.9	4.1	2.4	2.6	6.1	5.2	1.8	2.9
<i>Mycobacteriaceae</i>	0.4	1.0	0.6	1.2	3.9	3.9	3.3	7.5
<i>Brevibacteriaceae</i>	0.8	0.9	0.4	0.5	0.0	0.1	0.1	0.2
<i>Dermabacteraceae</i>	0.1	0.2	0.1	0.2	0.0	0.0	0.0	0.1
<i>Intrasporangiaceae</i>	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.1
<i>Microbacteriaceae</i>	0.5	0.5	0.3	0.3	0.5	0.4	0.4	0.3
<i>Micrococcaceae</i>	0.9	0.8	0.6	0.4	0.2	0.3	0.3	0.3
<i>Nocardoidaceae</i>	0.0	0.0	0.0	0.0	0.3	1.0	0.0	0.0
<i>Propionibacteriaceae</i>	0.0	0.1	0.2	0.2	0.1	0.2	0.2	0.3
<i>Coriobacteriaceae</i>	0.4	0.9	0.0	0.1	0.4	0.8	0.1	0.3
<i>Solirubrobacterales; Elev-16S-1332</i>	0.2	0.8	0.0	0.0	0.0	0.0	0.0	0.0
<i>Bacteroidaceae</i>	0.1	0.4	0.0	0.1	0.0	0.1	0.2	0.4
<i>Prevotellaceae</i>	0.9	3.1	0.0	0.1	0.0	0.0	0.1	0.2
<i>Flavobacteriaceae</i>	0.1	0.1	0.1	0.1	0.3	0.2	0.4	0.4
<i>Chitinophagaceae</i>	0.0	0.0	0.0	0.0	0.2	0.3	0.1	0.2
<i>Chlamydiales; Other</i>	0.1	0.2	0.4	0.4	1.1	1.0	1.6	1.6
<i>Chloroflexi; KD4-96; uncultured bacterium</i>	0.0	0.0	0.0	0.1	0.1	0.2	0.1	0.2
<i>Chloroplast; Other</i>	1.5	2.8	0.2	0.5	5.4	4.0	1.9	2.5
<i>Bacillaceae</i>	1.0	0.7	0.8	0.5	0.9	0.6	0.6	0.4
<i>Paenibacillaceae</i>	0.1	0.2	0.0	0.1	0.1	0.2	0.1	0.1
<i>Planococcaceae</i>	0.1	0.2	0.1	0.1	0.2	0.2	0.1	0.1
<i>Staphylococcaceae</i>	1.4	1.0	0.9	0.4	0.6	1.3	0.3	0.5
<i>Bacillales; Other</i>	0.1	0.4	0.0	0.0	0.0	0.0	0.0	0.0
<i>Aerococcaceae</i>	0.1	0.1	0.1	0.1	0.0	0.1	0.0	0.1
<i>Carnobacteriaceae</i>	0.0	0.1	0.0	0.1	0.1	0.2	0.1	0.2

<i>Enterococcaceae</i>	0.4	0.3	0.3	0.2	0.4	0.4	0.4	0.4
<i>Lactobacillaceae</i>	77.9	16.1	86.5	4.4	61.3	12.4	67.6	12.2
<i>Leuconostocaceae</i>	0.5	0.8	1.0	1.1	0.5	1.3	3.0	2.7
<i>Streptococcaceae</i>	2.0	1.5	1.3	1.4	4.1	3.7	3.2	2.3
<i>Clostridiaceae 1</i>	0.7	0.7	0.2	0.2	0.4	0.4	0.4	0.2
<i>Clostridiaceae 2</i>	0.0	0.0	0.0	0.0	0.1	0.5	0.0	0.0
<i>Clostridiales; Family XI</i>	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.3
<i>Clostridiales; Family XIII</i>	0.1	0.2	0.1	0.3	0.0	0.0	0.0	0.0
<i>Lachnospiraceae</i>	0.6	2.1	0.2	0.3	0.1	0.2	0.4	0.5
<i>Peptostreptococcaceae</i>	0.2	0.3	0.1	0.1	0.1	0.2	0.1	0.2
<i>Ruminococcaceae</i>	1.1	4.1	0.0	0.1	0.0	0.0	0.4	0.6
<i>Clostridiales; Other</i>	0.1	0.3	0.1	0.2	0.6	0.9	0.5	0.8
<i>Erysipelotrichaceae</i>	0.2	0.6	0.0	0.0	0.0	0.0	0.1	0.1
<i>Acidaminococcaceae</i>	0.2	0.6	0.0	0.0	0.0	0.0	0.1	0.3
<i>Veillonellaceae</i>	0.2	0.3	0.0	0.1	0.2	0.3	0.1	0.2
<i>Gracilibacteria; Other</i>	0.0	0.0	0.1	0.3	0.2	0.9	0.0	0.0
<i>Planctomycetaceae</i>	0.1	0.3	0.3	0.2	0.8	0.9	1.4	1.7
<i>Bradyrhizobiaceae</i>	0.0	0.0	0.0	0.0	0.7	1.6	0.1	0.3
<i>Brucellaceae</i>	0.0	0.0	0.0	0.0	0.2	0.5	0.1	0.2
<i>Hyphomicrobiaceae</i>	0.1	0.2	0.0	0.0	0.0	0.1	0.1	0.5
<i>Phyllobacteriaceae</i>	0.2	0.5	0.1	0.2	0.2	0.5	0.5	1.8
<i>Rhizobiaceae</i>	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.3
<i>Rhizobiales; Other</i>	0.0	0.0	0.0	0.1	0.2	0.3	0.2	0.2
<i>Rhodobacteraceae</i>	0.0	0.1	0.0	0.1	0.2	0.3	0.3	0.4
<i>Acetobacteraceae</i>	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.9
<i>Mitochondria</i>	0.3	0.6	0.1	0.2	0.6	0.8	0.2	0.4
<i>Sphingomonadaceae</i>	0.0	0.2	0.0	0.0	0.1	0.2	0.1	0.1
<i>Comamonadaceae</i>	0.2	0.4	0.0	0.1	0.3	0.4	0.1	0.2
<i>Oxalobacteraceae</i>	0.0	0.0	0.0	0.1	0.2	0.4	0.0	0.1
<i>Helicobacteraceae</i>	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.6
<i>Aeromonadaceae</i>	0.2	0.5	0.0	0.0	0.0	0.0	0.0	0.0
<i>Succinivibrionaceae</i>	0.3	1.3	0.0	0.0	0.0	0.0	0.0	0.0
<i>Enterobacteriaceae</i>	0.3	0.7	0.1	0.1	0.2	0.3	0.2	0.3
<i>Gammaproteobacteria; HTA4; Other</i>	0.1	0.2	0.0	0.1	0.4	1.0	0.3	0.6
<i>Coxiellaceae</i>	0.1	0.2	0.1	0.1	0.5	0.9	0.4	0.4
<i>Legionellaceae</i>	0.1	0.2	0.3	0.3	1.4	2.0	1.0	0.9
<i>Moraxellaceae</i>	0.1	0.3	0.1	0.1	0.2	0.2	0.1	0.3
<i>Vibrionaceae</i>	0.3	1.2	0.0	0.1	0.3	0.6	1.6	6.1
<i>Xanthomonadaceae</i>	0.0	0.0	0.0	0.0	0.1	0.2	0.2	0.3
<i>Gammaproteobacteria; Other</i>	0.0	0.1	0.1	0.2	1.2	1.1	0.8	1.0
<i>Saccharibacteria; uncultured bacterium</i>	0.3	0.5	0.2	0.4	0.6	1.3	0.8	1.1
<i>Brevinemataceae</i>	0.0	0.0	0.0	0.0	0.5	1.7	0.0	0.0

<i>TM6 (Dependentiae); uncultured bacterium</i>	0.0	0.0	0.0	0.0	0.1	0.3	0.1	0.2
<i>Verrucomicrobiaceae</i>	0.0	0.0	0.0	0.2	0.1	0.2	0.0	0.0
<i>WS6;Other</i>	0.0	0.0	0.0	0.1	0.1	0.6	0.0	0.0
<i>Unassigned;Other</i>	0.0	0.1	0.1	0.2	0.2	0.8	0.2	0.4
Genus								
<i>Acidimicrobiales; OMI clade; uncultured bacterium</i>	0.0	0.0	0.0	0.0	0.2	0.4	0.4	0.8
<i>Acidimicrobiales; uncultured;Other</i>	0.0	0.0	0.0	0.0	0.0	0.1	0.4	1.3
<i>Bifidobacterium</i>	0.2	0.5	0.1	0.1	0.3	0.7	0.3	0.4
<i>Corynebacterium 1</i>	2.8	4.1	2.3	2.5	6.1	5.2	1.8	2.9
<i>Mycobacterium</i>	0.4	1.0	0.6	1.2	3.9	3.9	3.3	7.5
<i>Nocardia</i>	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
<i>Brevibacterium</i>	0.8	0.9	0.4	0.5	0.0	0.1	0.1	0.2
<i>Brachybacterium</i>	0.1	0.2	0.1	0.2	0.0	0.0	0.0	0.1
<i>Intrasporangiaceae;Other</i>	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.1
<i>Leucobacter</i>	0.3	0.4	0.1	0.2	0.2	0.4	0.1	0.2
<i>Microbacteriaceae;Other</i>	0.2	0.5	0.2	0.3	0.3	0.4	0.2	0.3
<i>Arthrobacter</i>	0.2	0.4	0.3	0.4	0.1	0.2	0.1	0.2
<i>Glutamicibacter</i>	0.2	0.6	0.1	0.2	0.0	0.1	0.0	0.1
<i>Kocuria</i>	0.3	0.4	0.2	0.2	0.1	0.3	0.2	0.2
<i>Micrococcaceae;Other</i>	0.1	0.1	0.0	0.0	0.0	0.1	0.0	0.1
<i>Nocardioides</i>	0.0	0.0	0.0	0.0	0.3	1.0	0.0	0.0
<i>Propionibacterium</i>	0.0	0.1	0.1	0.1	0.1	0.1	0.2	0.3
<i>Collinsella</i>	0.2	0.8	0.0	0.0	0.0	0.0	0.0	0.1
<i>Enterorhabdus</i>	0.0	0.1	0.0	0.0	0.3	0.8	0.0	0.0
<i>Coriobacteriaceae; uncultured</i>	0.1	0.5	0.0	0.0	0.0	0.0	0.0	0.0
<i>Solirubrobacterales; Elev-16S-1332 uncultured bacterium</i>	0.2	0.8	0.0	0.0	0.0	0.0	0.0	0.0
<i>Bacteroides</i>	0.1	0.4	0.0	0.1	0.0	0.1	0.2	0.4
<i>Bacteroidales S24-7 group; uncultured bacterium</i>	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0
<i>Prevotella 2</i>	0.1	0.5	0.0	0.0	0.0	0.0	0.0	0.0
<i>Prevotella 9</i>	0.7	2.2	0.0	0.1	0.0	0.0	0.1	0.2
<i>Cloacibacterium</i>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
<i>Flavobacterium</i>	0.0	0.1	0.0	0.0	0.1	0.2	0.2	0.2
<i>Flavobacteriaceae;Other</i>	0.0	0.1	0.0	0.1	0.1	0.2	0.1	0.3
<i>Sediminibacterium</i>	0.0	0.0	0.0	0.0	0.1	0.2	0.1	0.2
<i>Chlamydiales;Other</i>	0.1	0.2	0.4	0.4	1.1	1.0	1.6	1.6
<i>Chloroflexi; KD4-96; uncultured bacterium</i>	0.0	0.0	0.0	0.1	0.1	0.2	0.1	0.2
<i>Chloroplast;Other</i>	1.5	2.8	0.2	0.5	5.4	4.0	1.9	2.5

<i>Bacillus</i>	0.5	0.4	0.5	0.4	0.8	0.6	0.5	0.4
<i>Bacillaceae; Other</i>	0.4	0.3	0.3	0.3	0.0	0.1	0.1	0.1
<i>Brevibacillus</i>	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.1
<i>Paenibacillus</i>	0.0	0.0	0.0	0.0	0.1	0.2	0.0	0.1
<i>Planococcaceae; Other</i>	0.1	0.1	0.1	0.1	0.2	0.2	0.0	0.1
<i>Staphylococcus</i>	1.3	1.0	0.8	0.4	0.5	1.3	0.3	0.5
<i>Staphylococcaceae; Other</i>	0.1	0.2	0.1	0.2	0.1	0.2	0.0	0.1
<i>Bacillales; Other</i>	0.1	0.4	0.0	0.0	0.0	0.0	0.0	0.0
<i>Granulicatella</i>	0.0	0.1	0.0	0.1	0.1	0.2	0.1	0.2
<i>Enterococcus</i>	0.3	0.3	0.3	0.2	0.4	0.4	0.4	0.4
<i>Lactobacillus</i>	77.9	16.1	86.5	4.4	61.3	12.4	67.6	12.2
<i>Leuconostoc</i>	0.0	0.1	0.3	0.3	0.1	0.2	0.1	0.3
<i>Weissella</i>	0.4	0.8	0.7	0.9	0.4	1.3	2.8	2.8
<i>Lactococcus</i>	0.3	0.4	0.1	0.2	0.5	0.4	0.3	0.3
<i>Streptococcus</i>	1.6	1.6	1.2	1.4	3.6	3.5	2.9	2.2
<i>Clostridium sensu stricto 1</i>	0.4	0.7	0.1	0.1	0.0	0.1	0.1	0.1
<i>Clostridiaceae 1; Other</i>	0.2	0.4	0.1	0.2	0.3	0.4	0.2	0.3
<i>Alkaliphilus</i>	0.0	0.0	0.0	0.0	0.1	0.5	0.0	0.0
<i>Clostridiales; Family XI; uncultured</i>	0.1	0.2	0.0	0.1	0.0	0.2	0.0	0.1
<i>Clostridiales; Family XI; Other</i>	0.1	0.2	0.2	0.2	0.1	0.2	0.2	0.3
<i>Blautia</i>	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.2
<i>Roseburia</i>	0.2	0.6	0.0	0.2	0.0	0.0	0.0	0.0
<i>Peptostreptococcaceae; Other</i>	0.2	0.3	0.0	0.1	0.1	0.2	0.0	0.1
<i>Faecalibacterium</i>	0.3	1.0	0.0	0.1	0.0	0.0	0.1	0.3
<i>Ruminococcaceae UCG-002</i>	0.1	0.6	0.0	0.0	0.0	0.0	0.0	0.1
<i>Ruminococcaceae UCG-005</i>	0.2	0.7	0.0	0.0	0.0	0.0	0.0	0.0
<i>Ruminococcus 2</i>	0.1	0.4	0.0	0.1	0.0	0.0	0.0	0.1
<i>[Eubacterium] coprostanoligenes group</i>	0.1	0.4	0.0	0.0	0.0	0.0	0.0	0.0
<i>Ruminococcaceae; uncultured</i>	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.3
<i>Clostridiales; Other</i>	0.1	0.3	0.1	0.2	0.6	0.9	0.5	0.8
<i>Phascolarctobacterium</i>	0.2	0.6	0.0	0.0	0.0	0.0	0.0	0.1
<i>Acidaminococcaceae; Other</i>	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.2
<i>Megasphaera</i>	0.0	0.1	0.0	0.1	0.2	0.2	0.1	0.1
<i>Gracilibacteria; Othe</i>	0.0	0.0	0.1	0.3	0.2	0.9	0.0	0.0
<i>Planctomycetaceae; Pir4 lineage</i>	0.0	0.0	0.1	0.1	0.2	0.2	0.2	0.0
<i>Planctomyces</i>	0.0	0.1	0.1	0.2	0.3	0.3	0.6	0.0
<i>Planctomycetaceae; uncultured</i>	0.1	0.3	0.1	0.2	0.2	0.4	0.4	0.1
<i>Bradyrhizobium</i>	0.0	0.0	0.0	0.0	0.7	1.6	0.1	0.0
<i>Ochrobactrum</i>	0.0	0.0	0.0	0.0	0.2	0.5	0.1	0.0
<i>Hyphomicrobium</i>	0.1	0.2	0.0	0.0	0.0	0.0	0.1	0.0
<i>Mesorhizobium</i>	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1
<i>Phyllobacteriaceae; Other</i>	0.2	0.5	0.1	0.2	0.2	0.5	0.4	0.4

<i>Rhizobiales; Other</i>	0.0	0.0	0.0	0.1	0.2	0.3	0.2	0.2
<i>Rhodobacteraceae; Other</i>	0.0	0.0	0.0	0.1	0.1	0.2	0.2	0.0
<i>Acetobacteraceae; Other</i>	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.1
<i>Mitochondria;Other</i>	0.3	0.6	0.1	0.2	0.6	0.8	0.2	0.0
<i>Delftia</i>	0.0	0.0	0.0	0.1	0.3	0.4	0.1	0.0
<i>Comamonadaceae;Other</i>	0.1	0.4	0.0	0.0	0.0	0.0	0.0	0.0
<i>Oxalobacteraceae;Other</i>	0.0	0.0	0.0	0.1	0.2	0.4	0.0	0.0
<i>Succinivibrio</i>	0.3	1.3	0.0	0.0	0.0	0.0	0.0	0.6
<i>Escherichia-Shigella</i>	0.2	0.7	0.0	0.0	0.0	0.1	0.1	0.0
<i>Serratia</i>	0.1	0.2	0.0	0.0	0.2	0.2	0.1	0.0
<i>Gammaproteobacteria; HTA4;Other</i>	0.1	0.2	0.0	0.1	0.4	1.0	0.3	0.0
<i>Aquicella</i>	0.0	0.1	0.0	0.1	0.2	0.4	0.2	0.2
<i>Coxiella</i>	0.1	0.2	0.0	0.1	0.3	0.5	0.2	0.0
<i>Legionella</i>	0.1	0.2	0.3	0.3	1.3	2.0	0.8	0.3
<i>Legionellaceae; Other</i>	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.0
<i>Acinetobacter</i>	0.1	0.2	0.1	0.1	0.2	0.2	0.1	0.5
<i>Photobacterium</i>	0.2	1.0	0.0	0.0	0.2	0.6	0.4	0.0
<i>Vibrio</i>	0.1	0.2	0.0	0.1	0.0	0.0	1.2	0.2
<i>Stenotrophomonas</i>	0.0	0.0	0.0	0.0	0.1	0.2	0.1	0.1
<i>Gammaproteobacteria;Other;</i>	0.0	0.1	0.1	0.2	1.2	1.1	0.8	4.6
<i>Saccharibacteria; uncultured bacterium;</i>	0.3	0.5	0.2	0.4	0.6	1.3	0.8	0.1
<i>Brevinema</i>	0.0	0.0	0.0	0.0	0.5	1.7	0.0	0.1
<i>TM6 (Dependentiae); uncultured bacterium</i>	0.0	0.0	0.0	0.0	0.1	0.3	0.1	0.1
<i>WS6;Other</i>	0.0	0.0	0.0	0.1	0.1	0.6	0.0	0.0
<i>Unassigned;Other</i>	0.0	0.1	0.1	0.2	0.2	0.8	0.2	0.0