

A transcriptomic analysis of diploid and triploid Atlantic salmon lenses with and without cataracts

Pål A. Olsvik^{a,b,*}, Roderick Nigel Finn^{c,d}, Sofie C. Remø^b, Per G. Fjellidal^b, François Chauvigné^d, Kevin A. Glover^{b,c}, Tom Hansen^b, Rune Waagbø^{b,c}

^a Faculty of Biosciences and Aquaculture, Nord University, Bodø, Norway

^b Institute of Marine Research, Nordnes, Bergen, Norway

^c Department of Biological Sciences, University of Bergen, Bergen, Norway

^d IRTA-Institute of Biotechnology and Biomedicine (IBB), Universitat Autònoma de Barcelona, 08193, Bellaterra, Cerdanyola del Vallès, Spain

ARTICLE INFO

Keywords:

Triploid Atlantic salmon
Cataracts
Transcriptional responses
RNA-seq

ABSTRACT

To avoid negative environmental impacts of escapees and potential inter-breeding with wild populations, the Atlantic salmon farming industry has and continues to extensively test triploid fish that are sterile. However, they often show differences in performance, physiology, behavior and morphology compared to diploid fish, with increased prevalence of vertebral deformities and ocular cataracts as two of the most severe disorders. Here, we investigated the mechanisms behind the higher prevalence of cataracts in triploid salmon, by comparing the transcriptional patterns in lenses of diploid and triploid Atlantic salmon, with and without cataracts. We assembled and characterized the Atlantic salmon lens transcriptome and used RNA-seq to search for the molecular basis for cataract development in triploid fish. Transcriptional screening showed only modest differences in lens mRNA levels in diploid and triploid fish, with few uniquely expressed genes. In total, there were 165 differentially expressed genes (DEGs) between the cataractous diploid and triploid lenses. Of these, most were expressed at lower levels in triploid fish. Differential expression was observed for genes encoding proteins with known function in the retina (phototransduction) and proteins associated with repair and compensation mechanisms. The results suggest a higher susceptibility to oxidative stress in triploid lenses, and that mechanisms connected to the ability to handle damaged proteins are differentially affected in cataractous lenses from diploid and triploid salmon.

1. Introduction

Recent years have seen renewed efforts to establish commercial farming of triploid Atlantic salmon (*Salmo salar*) – hereafter referred to as salmon (Benfey, 2016; Stien et al., 2019). Farming triploid salmon has two major advantages. The first is that triploid females do not mature sexually and can diverge energy into somatic growth (Piferrer et al., 2009). The second is related to the fact that domesticated salmon escapees and genetic interactions with wild conspecifics represents one of the most significant environmental challenges to salmon aquaculture (Glover et al., 2017). Rearing sterile triploid fish reduced this threat, and is an effective way to mitigate further genetic interactions.

Although production of triploid salmon has potential benefits, the global Atlantic salmon aquaculture industry is still primarily based upon rearing diploid fish. While there are several reasons for this, in part, this

is due to the fact that triploid salmon often show poor performance. For example, in comparison with diploid salmon, they display differences in physiology, behavior and morphology, with increased prevalence of vertebral deformity and ocular cataracts as two of the most severe disorders (Wall and Richards, 1992; Piferrer et al., 2009; Taranger et al., 2010; Taylor et al., 2015; Sambraus et al., 2017). Cataracts are defined as the loss of transparency of the lens and can appear both as reversible osmotic cataracts and permanent cataracts, which can have multiple causes (Hejtmancik, 2008). In farmed salmon, cataract formation has been linked to genetic predispositions and several nutritional and environmental factors (reviewed by Bjerkås et al., 2006). Cataract has been observed in both freshwater and seawater, however, farmed salmon are particularly prone to cataract development during the smolt transition from fresh to saltwater (Waagbø et al., 1998; Breck and Sveier, 2001; Breck et al., 2005a; Remø et al., 2014) and during periods of rapid

* Corresponding author. Faculty of Biosciences and Aquaculture, Nord University, Bodø, Norway.

E-mail address: pal.a.olsvik@nord.no (P.A. Olsvik).

<https://doi.org/10.1016/j.yexer.2020.108150>

Received 21 January 2020; Received in revised form 8 July 2020; Accepted 9 July 2020

Available online 28 July 2020

0014-4835/© 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

growth (Bjerkås et al., 1996; Breck and Sveier, 2001; Waagbø et al., 1996, 2010; Remø et al., 2014, 2017). Increased prevalence of cataracts in triploid fish is not well understood but may partly rely on altered metabolism due to differences in cellular morphology (Benfey, 1999).

A sub-optimal level of dietary histidine is currently considered the most important causative factor for cataract development in farmed Atlantic salmon (Breck et al., 2003, 2005b; Remø et al., 2014, 2017; Waagbø et al., 2010). Taylor et al. (2015) investigated the preventive effects of dietary histidine supplementation in triploid Atlantic salmon during seawater grow-out. Although the severity was higher in triploids compared to diploids irrespective of diet, applying a high histidine diet mitigated further cataract development in triploids. Similarly, dietary histidine supplementation reduced the severity of cataracts in diploid and triploid yearling smolt, but also with a higher severity in triploids compared to diploids at the highest dietary level (Sambraus et al., 2017). The cataract preventative effect of dietary histidine has been attributed to the functional roles of histidine and the derivative N-acetyl-histidine (NAH) as buffer component (Breck et al., 2005a), osmolyte (Rhodes et al., 2010) and possibly antioxidant (Remø et al., 2014), therefore being important to maintain cell integrity and water balance. The lens concentration of NAH was lower in triploids compared to diploids given the same high histidine diet (Sambraus et al., 2017), suggesting that the triploid lens may be more vulnerable to cataract development, possibly due to lower protection of the triploid lens through lower ability to synthesize NAH, or a higher requirement to maintain water balance in the lens. The latter might be linked to larger cell size in triploids (Wu et al., 2010). Thus, differences in susceptibility to cataracts, as well as the apparent higher requirement of histidine to mitigate (but not eliminate) cataract development in triploids, may be hypothesized to be due to alterations or weakness in the lens of triploids.

Thus far, no attempts have been conducted to evaluate the mechanisms behind increased prevalence of cataracts in triploid fish at the molecular level. Relatively few genome-wide examinations of the molecular mechanisms behind cataract formation have been performed on healthy and cataractous lenses in vertebrates (Sousounis and Tsonis, 2012), possibly due to the biased lens transcriptome, where the expression of structural genes, such as crystallins, predominates over genes that regulate cell function and phenotype (Wistow, 2006; Manthey et al., 2014). Global transcriptional examinations of the mammalian cataractous lens have revealed differential regulation on numerous types of genes, including crystallins and heat shock proteins, cytochrome oxidases, growth factors, metalloproteinases and collagen, as well as various transcription factors (Wistow et al., 2002; Wride et al., 2003; Hawse et al., 2003; Mansergh et al., 2004; Medvedovic et al., 2006; Hejtmancik, 2008; Shiels et al., 2010; Shiels and Hejtmancik, 2017, 2019). In Atlantic salmon, Tröbe et al. (2009) used a 16K salmonid microarray to screen for transcriptional responses to histidine related cataracts in lenses of Atlantic salmon and reported differences in genes encoding proteins linked to lipid metabolism, carbohydrate metabolism, and protein degradation. Among the significantly differentially regulated genes were gamma crystallin M2 (homolog to mammalian *crygb*), lens fiber membrane *lim2*, secreted protein, acidic, cysteine-rich (*sparc*), metallothionein B (*mt-b*), heat-shock cognate 70 (*hsc70a*), calpain (*capns1*), Na/K ATPase alpha subunit isoform 1c (*atpa1c*) and fatty acid binding protein 2 (*fabp2*), of which several have been linked to cataracts before.

In the present study, we used transcriptomics (RNA-seq) to examine why triploid fish are more prone to cataract development than diploid fish. To do so, we compared the transcriptional patterns in the lens of diploid and triploid Atlantic salmon originating from both a domesticated strain and a wild population, with and without mature cataract, as assessed by a slit-lamp biomicroscope.

2. Materials and methods

2.1. Experimental animals and set-up

The salmon used in this experiment originated from females (f) and males (m) from the domesticated Mowi strain (M) crossed with females and males from a wild population in the river Figgjo (F) in November 2011. Eight groups were made as diploid and triploid of the systematic breeding of the farmed and wild strains: mM × fM, mM × fF, mF × fF and mF × fM. The offspring groups were start-fed with a commercial feed (Skretting, Stavanger, Norway) at March 26th 2012 and were held at 12 °C water temperature from start feeding to mid-summer. Thereafter, the groups were reared at ambient temperature. Fish were reared under continuous light from start feeding to October 1st, followed by a simulated natural photoperiod to initiate parr-smolt transformation. Experimental groups were held separately in eight tanks until November 27th 2012, when they were individually passive integrated transponder-tagged (PIT-tags, Electronic I, Inc., Dallas, TX, USA) and the groups distributed equally into three replicate tanks. Fish were transferred to seawater at May 10th, 2013. In sea, the fish were fed Skretting Spirit 75-50A. The experiment was terminated October 16th 2013.

Fish were sampled as post smolts in seawater at a mean body weight of 143 ± 8 g ($n = 46$). Upon sampling, the fish were inspected for cataracts, and weight and length measured. From each fish, two lenses and heart tissues were dissected and immediately frozen on liquid nitrogen. The left lens was used for transcriptome *de novo* assembly and transcriptomics.

2.2. Cataract determination

Cataract assessment was performed on anaesthetized fish by use of a Kowa SL-15 slit-lamp biomicroscope (Kowa, Tokyo, Japan). The type, position and severity of the observed cataractous changes were determined according to Wall and Richards (1992), but with a maximum severity extended from 3 to 4 per eye to match the amplitude of the macroscopic scale (microscopic cataract score 0: absent, 1: slight, 2: moderate, 3: severe, 4: total cataract).

2.3. Histidine determination

Heart tissue from the sampled fish was used as status organ for histidine and NAH (Remø et al., 2014). NAH and free histidine in the heart tissue were analyzed by reverse phase HPLC and UV detection at 210 nm, with modifications according to Breck et al. (2005b).

2.4. RNA isolation

Lens tissue was thoroughly homogenized before RNA extraction using a Precellys 24 homogenizer and ceramic beads CK28 (Bertin Technologies, Montigny-le-Bretonneux, France). Total RNA was extracted using the BioRobot EZ1 and RNA Tissue Mini Kit (Qiagen, Hilden, Germany), treated with DNase according to the manufacturer's instructions and eluted in 50 µL RNase-free MilliQ H₂O. RNA quality and integrity were assessed with the NanoDrop ND-1000 UV-Vis Spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA) and the Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA). The RNA 6000 Nano LabChip kit (Agilent Technologies, Palo Alto, CA, USA) was used to evaluate the RNA integrity of the lens samples. The RNA integrity number (RIN) of RNA extracted for transcriptome assembly (2N: $n = 10$, 3N: $n = 10$) and RNA-seq (2N: $n = 12$, 3N: $n = 14$) were 8.5 ± 0.0 ($n = 20$) and 8.2 ± 0.1 ($n = 26$), respectively (mean \pm SEM).

2.5. Lens transcriptome *de novo* assembly

Since no Atlantic salmon reference genome was available at the time

of analysis, transcriptome *de novo* assembly had to be conducted using Illumina paired-end reads before RNA-seq analyses. RNA extracted from 10 diploid and 10 triploid lenses ($n = 20$) was mixed and used to generate the assembly. Transcriptome *de novo* assembly was conducted using the short reads assembling software Trinity as described by Grabherr et al. (2011). Trinity combines three independent software modules, Inchworm, Chrysalis, and Butterfly, to process the RNA-seq reads into unigenes. The output sequences were aligned to the databases of NR, NT, SwissProt, KEGG, COG and GO using Blastx, and the best aligning result was used to decide sequence direction. Sequence direction of unigenes not aligned to any of the above-mentioned databases was determined with the ESTScan software (Iseli et al., 1999).

2.6. RNA-seq analysis

Direct RNA sequencing (RNA-seq) was used to screen for differentially expressed genes (DEGs) in lenses of both diploid and triploid individuals. As the two strains used here are known to display divergent transcription patterns (Bicskei et al., 2014, 2016), fish from both groups were randomly mixed and pooled prior to any analysis. Individual left lenses from 26 salmon were used for RNA-seq examination (2N-: $n = 6$, 2N+: $n = 6$, 3N-: $n = 6$, 3N+: $n = 8$). Poly (A) mRNA was isolated using magnetic beads with oligo (dT) from total RNA obtained from the lens samples. Fragmentation buffer was added to shred mRNA to short reads. Using these short fragments (about 200 bp) as templates, random hexamer primers were applied to synthesize first-strand cDNA. Second-strand cDNA was synthesized using buffer, dNTPs, RNaseH, and DNA polymerase I. QiaQuick PCR extraction kit (Qiagen) was used to purify short double-stranded cDNA fragments according to manufacturer's instructions. These fragments were then resolved with EB buffer for end reparation, added poly (A), and ligated to the sequencing adapters. After agarose gel electrophoresis, the suitable fragments were selected for PCR amplification as templates. Finally, the libraries were sequenced using Illumina HiSeq™ 2000 (San Diego, CA, USA).

The *de novo* lens transcriptome described above was thereafter used as a reference for alignment of the RNA-seq data. Unigenes were annotated with Blastx alignment using an e-value cut-off of 10^{-5} between unigenes and the databases of NR, NT, SwissProt, KEGG, COG and GO. The NOISeq software package (Tarazona et al., 2011) was used to screen for differentially expressed genes (DEGs). NOISeq is a novel non-parametric method for the identification of DEGs, which shows a good performance when compared to other differential expression methods, like Fisher's Exact Test, edgeR, DESeq and baySeq. All RNA-seq work was performed by staff at the Beijing Genome Institute (BGI, Hong Kong).

2.7. Ploidy verification

Fish from each ploidy were sampled and measured for erythrocyte diameter to verify their ploidy status (Benfey et al., 1984). Blood smears were used to measure the relative diameters of 10 erythrocytes per fish (Image-Pro Plus, version 4.0, Media Cybernetics Silver Spring). The triploid fish had significantly (22%) larger blood cells than diploid fish.

2.8. Statistics

To calculate differential expression, NOISeq default settings were used (Tarazona et al., 2011). NOISeq empirically models the noise distribution of count changes by contrasting fold-change differences (M) and absolute expression differences (D) for all the features in samples within the same condition. This reference distribution is used to assess whether the M-D values computed between two conditions for a given gene is likely to be part of the noise or represent a true differential expression. Instead of using a false discovery rate (FDR) or a q-value cut-off, the NOISeq method calculates a differential expression probability value. A gene is declared as differentially expressed if this

probability is higher than q . The threshold q is set to 0.8 by default, since this value is equivalent to an odd of 4:1 (the gene is 4 times more likely to be differentially expressed than not). In this work we used a \log_2 M-value cut-off of ≥ 2 (fold-change ≥ 2). For genes not expressed in some samples, the gene expression value (D) of 0.001 was used. Functional pathway analyses, including prediction of activation and inhibition of upstream transcription factors and downstream effects, were generated through the use of QIAGEN's Ingenuity Pathway Analysis (IPA, QIAGEN Redwood City, www.qiagen.com/ingenuity). Since IPA only can map mammalian homolog entries, identifiers were obtained with Blast alignment against the RefSeq databases (cut-off $E10^{-5}$) and assuming orthologous genes have the same function. A limited number of fish-specific genes with no mammalian homologs were for this reason not included in the IPA pathway analysis. This may have skewed the interpretation of the transcriptomic data.

3. Data availability

The RNA-seq dataset discussed in this publication has been deposited in NCBI's Gene Expression Omnibus and is accessible through GEO Series accession number GSE153933 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE153933>).

4. Results

4.1. Growth and heart histidine levels

Growth of the farmed and wild stocks is reported in detail by Harvey et al. (2017), and therefore not reported here. At the present sampling, there was no significant difference in weight between the diploid and triploid fish. Fig. 1 shows the concentrations of L-histidine and N-acetyl-L-histidine (NAH) analyzed in salmon heart as a measure on the ambient histidine status. Two-way ANOVA analysis showed that there




Fig. 1. Heart concentrations of A) L-histidine and B) N-acetyl-L-histidine in diploid and triploid Atlantic salmon. Significance levels of two-factor analyses are shown in figures (2-way ANOVA). Capped lines indicate significance levels of direct comparisons of the compounds in 2N and 3N fish (uncorrected Fisher's test, $P < 0.05$). ns = not significant.

was a significant effect of cataract on L-histidine (Fig. 1A, $p = 0.016$), while there were no significant effects of either cataract or ploidy on NAH (Fig. 1B). Posthoc tests showed reduced levels of L-histidine ($p = 0.019$) and increased levels of N-acetyl-L-histidine ($p = 0.046$) in diploid fish with cataract.

4.2. Cataract status

The mean cataract scores of the left lenses used to generate the lens transcriptome (2N+ had score 2.5 and 3N+ had score 1.5, while both 2N- and 3N- had score 0) and the right lenses used for the RNA-seq analysis (both 2N+ and 3N+ had score 3, while both 2N- and 3N- had score 0) are shown in Fig. 2A and B.

4.3. Lens de novo transcriptome assembly

Total RNA extracted from lenses of 20 domesticated and wild salmon, 10 of which were diploid and 10 were triploid, was mixed (Fig. 2A), sequenced and used to assemble the lens transcriptome. Using the Illumina HiSeq 2000 platform, a total of 68,403,252 raw reads and 63,098,790 clean reads were sequenced. The clean reads were

assembled into 78,306 contigs with mean length of 284 nucleotides (nts). Of these, 29,177 contigs with mean length of 659 nts mapped to UniGene entries. Distinct clusters, which contained highly similar UniGene entries (more than 70% that may come from the same gene or homologous genes), were 7391. Distinct singletons representing a single UniGene were 21,786. Using blastx and blastn, a cut-off of 10^{-5} and the following priority order, 15,711, 21,084, 14,445, 11,680, 5231, 11,327 UniGene entries were functionally annotated to the NR, NT, Swiss-Prot, KEGG, COG and GO databases, respectively. UniGene entries aligned to a higher-priority database were not aligned to a lower-priority database. In total, 22,160 out of the 29,177 UniGene entries were given a functional annotation in these databases.

For protein coding region prediction analysis, the number of coding DNA sequences (CDS) that mapped to the protein database was 15,428. The number of predicted CDS, Unigene entries that could not be aligned to any database and were scanned by ESTScan, was 701. The total number of CDS was 16,129. According to the microsatellite analysis conducted with the MicroSATellite (MISA) software and using Unigenes as reference, there were 7274 simple sequence repeat (SSR) in the transcriptome. Heterozygous analysis using SOAPSnp, a member of the Short Oligonucleotide Analysis Package (SOAP), revealed 23,759 single-




Fig. 2. Cataract score of the Atlantic salmon lenses used to A) generate a lens transcriptome and B) quantification of DEGs. C) Number of DEGs in lenses of diploid (2N) and triploid (3N) Atlantic salmon without (-) and with cataracts (+). Exact numbers of genes are given above the individual bars. D) Venn diagram showing the degree of overlap of DEGs in the four treatment groups based on a four-way comparison of genes.

nucleotide polymorphisms (SNPs) in the transcriptome.

A summary of the NR annotation is shown in Fig. S1 (Supplementary File 1). Fig. S1A shows E-value distribution, while similarity distribution is shown in Fig. S1B. Fig. S1C shows the species distribution of UniGene entries annotated to the NR database. Most UniGene entries mapped to Atlantic salmon, followed by hits against Nile tilapia (*Oreochromis niloticus*), zebrafish (*Danio rerio*), Japanese medaka (*Oryzias latipes*) and other fish species. Gene ontology (GO) annotation of the NR unigenes was obtained with the Blast2GO and WEGO software's (Conesa et al., 2005; Ye et al., 2006). Fig. S2 shows the major GOs from the salmon lens transcriptome, divided into the three ontologies biological process, molecular function and cellular component. Of the more specialized molecular process worth mentioning is the antioxidant activity, while the biological function GO annotation indicate that the lens cells are relatively metabolic active.

4.4. Differentially expressed genes (DEGs)

To search for differentially expressed genes (DEGs) in diploid and triploid salmon with and without cataracts (Fig. 2B), the left lens from 26 individual fish were selected for RNA-seq analysis. The selection was based on cataract score (score 0 vs 3) and ploidy (2N vs 3N). A total of 634,610,512 single-end reads were sequenced with the Illumina HiSeq™ 2000 system. In average, 24,023,481 ± 199,810 single-end reads were sequenced per sample ($n = 26$, mean ± SEM). Average total reads mapped to the in-house made lens transcriptome were 17,882,297 ± 161,561 ($n = 26$, mean ± SEM), representing 74.4% of the total reads. As expected for fish, some contigs had redundant annotations.

Using the default NOISeq setting for calculation of differential expression ($q \geq 0.8$ and $\log 2 \geq 1$), the comparison between diploid fish without and with cataracts (2N- vs 2N+) showed that 182 DEGs were more highly expressed in 2N- lenses and 25 DEGs were more highly expressed in 2N+ lenses (Fig. 2C). For the comparison between triploid fish without and with cataracts (3N- vs 3N+), 74 DEGs were more highly expressed in 3N- and 78 DEGs were more highly expressed in 3N+. Comparison of healthy diploid lenses vs healthy triploid lenses (2N- vs 3N-) yielded 107 DEGs, with 93 genes more highly expressed in 2N-, and 14 more highly expressed in 3N-. Comparison of cataractous diploid lenses vs cataractous triploid lenses (2N+ vs 3N+) yielded 165 significant DEGs, with 9 genes more highly expressed in 2N+, and 156 genes more highly expressed in 3N+. All significant DEGs in the four comparisons, including fold changes, significance levels and best annotation, which were used in downstream functional analyses, are shown in Supplementary File 2. Annotations were given to about 52% of the DEGs.

Very few DEGs with unique expression were found in the lenses from the four treatment groups. Fig. 2D shows a Venn diagram of the number of unique and shared DEGs determined with a four-way comparison. There were 4 unique DEGs in the 2N- group, 17 in the 2N+ group, 15 in the 3N- group and 245 in the 3N+ group. 98.7% of the DEGs were shared between all treatment groups. Most of these unique DEGs were expressed only in one or a few of the lenses from their respective group. Annotations of unique DEGs are shown in Supplementary File 3.

4.5. Functional analysis

Two pathway analysis methods, KEGG and IPA pathway analysis, were employed for functional analysis of DEGs in cataractous lenses from diploid and triploid fish. KEGG pathway enrichment analysis identifies significantly enriched metabolic pathways or signal transduction pathways in DEGs by comparison to the whole genome. Table 1 shows the most significant KEGG pathways from the four comparisons based on a q -value cut-off of 0.05. The top three pathways in both diploid and triploid fish with cataracts were “Phototransduction”, “Carbohydrate digestion and absorption” and “Proximal tubule

bicarbonate reclamation”. Interestingly, for the phototransduction pathway (KEGG pathway ko04744), the significant DEGs linked to this system, 13 DEGs in the diploid fish and 17 DEGs in the triploid fish, (DEGs only found in triploid fish were *gnb1*, *arrb1* and *arr3*), were all up-regulated in the diploid cataractous lens (Fig. 3A) and down-regulated in the triploid cataractous lens (Fig. 3B). As expected, direct comparisons between diploid and triploid lens from fish without and with cataract gives similar patterns. The “ECM-receptor interaction” and “PPAR signaling” pathways were two the most significantly affected KEGG entries based on a direct comparison of DEGs in diploid and triploid salmon with cataracts not listed in the other comparisons.

IPA Core analysis and the IPA Compare function were used for evaluation of biological processes, pathways and networks. In order to use IPA, all identifiers must be recognized as mammalian homologs. Some fish-specific genes obviously cannot be given human ortholog names recognized by IPA, and thus were omitted from the IPA-Core analysis. About 52% of the DEGs from the four gene lists were given automatic annotation as described above (Supplementary File 2). In addition, all unknown DEGs were manually aligned against the core nucleotide and EST databases, and given annotation based on hits against NCBI Unigene entries (Blastn cut-off E10-5). This way, 64.4% of the DEGs used for the functional analysis had IPA identifiers. Table 2 shows annotated salmon genes with human identifiers used in these functional analyses which were significantly differently expressed according to the four comparisons (2N+ vs 2N-, 3N+ vs. 3N-, 3N+ vs 2N+ and 3N- vs 2N-). Highlighted in the table are cataract-linked genes that are differentially regulated in various mice knockout models (data obtained from the iSyTE (integrated Systems Tool for Eye gene discovery) database (URL: <http://research.bioinformatics.udel.edu/iSyTE>).

4.6. Impact of cataracts

To get an idea of the mechanistic basis for cataract development in the salmon lens and the impact of ploidy, we used IPA Core Analysis with the predicted upstream regulators function and the categorical annotations of disease or function to search for differences in the four comparisons described above. By sorting with an activation z-score >2 and p -value of overlap <10.5, IPA Core Analysis predicted six upstream regulators that may explain the observed DEGs in lenses of diploid salmon with cataracts. These were CRX, GTF2IRD1, HIF1A, EDN1, hexachlorobenzene and EPO (Supplementary File 4). The dataset for the most significant transcriptional regulator, CRX with a z-score of 2.43 and a p -value of overlap of 8,35E-19, was made up of the DEGs *arr3*, *gnat1*, *gnat2*, *opn1lw*, *opn1sw*, *pdg*, *pde6g*, *prph2*, *rcvrn* and *rho*. For GTF2IRD1, which had a z-score of 2.10 and p -value of overlap of 2,89E-10, the dataset consisted of *arr3*, *gnat1*, *gnat2*, *opn1lw*, *opn1sw*, *pdg* and *rho*. For the disease of functional annotation, the analysis predicted eight categories with a z-score above 2 and p -value >10.5. These were “Cellular Movement, Immune Cell Trafficking-leukocyte migration”, “Cellular Movement-cell movement”, “Cellular Movement, Hematological System Development and Function, Immune Cell Trafficking-cell movement of leukocytes”, “Cellular Movement-migration of cells”, “Cell Death and Survival-cell viability”, “Cell Death and Survival-cell survival”, “Tissue Morphology-quantity of cells” and “Cellular Movement-migration of brain cancer cell lines”.

In the lenses of triploid salmon with cataracts, five upstream regulators had a predicted activation state based on the same cut-off as described above. These were CRX, GTF2IRD1, beta-estradiol, trichostatin A and decitabine (Supplementary File 5). CRX, the most significantly regulator with a z-score of -2.73 and a p -value of overlap of 8,90E-20, was predicted affected based on the same DEGs as in diploid fish, i.e. *arr3*, *gnat1*, *gnat2*, *opn1lw*, *opn1sw*, *pdg*, *pde6g*, *prph2*, *rcvrn* and *rho*. The predicted activation state for GTF2IRD1 (z-score: -2.10, p -value of overlap: 6,33E-11) in lenses of triploid salmon was based on the same DEGs as in diploid fish. Using the same cut-off, no disease or function categories had a predicted activation state in the

Table 1
KEGG pathway enrichment analysis of DEGs.

| # | Treatment comparison/Pathway | DEGs with pathway annotation | All genes with pathway annotation | p-value | Q-value | Pathway ID |
|----|-----------------------------------------------------------|------------------------------|-----------------------------------|--------------|--------------|------------|
| | 2N- vs 2N+ | 72 | 11680 | | | |
| 1 | Phototransduction | 13 (18.06%) | 67 (0.57%) | 1.161234e-16 | 9.638242e-15 | ko04744 |
| 2 | Carbohydrate digestion and absorption | 8 (11.11%) | 39 (0.33%) | 7.384585e-11 | 3.064603e-09 | ko04973 |
| 3 | Proximal tubule bicarbonate reclamation | 7 (9.72%) | 27 (0.23%) | 2.019998e-10 | 5.588661e-09 | ko04964 |
| 4 | Mineral absorption | 6 (8.33%) | 48 (0.41%) | 4.43982e-07 | 7.370101e-06 | ko04978 |
| 5 | Glutamatergic synapse | 7 (9.72%) | 95 (0.81%) | 1.803348e-06 | 1.663088e-05 | ko04724 |
| 6 | Aldosterone-regulated sodium reabsorption | 5 (6.94%) | 35 (0.3%) | 2.173693e-06 | 1.804165e-05 | ko04960 |
| 7 | Chemokine signaling pathway | 9 (12.5%) | 231 (1.98%) | 1.141543e-05 | 7.895672e-05 | ko04062 |
| 8 | Dopaminergic synapse | 8 (11.11%) | 178 (1.52%) | 1.295758e-05 | 8.272916e-05 | ko04728 |
| 9 | GABAergic synapse | 6 (8.33%) | 86 (0.74%) | 1.414778e-05 | 8.387612e-05 | ko04727 |
| 10 | Endocrine and other factor-regulated calcium reabsorption | 5 (6.94%) | 70 (0.6%) | 6.855952e-05 | 3.793627e-04 | ko04961 |
| 11 | Retrograde endocannabinoid signaling | 5 (6.94%) | 77 (0.66%) | 0.0001082454 | 5.615230e-04 | ko04723 |
| 12 | Cholinergic synapse | 5 (6.94%) | 86 (0.74%) | 0.0001827854 | 8.924228e-04 | ko04725 |
| 13 | Serotonergic synapse | 5 (6.94%) | 92 (0.79%) | 0.0002508305 | 1.095733e-03 | ko04726 |
| 14 | Protein digestion and absorption | 5 (6.94%) | 122 (1.04%) | 0.0009166398 | 3.732900e-03 | ko04974 |
| 15 | Cardiac muscle contraction | 5 (6.94%) | 140 (1.2%) | 0.001692799 | 6.108796e-03 | ko04260 |
| 16 | MAPK signaling pathway | 6 (8.33%) | 302 (2.59%) | 0.01064676 | 3.398773e-02 | ko04010 |
| 17 | Endocytosis | 6 (8.33%) | 324 (2.77%) | 0.01466203 | 4.507217e-02 | ko04144 |
| 1 | 3N- vs 3N+ Proximal tubule bicarbonate reclamation | 63 7 (11.11%) | 11680 27 (0.23%) | 7.690174e-11 | 5.998336e-09 | ko04964 |
| 2 | Carbohydrate digestion and absorption | 6 (9.52%) | 39 (0.33%) | 5.481106e-08 | 1.654731e-06 | ko04973 |
| 3 | Phototransduction | 7 (11.11%) | 67 (0.57%) | 6.364351e-08 | 1.654731e-06 | ko04744 |
| 4 | Mineral absorption | 6 (9.52%) | 48 (0.41%) | 1.985252e-07 | 3.871241e-06 | ko04978 |
| 5 | Aldosterone-regulated sodium reabsorption | 5 (7.94%) | 35 (0.3%) | 1.113271e-06 | 1.010844e-05 | ko04960 |
| 6 | GABAergic synapse | 6 (9.52%) | 86 (0.74%) | 6.486509e-06 | 4.599525e-05 | ko04727 |
| 7 | Glutamatergic synapse | 6 (9.52%) | 95 (0.81%) | 1.154713e-05 | 7.505634e-05 | ko04724 |
| 8 | Endocrine and other factor-regulated calcium reabsorption | 5 (7.94%) | 70 (0.6%) | 3.590785e-05 | 2.154471e-04 | ko04961 |
| 9 | Protein digestion and absorption | 6 (9.52%) | 122 (1.04%) | 4.796446e-05 | 2.672306e-04 | ko04974 |
| 10 | Retrograde endocannabinoid signaling | 5 (7.94%) | 77 (0.66%) | 5.694637e-05 | 2.961211e-04 | ko04723 |
| 11 | Cholinergic synapse | 5 (7.94%) | 86 (0.74%) | 9.671235e-05 | 4.714727e-04 | ko04725 |
| 12 | Serotonergic synapse | 5 (7.94%) | 92 (0.79%) | 0.0001332214 | 6.112511e-04 | ko04726 |
| 13 | Chemokine signaling pathway | 7 (11.11%) | 231 (1.98%) | 0.0002331504 | 1.010318e-03 | ko04062 |
| 14 | Dopaminergic synapse | 6 (9.52%) | 178 (1.52%) | 0.0003808976 | 1.555533e-03 | ko04728 |
| 15 | Arginine and proline metabolism | 4 (6.35%) | 67 (0.57%) | 0.00045676 | 1.696537e-03 | ko00330 |
| 16 | Cardiac muscle contraction | 5 (7.94%) | 140 (1.2%) | 0.0009267016 | 3.142727e-03 | ko04260 |
| 17 | PPAR signaling pathway | 3 (4.76%) | 90 (0.77%) | 0.01259403 | 3.929337e-02 | ko03320 |
| 1 | 2N- vs 3N- Phototransduction | 45 9 (20%) | 11680 67 (0.57%) | 2.900818e-12 | 1.885532e-10 | ko04744 |

(continued on next page)

Table 1 (continued)

| # | Treatment comparison/Pathway | DEGs with pathway annotation | All genes with pathway annotation | p-value | Q-value | Pathway ID |
|----|-----------------------------------------------------------|------------------------------|-----------------------------------|--------------|--------------|------------|
| | 2N- vs 2N+ | 72 | 11680 | | | |
| 2 | Carbohydrate digestion and absorption | 7 (15.56%) | 39 (0.33%) | 1.084661e-10 | 3.525148e-09 | ko04973 |
| 3 | Proximal tubule bicarbonate reclamation | 6 (13.33%) | 27 (0.23%) | 6.445801e-10 | 1.396590e-08 | ko04964 |
| 4 | Aldosterone-regulated sodium reabsorption | 5 (11.11%) | 35 (0.3%) | 2.011263e-07 | 2.614642e-06 | ko04960 |
| 5 | Mineral absorption | 5 (11.11%) | 48 (0.41%) | 1.022166e-06 | 9.305999e-06 | ko04978 |
| 6 | Serotonergic synapse | 6 (13.33%) | 92 (0.79%) | 1.288523e-06 | 9.305999e-06 | ko04726 |
| 7 | Chemokine signaling pathway | 8 (17.78%) | 231 (1.98%) | 2.378076e-06 | 1.405227e-05 | ko04062 |
| 8 | Protein digestion and absorption | 6 (13.33%) | 122 (1.04%) | 6.701316e-06 | 3.150043e-05 | ko04974 |
| 9 | Endocrine and other factor-regulated calcium reabsorption | 5 (11.11%) | 70 (0.6%) | 6.784707e-06 | 3.150043e-05 | ko04961 |
| 10 | Retrograde endocannabinoid signaling | 5 (11.11%) | 77 (0.66%) | 1.085652e-05 | 4.704492e-05 | ko04723 |
| 11 | Cholinergic synapse | 5 (11.11%) | 86 (0.74%) | 1.865061e-05 | 7.131116e-05 | ko04725 |
| 12 | GABAergic synapse | 5 (11.11%) | 86 (0.74%) | 1.865061e-05 | 7.131116e-05 | ko04727 |
| 13 | Glutamatergic synapse | 5 (11.11%) | 95 (0.81%) | 3.024149e-05 | 1.092054e-04 | ko04724 |
| 14 | Dopaminergic synapse | 6 (13.33%) | 178 (1.52%) | 5.726626e-05 | 1.959109e-04 | ko04728 |
| 15 | Cardiac muscle contraction | 5 (11.11%) | 140 (1.2%) | 0.0001913906 | 5.923995e-04 | ko04260 |
| 16 | Retinol metabolism | 2 (4.44%) | 19 (0.16%) | 0.002380689 | 6.447699e-03 | ko00830 |
| 17 | Neuroactive ligand-receptor interaction | 2 (4.44%) | 52 (0.45%) | 0.01703389 | 4.258472e-02 | ko04080 |
| 1 | 2N + vs 3N+ Phototransduction | 68 14 (20.59%) | 11680 67 (0.57%) | 9.57472e-19 | 5.170349e-17 | ko04744 |
| 2 | Protein digestion and absorption | 16 (23.53%) | 122 (1.04%) | 6.816708e-18 | 1.840511e-16 | ko04974 |
| 3 | Carbohydrate digestion and absorption | 8 (11.76%) | 39 (0.33%) | 4.604135e-11 | 6.215582e-10 | ko04973 |
| 4 | Proximal tubule bicarbonate reclamation | 7 (10.29%) | 27 (0.23%) | 1.337376e-10 | 1.444366e-09 | ko04964 |
| 5 | Dopaminergic synapse | 9 (13.24%) | 178 (1.52%) | 8.241675e-07 | 5.454431e-06 | ko04728 |
| 6 | Aldosterone-regulated sodium reabsorption | 5 (7.35%) | 35 (0.3%) | 1.633423e-06 | 8.820484e-06 | ko04960 |
| 7 | ECM-receptor interaction | 8 (11.76%) | 146 (1.25%) | 1.929870e-06 | 8.874621e-06 | ko04512 |
| 8 | Chemokine signaling pathway | 9 (13.24%) | 231 (1.98%) | 7.078719e-06 | 2.940391e-05 | ko04062 |
| 9 | Mineral absorption | 5 (7.35%) | 48 (0.41%) | 8.12637e-06 | 3.134457e-05 | ko04978 |
| 10 | GABAergic synapse | 6 (8.82%) | 86 (0.74%) | 1.014675e-05 | 3.652830e-05 | ko04727 |
| 11 | Glutamatergic synapse | 6 (8.82%) | 95 (0.81%) | 1.800377e-05 | 6.076272e-05 | ko04724 |
| 12 | Endocrine and other factor-regulated calcium reabsorption | 5 (7.35%) | 70 (0.6%) | 5.203389e-05 | 1.652841e-04 | ko04961 |
| 13 | Retrograde endocannabinoid signaling | 5 (7.35%) | 77 (0.66%) | 8.231658e-05 | 2.339524e-04 | ko04723 |
| 14 | Cholinergic synapse | 5 (7.35%) | 86 (0.74%) | 0.000139355 | 3.762585e-04 | ko04725 |
| 15 | Serotonergic synapse | 5 (7.35%) | 92 (0.79%) | 0.0001915559 | 4.925723e-04 | ko04726 |
| 16 | Arginine and proline metabolism | 4 (5.88%) | 67 (0.57%) | 0.0006112079 | 1.500238e-03 | ko00330 |
| 17 | Cardiac muscle contraction | 5 (7.35%) | 140 (1.2%) | 0.001310250 | 2.948062e-03 | ko04260 |
| 18 | PPAR signaling pathway | 4 (5.88%) | 90 (0.77%) | 0.001843347 | 3.828490e-03 | ko03320 |
| 19 | Focal adhesion | 8 (11.76%) | 418 (3.58%) | 0.002838214 | 5.676428e-03 | ko04510 |




Fig. 3. Six DEGs encoding protein involved in phototransduction (KEGG pathway ko04744) were up-regulated in cataractous lenses from diploid (2N+) Atlantic salmon (A) and down-regulated in cataractous lenses from triploid (3N+) Atlantic salmon (B). Border color indicates up-regulated genes (red) and down-regulated genes (green). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

lenses of triploid salmon.

By comparing the transcriptional patterns in cataractous lenses of diploid and triploid salmon indirectly (IPA Compare Analysis of 2N- versus 2N+ and 3N- versus 3N+), the most pronounced differences were seen for the “IL8” and “Production of nitric oxide and reactive oxygen species in macrophage” canonical pathways (data not shown). These pathways had higher activation z-scores in lenses of the diploid fish compared to the triploid fish. A predicted regulator network generated from the comparison of DEGs in cataractous lenses of diploid and triploid salmon is shown in Fig. 4. This network, which had a consistency score of 13.87 and was based on target DEGs *apoe*, *clu*, *gal*, *igfbp2*, *junb*, *krt18*, *lep*, *mmp2*, *plaur*, *rbp4* and *snca*, and on upstream regulators AGT, CREB1, ERK, HIF1A and P38 MAPK, predicted that synthesis of nitric oxide and chemotaxis of cells might be different in cataractous lenses of diploid and triploid salmon. Based on analysis of predicted upstream regulators and hierarchical clustering, the most pronounced difference in triploid fish was seen for CRX, GTF2IRD1, SRC and RHO (Table 3). Interestingly, these upstream regulators were predicted activated in diploid fish (positive z-score) and predicted inhibited in triploid fish (negative z-score) with cataracts. Fig. 5 shows the molecules in these four networks. Except *krt18* and *ckm* in the SRC network, all genes in these networks were up-regulated by cataracts in diploid fish and down-regulated in triploid fish.

4.7. Impact of ploidy

Comparison of the transcriptional patterns in lenses from diploid and triploid salmon without cataracts revealed two upstream regulators with predicted activation scores above 2 and *p*-values of overlap >10⁻⁴ (Supplementary File 6). According to the IPA Core Analysis, both the transcription regulator CRX and the chemical drug trichostatin A were predicted activated with z-scores of 2.55 and 2.40, respectively. Targeted DEGs for CRX were *gnat1*, *gnat2*, *opn1lw*, *opn1sw*, *pdg*, *pde6g*, *prph2*, *rcvrm* and *rho*, while *cdh1*, *hba1/hba2*, *hbb*, *hbx*, *ndrg1* and *slc1a2* made up the dataset for the trichostatin A prediction.

A comparison of transcriptional patterns in lenses from diploid and triploid fish without cataracts showed that two disease or function annotations had prediction scores above 2 and a *p*-value of overlap >10⁻⁵ (Supplementary File 6). “Organ degeneration” (z-score 2.19) and “Degeneration of cells” (z-score 2.19) showed significant differential prediction scores in lenses of triploid and diploid salmon without cataracts. The “Organ degeneration” z-score was based on differential transcription of *prph2*, *rho*, *crb1*, *slc1a2*, *pde6g*, *gngt1*, *rpe65*, *rcvrm*, *ca1*, *gnat1*, *opn1mw* (includes others) and *ca2*, whereas the “Degeneration of cells” z-score was based on differential transcription of *rho*, *slc1a2*, *pde6g*, *gngt1*, *rpe65*, *gnat1* and *prph2*. All these genes were more highly expressed in lenses of diploid fish compared to triploid fish. This could reflect differential transcription in non-cataractous lenses from diploid and triploid salmon, or indicate that mechanisms leading to cataracts

Table 2

Annotated salmon genes with human orthologs used in the IPA functional analyses. Genes common for all four comparisons were all up-regulated in diploid cataractous lenses and down-regulated in triploid cataractous lenses. Genes in bold are differentially regulated in mice cataract mutants (>2.0 fold) according to the iSyTE database (Kakrana et al., 2018). Underlined are genes that are differentially regulated by more than one mutant type.

| Comparison | Genes |
|---------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 2N + vs 2N- (diploid cataract) | <u>ada</u> , <u>agbl4</u> , <u>anxa2</u> , <u>anxa5</u> , <u>apoc1</u> , <u>apoe</u> , <u>arl14ep</u> , <u>arr3</u> , <u>arrb2</u> , <u>atp1a3</u> , <u>atp1a4</u> , <u>aurka</u> , <u>btbd17</u> , <u>c18orf25</u> , <u>c4orf33</u> , <u>ca1</u> , <u>ca2</u> , <u>cars</u> , <u>ccl28</u> , <u>cdca8</u> , <u>ckmt2</u> , <u>clu</u> , <u>col11a1</u> , <u>cyb5r1</u> , <u>depor</u> , <u>dnajc3</u> , <u>egfl7</u> , <u>f11r</u> , <u>gal</u> , <u>glul</u> , <u>gnat1</u> , <u>gnat2</u> , <u>gnb1</u> , <u>gnb3</u> , <u>gnb5</u> , <u>nggt1</u> , <u>nggt2</u> , <u>hamp</u> , <u>harbi1</u> , <u>hba1</u> , <u>hbb</u> , <u>hzb</u> , <u>homer2</u> , <u>hspa8</u> , <u>ifitm5</u> , <u>igfbp2</u> , <u>junb</u> , <u>krt18</u> , <u>ldhb</u> , <u>lep</u> , <u>linc00998</u> , <u>lurap1</u> , <u>lzt1</u> , <u>mlkl</u> , <u>mlt11</u> , <u>mmp2</u> , <u>ndrg1</u> , <u>ndrg4</u> , <u>nmt1</u> , <u>opn1lw</u> , <u>opn1mw2</u> , <u>opn1sw</u> , <u>parp15</u> , <u>pdac</u> , <u>pde6g</u> , <u>plaur</u> , <u>ppdpf</u> , <u>ppp2ca</u> , <u>prdm9</u> , <u>prph2</u> , <u>rab32</u> , <u>rasd2</u> , <u>rbp4</u> , <u>rcvrn</u> , <u>rho</u> , <u>rpe65</u> , <u>rtbdn</u> , <u>s100a1</u> , <u>s100p</u> , <u>slc27a2</u> , <u>smarcd1</u> , <u>snc</u> , <u>specc1</u> , <u>stc1</u> , <u>tmsb10</u> , <u>tp52l1</u> , <u>trpm7</u> , <u>wpp2</u> , <u>znf391</u> , <u>znf501</u> |
| 3N + vs 3N- (triploid cataract) | <u>anxa2</u> , <u>anxa5</u> , <u>arl14ep</u> , <u>arrb2</u> , <u>atp1a3</u> , <u>atp1a4</u> , <u>btbd17</u> , <u>ca1</u> , <u>ca2</u> , <u>cars</u> , <u>ccl28</u> , <u>ckb</u> , <u>bckm</u> , <u>ckmt2</u> , <u>clu</u> , <u>cdnp2</u> , <u>ctnnap5</u> , <u>dnal</u> , <u>defb4b</u> , <u>erbb2ip</u> , <u>f11r</u> , <u>fabp4</u> , <u>fabp7</u> , <u>gal</u> , <u>gfap</u> , <u>glul</u> , <u>gnat1</u> , <u>gnat2</u> , <u>gnb3</u> , <u>nggt1</u> , <u>nggt2</u> , <u>hamp</u> , <u>harbi1</u> , <u>ifitm5</u> , <u>igfbp2</u> , <u>junb</u> , <u>krt18</u> , <u>krt8</u> , <u>lcn1</u> , <u>ldhb</u> , <u>lep</u> , <u>lgals3</u> , <u>lmna</u> , <u>mlkl</u> , <u>mxra7</u> , <u>ndrg1</u> , <u>nfyb</u> , <u>nmt1</u> , <u>oaz2</u> , <u>opn1lw</u> , <u>opn1mw2</u> , <u>opn1sw</u> , <u>pdac</u> , <u>pde6g</u> , <u>plaur</u> , <u>prdm9</u> , <u>prph2</u> , <u>rcvrn</u> , <u>rho</u> , <u>rpe65</u> , <u>rtbdn</u> , <u>s100a1</u> , <u>s100p</u> , <u>serf2</u> , <u>slc27a2</u> , <u>snc</u> , <u>tmsb10</u> , <u>trpm7</u> , <u>vps13d</u> , <u>znf391</u> , <u>znf501</u> |
| 3N + vs 2N+ | <u>actg2</u> , <u>aebp1</u> , <u>alpl</u> , <u>apoe</u> , <u>arr3</u> , <u>arrb2</u> , <u>atp1a1</u> , <u>atp1a3</u> , <u>atp1a4</u> , <u>btbd17</u> , <u>ca1</u> , <u>ca2</u> , <u>ctgf</u> , <u>ckb</u> , <u>ckm</u> , <u>ckmt2</u> , <u>cdnp2</u> , <u>ctnnap5</u> , <u>coch</u> , <u>col2a1</u> , <u>col9a1</u> , <u>col9a2</u> , <u>dnase1l3</u> , <u>eef1a1</u> , <u>fabp4</u> , <u>fabp7</u> , <u>gja1</u> , <u>glul</u> , <u>gnat1</u> , <u>gnat2</u> , <u>gnb1</u> , <u>gnb3</u> , <u>gnb5</u> , <u>nggt1</u> , <u>nggt2</u> , <u>hamp</u> , <u>harbi1</u> , <u>igsf11</u> , <u>lcn1</u> , <u>ldhb</u> , <u>lrrn1</u> , <u>mgst3</u> , <u>nbl1</u> , <u>ndrg1</u> , <u>ndufa4</u> , <u>opn1lw</u> , <u>opn1mw2</u> , <u>opn1sw</u> , <u>pdac</u> , <u>pde6g</u> , <u>pkm</u> , <u>plaur</u> , <u>ppdpf</u> , <u>prph2</u> , <u>ptma</u> , <u>pygm</u> , <u>rbp4</u> , <u>rcvrn</u> , <u>rgmb</u> , <u>rho</u> , <u>rpe65</u> , <u>rtbdn</u> , <u>serpinh1</u> , <u>sirt1</u> , <u>slc39a11</u> , <u>snc</u> , <u>sparc</u> , <u>spon1</u> , <u>stmn1</u> , <u>suclg1</u> , <u>trpm7</u> , <u>ube2j1</u> |
| 3N- vs 2N- | <u>arrb2</u> , <u>atp1a3</u> , <u>atp1a4</u> , <u>btbd17</u> , <u>ca1</u> , <u>ca2</u> , <u>cdh1</u> , <u>ckm</u> , <u>ckmt2</u> , <u>crb1</u> , <u>crygs</u> , <u>cyp2c19</u> , <u>erbb2ip</u> , <u>exosc2</u> , <u>glra1</u> , <u>glul</u> , <u>gnat1</u> , <u>gnat2</u> , <u>gnb3</u> , <u>nggt1</u> , <u>nggt2</u> , <u>harbi1</u> , <u>hba1</u> , <u>hbb</u> , <u>hzb</u> , <u>igsf11</u> , <u>ldhb</u> , <u>mvp</u> , <u>ndrg1</u> , <u>opn1lw</u> , <u>opn1mw2</u> , <u>opn1sw</u> , <u>pdac</u> , <u>pde6g</u> , <u>plaur</u> , <u>polr3c</u> , <u>prph2</u> , <u>rbp4</u> , <u>rcvrn</u> , <u>rho</u> , <u>rpe65</u> , <u>rps12</u> , <u>rtbdn</u> , <u>slc1a2</u> , <u>stc1</u> , <u>suclg1</u> , <u>trpm7</u> |
| Common in 2N+ and 3N+ | <u>anxa2</u> , <u>anxa5</u> , <u>arl14ep</u> , <u>arrb2</u> , <u>atp1a3</u> , <u>atp1a4</u> , <u>btbd17</u> , <u>ca1</u> , <u>ca2</u> , <u>cars</u> , <u>ccl28</u> , <u>ckmt2</u> , <u>clu</u> , <u>f11r</u> , <u>gal</u> , <u>glul</u> , <u>gnat1</u> , <u>gnat2</u> , <u>gnb3</u> , <u>nggt1</u> , <u>nggt2</u> , <u>hamp</u> , <u>harbi1</u> , <u>ifitm5</u> , <u>igfbp2</u> , <u>junb</u> , <u>krt18</u> , <u>ldhb</u> , <u>lep</u> , <u>mlkl</u> , <u>ndrg1</u> , <u>nmt1</u> , <u>opn1lw</u> , <u>opn1mw2</u> , <u>opn1sw</u> , <u>pdac</u> , <u>pde6g</u> , <u>plaur</u> , <u>prdm9</u> , <u>prph2</u> , <u>rcvrn</u> , <u>rho</u> , <u>rpe65</u> , <u>rtbdn</u> , <u>s100a1</u> , <u>s100p</u> , <u>slc27a2</u> , <u>snc</u> , <u>tmsb10</u> , <u>trpm7</u> , <u>znf391</u> , <u>znf501</u> |
| Common for all 4 comparisons | <u>arrb2</u> , <u>atp1a3</u> , <u>atp1a4</u> , <u>btbd17</u> , <u>ca1</u> , <u>ca2</u> , <u>ckmt2</u> , <u>glul</u> , <u>gnat1</u> , <u>gnat2</u> , <u>gnb3</u> , <u>nggt1</u> , <u>nggt2</u> , <u>harbi1</u> , <u>ldhb</u> , <u>ndrg1</u> , <u>opn1lw</u> , <u>opn1mw2</u> , <u>opn1sw</u> , <u>pdac</u> , <u>pde6g</u> , <u>plaur</u> , <u>prph2</u> , <u>rcvrn</u> , <u>rho</u> , <u>rpe65</u> , <u>rtbdn</u> , <u>trpm7</u> |

may be affected in some of the lenses even without visible signs of damage.

The direct comparison of transcriptional patterns in lenses of diploid and triploid salmon with cataracts yielded five predicted upstream regulators with activation z-score >2 and p-value of overlap >10⁻⁵ (Supplementary File 7). These were CRX, GTF2IRD1, beta-estradiol, trichostatin A and decitabine. Targeted DEGs in the most significant regulator (transcription regulator CRX, p-value of overlap 8,90E-20) were *arr3*, *gnat1*, *gnat2*, *opn1lw*, *opn1sw*, *pdac*, *pde6g*, *prph2*, *rcvrn* and *rho*. A significant result for the transcription regulator GTF2IRD1 was based on the DEGs *arr3*, *gnat1*, *gnat2*, *opn1lw*, *opn1sw*, *pdac* and *rho*. Five categories with disease or functional annotation had predicted activation state based on z-score >2 and p-value of overlap >10⁻⁵ (Supplementary File 7). These were “DNA Replication, Recombination, and Repair, Nucleic Acid Metabolism, Small Molecule Biochemistry-hydrolysis of nucleotide”, “Behavior-”, “Cellular-Movement-migration of blood cells”, “Cellular Movement-cell movement” and “Organismal Development-size of body”.

5. Discussion

This is the first study to investigate the transcriptomics of salmon lenses in diploid and triploid salmon with and without cataracts. Functional analysis showed that retina-associated genes were differentially affected in diploid and triploid fish. Predicted differential effects of NO-induced oxidative stress, modified cytoskeleton stability and lipid metabolism, possibly affecting cellular metabolism, indicate that the triploid lens might be more vulnerable to cataract due to altered protein degradation and turnover.

Overall, this study indicates that the transcriptional patterns in the lenses of diploid and triploid Atlantic salmon are very similar. This is consistent with the results from a recent study, which showed that the vast majority of genes in liver tissue had similar expression levels between diploid and triploid coho salmon (*Oncorhynchus kisutch*) (Christensen et al., 2019). Similar results have been shown for other fish species (Chatchaiphan et al., 2017). At the protein level there also appears to be small differences in expression between diploid and triploid salmon (Nuez-Ortin et al., 2017). Relatively few significant DEGs were found in the current dataset. Most of the significant DEGs in the cataractous triploid lenses were higher expressed compared to the cataractous diploid lens (156 vs 9). In healthy lenses the pattern was opposite, with more of the significant DEGs being lower expressed in the diploid lenses (93 vs 14). According to the functional analysis, the most distinct difference between diploid and triploid cataractous lenses in transcript levels were seen for genes encoding proteins involved in the photo-transduction pathway. Whether this reflects a direct effect of ploidy on the transcription of these genes is unknown.

The N3+ vs N2+ comparison list contained a gene associated with heat shock protein (HSP) activity, e.g. *hsp47/serpinh1*. Furthermore, two heat shock protein genes, annotated to *hspa8* and *hspa8b*, were up-regulated in diploid cataractous lenses but not in diploid non-cataractous lenses. These findings potentially suggest a different ability to handle damaged proteins and protein turnover. Crystallins, water-soluble structural protein found in the lens and the cornea of the eye accounting for the transparency, are relatively similar to HSPs, and have similar chaperone activity (Wang and Spector, 1995; Slingsby et al., 2013). “Protein digestion and absorption” (KEGG pathway ko04974) was the second most significantly affected pathway, after photo-transduction, in triploid cataractous lenses compared to diploid cataractous lenses, according to the functional analysis. In humans, ROS-generated protein oxidation may lead to cataract formation in the aged lens (Taylor and Davies, 1987). In addition to oxidative stress and the inflammatory response, an unfolded protein response is known to be activated in age-related ocular disorders such as cataracts (Lenox et al., 2015). Histidine has been shown to stimulate the proteasome and thereby protein degradation and turnover (Hamel et al., 2003; Breck et al., 2005a). With diminished antioxidant capacity and decreased proteolytic capabilities, the triploid lens may be less efficient in clearance of damaged proteins. Taken together, the results from the current study indicate that the higher susceptibility to cataract development in triploid vs diploid salmon may in part rely on how well the cells handle damaged proteins.

The heart histidine and NAH levels observed in the present study represent normal values obtained from a commercial salmon smolt feed (Remø et al., 2014). According to the factorial analysis, the histidine concentration was related to cataract status and not to ploidy, while NAH status did not indicate differences with neither cataract status nor ploidy. The result confirms the higher sensitivity of histidine relative to NAH status in heart tissue, but more importantly, a corresponding lens histidine status to the diploid cataract group and both the triploid groups indicates suboptimal conditions for salmon smolts (Remø et al., 2014). The present groups of salmon would therefore be prone to cataract development.

At the molecular level, functional analysis predicted that the upstream regulators cone-rod homeobox (CRX), GTF2I repeat domain-

2N+ vs 3N+




Fig. 4. Predicted upstream regulator network with highest consistency score based on IPA Compare Analysis, generated by comparison of DEGs in cataractous lenses of diploid and triploid Atlantic salmon (2N + vs 3N+). Up-regulated and down-regulated genes are highlighted in red and green, while predicted activated and inhibited regulators and relationships are highlighted in orange and blue, respectively. The color depth is correlated to the fold change. Solid lines show direct interactions between two gene products, while dashed lines represent indirect interactions among genes shown in the network. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Table 3

Comparison of predicted upstream regulators in lenses of diploid and triploid Atlantic salmon. Hierarchical clustering based on z-scores as determined with IPA Compare Analysis.

| Upstream regulator | 2N + vs 2N- | 3N + vs 3N- |
|---------------------|-------------|-------------|
| CRX | 2.73 | -2.55 |
| GTF2IRD1 | 2.10 | -1.85 |
| SRC | 1.70 | -1.73 |
| RHO | 1.89 | -1.63 |
| decitabine | 2.77 | 2.20 |
| IL6 | 2.25 | 3.06 |
| P38 MAPK | 2.11 | 1.44 |
| OSM | 1.23 | 2.36 |
| NFkB (complex) | 1.34 | 1.95 |
| APP | 1.25 | 1.94 |
| hexachlorobenzene | 2.0 | 1.98 |
| cisplatin | 1.77 | 1.86 |
| STAT3 | 1.76 | 2.20 |
| thioacetamide | 1.96 | 2.21 |
| lipopolysaccharide | 2.12 | 2.32 |
| pioglitazone | | -1.92 |
| curcumin | -0.73 | -1.71 |
| U0126 | -1.96 | -1.29 |
| LY294002 | -1.67 | -1.67 |
| N-acetyl-L-cysteine | -1.67 | -0.44 |
| MYC | -1.40 | 0.30 |
| CSF2 | -1.48 | |
| PD98059 | -1.82 | -0.01 |
| sirolimus | -2.0 | |
| ESR2 | -1.94 | 0.15 |
| cyclosporin A | 1.07 | 1.39 |
| nitrofurantoin | 0.65 | 1.48 |
| dihydrotestosterone | 0.78 | 1.26 |
| bucladesine | 1.27 | 1.19 |
| beta-estradiol | 1.41 | 1.15 |

containing 1 (GTF2IRD1), SRC proto-oncogene, non-receptor tyrosine kinase (SRC) and rhodopsin (RHO) could explain the differences observed in lens transcript levels between cataractous diploid and triploid fish. All these upstream regulators were predicted to be activated in cataractous lenses of diploid fish and inhibited in cataractous lenses of triploid fish. Expression of visual pigment-like proteins has been described in extraretinal tissue (Shichida and Yamashita, 2003). Differential expression of retina-associated genes has also been documented in zebrafish (*Danio rerio*) cataractous lens cells (Posner et al., 2019). Since the lenses in the current study were extracted through an incision in the cornea, without contacting retinal tissues, it seems unlikely that such visual pigment-like proteins were derived from retinal contamination. CRX is a photoreceptor-specific transcription factor, which plays a role in the differentiation of photoreceptor cells, controlling the maintenance of normal cone and rod function (GeneCards database). While not directly linked to cataracts, mutations in the gene encoding CRX have been linked to severe dystrophy of the human retina (Weleber et al., 1993). GTF2IRD1 function as a transcription factor under the control of retinoblastoma protein, and may be a transcription regulator involved in cell-cycle progression and skeletal muscle differentiation (GeneCards database). In human stromal cells, vitamin E treatment has been shown to down-regulate GTF2IRD1, suggesting a link to the lens antioxidative defense. GTF2IRD1 also respond to chemical exposure. For example, down-regulation of GTF2IRD1 has been shown in human liver cells after exposure to benzo(a)pyrene (Jennen et al., 2010), while the peroxisome proliferator-activated receptor alpha (PPARα) agonist pirinixic acid increases GTF2IRD1 mRNA levels in mice liver (Sanderson et al., 2008). The SRC proto-oncogene may play a role in cell growth and participates in signaling pathways that control a broad spectrum of biological activities including gene transcription, immune response, cell adhesion, cell cycle progression, apoptosis, migration, and transformation (GeneCards database). Of interest for lens damage, SRC plays an important role in the regulation of cytoskeletal organization through phosphorylation mechanisms (GeneCards database). Recent findings suggest that accumulation of crystallin

2N+ vs 3N+




Fig. 5. IPA Compare Analysis of transcripts differentially regulated in cataractous lenses of triploid salmon compared to diploid salmon (2N+ vs 3N+). The figure shows genes associated with four predicted regulators that were activated in diploid fish but inhibited in triploid fish, based on hierarchical clustering and z-score using Upstream Analysis. A) Cone-rod homeobox (CRX), B) GTF21 repeat domain containing 1 (GTF2IRD1), C) rhodopsin (RHO) and D) SRC proto-oncogene, non-receptor tyrosine kinase (SRC).

proteins, a prerequisite for refractive properties and transparency of the lens, in part is controlled by post-transcriptional mechanisms rather than by differential gene transcription (mRNA synthesis) (Terrell et al., 2015). This may explain why we did not see any differentially expressed crystallin genes in the current study. RHO is a photoreceptor primarily expressed in rod cells in the retina required for image-forming vision at low light intensity (GeneCards database). Of nutrients and essential elements, Zn is known to affect stability and folding of RHO (Stojanovic et al., 2004; Gleim et al., 2009), indicating that Zn imbalance might impact RHO activity in cataractous triploid lenses. Zn deficiency has been linked to cataract development in rainbow trout (Ketola, 1979). The functional implication of differential expression of RHO-associated transcripts in lens cells is unknown, as RHO protein is primarily expressed in rod photoreceptors in the retina. In agreement with the current examination, down-regulation of *rho* has been observed in the cataractous zebrafish lens (Posner et al., 2019). The observed effect on *rho* may be linked to dysregulation of vitamin A1 and A2 in the lens endothelial cells outside the retina (Enright et al., 2015), or alternatively may reflect signaling interactions between the lens and retina through hyaloid capillaries (Dhakal et al., 2015).

Lower activation z-scores for pathways linked to synthesis of nitric

oxide (NO) and chemotaxis in triploid fish, suggests that their lens cells may be more prone to oxidative damage and chemotactic cell movement than their diploid counterparts. It is well known that NO has a role in cataract formation in the mammalian lens (Ito et al., 2001; Ornek et al., 2003; Chamberlain et al., 2008). Oxidative stress is an important factor in the development of cataracts for both animals and humans (Ottonello et al., 2000; Williams, 2006). High concentrations of reactive oxygen species (ROS), produced from both endogenous and exogenous sources, cause oxidative damage to cellular constituents that results in interrupted physiological functions and oxidative stress-associated diseases such as cataracts (Lou, 2003). The decreased protein turnover towards the nucleus makes lenses especially vulnerable to increased ROS production in the epithelial cells (reviewed by Brennan and Kantorow, 2009). In human age-related cataracts, oxidation of membrane proteins has been found to precede the development of cataract formation (Wang and Spector, 1995). The accumulation of oxidized proteins further results in loss of cell function, apoptosis and necrosis (Brennan and Kantorow, 2009). Lens NAH has been suggested to be an important intracellular antioxidant in the salmon lens and may contribute to the cataract mitigating effect of dietary histidine (Remø et al., 2011). The present results suggest a higher susceptibility to oxidative stress in

triploid lenses, which may be hypothesized to be an underlying factor for the higher prevalence of cataracts compared to diploids, as well as the lower lens NAH status observed when reared under similar conditions in the studies by Taylor et al. (2015) and Sambraus et al. (2017).

Likewise, cataract and chemotactic activity have been extensively studied over the years (Rosenbaum et al., 1987; Schneider et al., 2012). Several of the predicted upstream regulators for this network, angiostatin (AGT), CAMP responsive element binding protein 1 (CREB1), extracellular signal-regulated kinase (ERK), HIF1A and P38 mitogen-activated protein kinase (P38 MAPK), have been linked to cataract formation (AGT: Taube et al., 2012; Ji et al., 2015; CREB1: Weng et al., 2008; ERK: Iyengar et al., 2007; HIF1A: Chen et al., 2014; P38 MAPK: Bai et al., 2015). Follow-up studies should look at how NO induce ROS and oxidative stress in the triploid fish lens, as well as the involvement of chemotaxis in the development of cataract in triploid salmon.

Two pathways, “ECM-receptor interaction” and “PPAR signaling pathway”, were among the most significantly affected KEGG entries based on a direct comparison of DEGs in diploid and triploid salmon lenses with cataracts. These were not listed in the other comparisons. The ECM-receptor interaction pathway, including collagen, type II, alpha 1 (*col2a1*) and secreted protein, acidic, cysteine-rich (osteonectin) *sparc*, has been linked to disorders of the eye characterized by early onset cataract (Bradshaw, 2009). SPARC is a key lens- and cataract-associated protein (Shiels et al., 2010; Sousounis and Tsonis, 2012; Terrell et al., 2015). SPARC is important for normal cellular proliferation and differentiation and is involved in maintaining lens transparency as shown for mice (Gilmour et al., 1998) and humans (Yan et al., 2000). SPARC is also one of at least 13 proteins harboring mutations that have been associated with a lens or cataract phenotype in mice but not yet in humans (Shiels et al., 2010). In Atlantic salmon, SPARC was suggested to be an “early” up-regulated marker for cataract development (Tröbe et al., 2009). Lower expression of *sparc* suggests that the cataractous triploid lenses might have impaired circulation of fluids, ions, and small molecules, possibly resulting in depolarized membrane resting voltage as shown in mice (Greiling et al., 2009). Cartilage extracellular matrix (ECM) is composed of type II collagen, fibrous proteins and proteoglycans, hyaluronic acid and chondroitin sulfate (Gao et al., 2014). The finding indicates a differential regulated mechanism linked to cytoskeleton disruption and NO-induced oxidative stress (Gao et al., 2014). Differential regulation of PPARs, which are transcription factors in control of many cellular processes, indicate an effect on lipid metabolism in the lens. An effect on lipid/cholesterol transport, previously reported in age-related cataract in humans (Utheim et al., 2008), is suggested by differential expression of apolipoprotein E (*apoe*). APOE is a major apoprotein that is essential for the normal catabolism of triglyceride-rich lipoprotein constituents (Genecards database), indicating a differential effect on lipoprotein metabolism. *Apoe*, together with *sparc*, was among the differentially regulated genes in cataractous lenses of Atlantic salmon fed a low-histidine diet compared to a high-histidine diet (Tröbe et al., 2009) and had a lower expression level in the lens of Atlantic salmon fed plant oils compared to fish oils (Remø et al., 2011).

Only five genes from the *Cat-Map* gene list, an online chromosome map and reference database for cataract in humans and mice (Shiels et al., 2010), showed overlap with the current gene list of cataractous diploid and triploid lenses from salmon (direct comparison). These were, in addition to *sparc* and *apoe*, *col2a1*, gap junction protein alpha 1, 43 kDa (*gja1*) and retinal pigment epithelium-specific protein 65 kDa (*rpe65*). Gap junction proteins, also called connexins, are constituents of gap junctions, channels specialized in cell-cell contacts that provide direct intracellular communication. They allow passive diffusion of molecules up to 1 kDa, including nutrients, metabolites (glucose), ions and second messengers (Genecards database). They are especially important for nutrition and intercellular communication in the avascular lens (Hejtmancik, 2008). Mutations in gap junction proteins such

as *GJA1*, present in the lens epithelium, have been linked to human cataracts (Hejtmancik, 2008). RPE65 is a protein located in the retinal pigment epithelium and involved in the production of 11-cis retinal and in visual pigment regeneration (Genecards database). Finally, RPE65 has been associated with leber congenital amaurosis (LCA), a severe dystrophy of the retina (Weleber et al., 1993). No genes associated with Mendelian (inherited) cataracts or cataracts caused by mutations in transcription factors or metabolic enzymes in humans (Shiels and Hejtmancik, 2019) were on the significant lists in this study. Several of the genes that were differentially expressed in cataractous lenses of triploid salmon have previously been documented to be affected by mutations in the mouse lens (Table 2). By comparing our significant genes with the responses of mammalian orthologs with lens defects or cataract as listed in the iSyTE database (Kakrana et al., 2018), it appears that several may be potential candidate markers for follow-up studies in salmon. Apart from the CBP:p300 E9.5 mutation, which seems to down-regulate many of these genes in mice (iSyTE database), several gene knockout mutation types impact the expression of genes from our lists.

Hsp47, also called *serpinh1*, was one of the genes that were lower expressed in cataractous triploid lenses than in cataractous diploid lenses. HSP47, localized in the endoplasmic reticulum, plays a role in collagen biosynthesis as a collagen-specific molecular chaperone (Genecards database). Heat shock proteins, found throughout the various tissues of the eye, protect and maintain cell viability under stressful conditions such as those occurring during thermal and oxidative challenges chiefly by refolding and stabilizing proteins (Urbak and Vorum, 2010). In the human eye, HSP47 has been suggested to aid the control of pro-collagen under stressful conditions and is induced by corneal structure damage (Urbak and Vorum, 2010). In the salmon lens, increased expression of *hsp70* has been shown after short-term handling stress (30 min), indicating that HSPs are transcriptionally controlled and act to protect the cells after stress-induced protein misfolding (Tröbe et al., 2010). Lower expression of *hsp47* in triploid lenses suggests a poorer ability to facilitate proper folding of proteins. It may be speculated that this is linked to the synthesis, accumulation, repair or breakdown of crystallins or other structural proteins, responsible for lens transparency (Hejtmancik, 2008). Crystallins make up about 80–90% of the soluble proteins in the lens (Hejtmancik, 2008). Mutations in crystalline genes is one of the major reasons for human cataract, and improper ability of chaperones to correct for misfolding or protein damage may render the triploid lens more vulnerable to imbalances responsible for cataract formation in the salmon lens.

When studying the lens transcriptome, it is important to note that the eye lens mostly consists of fiber cells without nuclei and organelles (Bassnett, 2002). With transcription restricted to metabolically active lens epithelial cells and young fiber cells (Hejtmancik et al., 2015), transcriptional differences between diploid and triploid cataractous fish lenses may generally be small. Furthermore, triploid salmon in general differ from diploids by containing fewer and larger cells in most organs (Swarup, 1959; Small and Benfey, 1987), possibly impacting transcriptional differences.

With 74.4% of the total reads mapped to the novel in-house made lens transcriptome, the mapping degree was similar to using a fully sequenced genome as reference. In total however, only 52% of the significant DEGs were annotated using the described pipeline. With manual annotation of all unknowns, about 64% of the DEGs were assigned annotation for IPA functional evaluation. The reason for this relatively poor annotation level is unknown. A good mapping score combined with a poor annotation level might suggest that the lens transcriptome contains a relatively high number of novel transcripts. Among the most strongly differentially regulated genes in both diploid and triploid salmon with cataracts was the CXC chemokine *cxcl1a*. This is a fish-specific chemokine with no mammalian ortholog (Chen et al., 2013), so its function was not included in the IPA functional analysis. This illustrates one of the limitations studying cataract mode of action in

non-model fish species.

In conclusion, this study shows only moderate differences in lens mRNA levels between diploid and triploid Atlantic salmon with score-3 cataract, and very few DEGs with unique expression. Several retina related genes were differentially expressed in the diploid and triploid lenses. The study indicates that the triploid lens may be more vulnerable to cataract than the diploid lens due to predicted effects of protein degradation and turnover, NO-induced oxidative stress, modified cytoskeleton stability and lipid metabolism, possibly linked to repair and compensation mechanisms. Overall, this study suggests that cataract formation is associated with modest changes in gene expression levels, and that transcriptional controls to a large degree regulate gene expression levels independent of chromosomal number in salmon.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

We like to acknowledge the valuable laboratory help from Leikny Fjeldstad (IMR). This project was funded by the Norwegian Research Council (grant number 224816/E40). The fish used in this study were produced in the Norwegian Research Council project INTERACT (grant number 200510).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.exer.2020.108150>.

References

- Bai, J., Zheng, Y., Dong, L., Cai, X., Wang, G., Liu, P., 2015. Inhibition of p38 mitogen-activated protein kinase phosphorylation decreases H(2)O(2)-induced apoptosis in human lens epithelial cells. *Graefes Arch. Clin. Exp. Ophthalmol.* 253, 1933–1940.
- Bassnett, S., 2002. Lens organelle degradation. *Exp. Eye Res.* 74, 1–6.
- Benfey, T.J., Sutterlin, A.M., Thompson, R.J., 1984. Use of erythrocyte measurements to identify triploid salmonids. *Can. J. Fish. Aquat. Sci.* 41, 980–984.
- Benfey, T.J., 1999. The physiology and behavior of triploid fishes. *Rev. Fish. Sci.* 7, 39–67.
- Benfey, T.J., 2016. Effectiveness of triploidy as a management tool for reproductive containment of farmed fish: Atlantic salmon (*Salmo salar*) as a case study. *Rev. Aquacult.* 8 (3), 264–282.
- Bicskei, B., Bron, J.E., Glover, K.A., Taggart, J.B., 2014. A comparison of gene transcription profiles of domesticated and wild Atlantic salmon (*Salmo salar* L.) at early life history stages under controlled conditions. *BMC Genom.* 15, 884.
- Bicskei, B., Taggart, J.B., Glover, K.A., Bron, J.E., 2016. Comparing the transcriptome of embryos from domesticated and wild Atlantic salmon (*Salmo salar*) stocks and examining factors influencing heritability of expression. *Gen. Select. Evol.* 48, 20.
- Bjerkås, E., Waagbø, R., Sveier, H., Bjerkås, I., Bjørnstad, E., Maage, A., 1996. Cataract development in Atlantic Salmon (*Salmo salar* L.) in fresh water. *Acta Vet. Scand.* 37, 351–360.
- Bjerkås, E., Breck, O., Waagbø, R., 2006. The role of nutrition in cataract formation in farmed fish. *CAB Rev* 1, 1–16.
- Bradshaw, A.D., 2009. The role of SPARC in extracellular matrix assembly. *J. Cell Commun. Signal.* 3, 239–246.
- Breck, O., Sveier, H., 2001. Growth and cataract development in two groups of Atlantic salmon (*Salmo salar* L.) post smolts transferred to sea with a four week interval. *Bull. Europ. Assoc. Fish Pathol.* 21, 91–103.
- Breck, O., Bjerkås, E., Campbell, P., Arnesen, P., Haldorsen, P., Waagbø, R., 2003. Cataract preventative role of mammalian blood meal, histidine, iron and zinc in diets for Atlantic salmon (*Salmo salar* L.) of different strains. *Aquacult. Nutr.* 9, 341–350.
- Breck, O., Bjerkås, E., Sanderson, J., Waagbø, R., Campbell, P., 2005a. Dietary histidine affects lens protein turnover and synthesis of N-acetylhistidine in Atlantic salmon (*Salmo salar* L.) undergoing parr-smolt transformation. *Aquacult. Nutr.* 11, 321–332.
- Breck, O., Bjerkås, E., Campbell, P., Rhodes, J.D., Sanderson, J., Waagbø, R., 2005b. Histidine nutrition and genotype affect cataract development in Atlantic salmon, *Salmo salar* L. *J. Fish. Dis.* 28, 357–371.
- Brennan, L.A., Kantorow, M., 2009. Mitochondrial function and redox control in the aging eye: role of MsrA and other repair systems in cataract and macular degenerations. *Exp. Eye Res.* 88, 195–203.
- Chamberlain, C.G., Mansfield, K.J., Cerra, A., 2008. Nitric oxide, a survival factor for lens epithelial cells. *Mol. Vision* 14, 983–991.
- Chatchaiphan, S., Srisapoom, P., Kim, J.H., Devlin, R.H., Na-Nakorn, U., 2017. De Novo transcriptome characterization and growth-related gene expression profiling of diploid and triploid bighead catfish (*Clarias macrocephalus* Gunther, 1864). *Mar. Biotechnol.* 19, 36–48.
- Chen, J., Xu, Q., Wang, T., Collet, B., Corripio-Miyar, Y., Bird, S., Xie, P., Nie, P., Secombes, C.J., Zou, J., 2013. Phylogenetic analysis of vertebrate CXC chemokines reveals novel lineage specific groups in teleost fish. *Dev. Comp. Immunol.* 41, 137–152.
- Chen, Y., Wu, Q., Miao, A., Jiang, Y., Wu, X., Wang, Z., Wu, F., Lu, Y., 2014. Effect of HSP4b on age related cataract may through its novel downstream target Hif1alpha. *Biochem. Biophys. Res. Commun.* 453, 674–678.
- Conesa, A., Gotz, S., Garcia-Gomez, J.M., Terol, J., Talon, M., Robles, M., 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21, 3674–3676.
- Christensen, K.A., Sakhrani, D., Rondeau, E.B., Richards, J., Koop, B.F., Devlin, R.H., 2019. Effect of triploidy on liver gene expression in chou salmon (*Oncorhynchus kisutch*) under different metabolic states. *BMC Genom.* 20, 336.
- Dhakal, S., Stevens, C.B., Sebbagh, M., Weiss, O., Frey, R.A., Adamson, S., Shelden, E.A., Inbal, A., Stenkamp, D.L., 2015. Abnormal retinal development in Cloche mutant zebrafish. *Dev. Dyn.* 244, 1439–1455.
- Enright, J.M., Toomey, M.B., Sato, S.Y., Temple, S.E., Allen, J.R., Fujiwara, R., Kramlinger, V.M., Nagy, L.D., Johnson, K.M., Xiao, Y., How, M.J., Johnson, S.L., Roberts, N.W., Kefalov, V.J., Guengerich, F.P., Corbo, J.C., 2015. Cyp27c1 red-shifts the spectral sensitivity of photoreceptors by converting vitamin A1 into A2. *Curr. Biol.* 25, 3048–3057.
- Gao, Y., Liu, S., Huang, J., Guo, W., Chen, J., Zhang, L., Zhao, B., Peng, J., Wang, A., Wang, Y., Xu, W., Lu, S., Yuan, M., Guo, Q., 2014. The ECM-cell interaction of cartilage extracellular matrix on chondrocytes. *BioMed Res. Int.* 2014, 648459.
- Gilmour, D.T., Lyon, G.J., Carlton, M.B., Sanes, J.R., Cunningham, J.M., Anderson, J.R., Hogan, B.L., Evans, M.J., Colledge, W.H., 1998. Mice deficient for the secreted glycoprotein SPARC/osteonectin/BM40 develop normally but show severe age-onset cataract formation and disruption of the lens. *EMBO J.* 17, 1860–1870.
- GeneCards database. <http://www.genecards.org>.
- Gleim, S., Stojanovic, A., Arehart, E., Byington, D., Hwa, J., 2009. Conserved rhodopsin intradiscal structural motifs mediate stabilization: effects of zinc. *Biochemistry* 48, 1793–1800.
- Glover, K.A., Solberg, M.F., McGinnity, P., Hindar, K., Verspoor, E., Coulson, M.W., Hansen, M.M., Araki, H., Skaala, Ø., Svåsand, T., 2017. Half a century of genetic interaction between farmed and wild Atlantic salmon: status of knowledge and unanswered questions. *Fish Fish.* 18 (5), 890–927.
- Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q.D., Chen, Z.H., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., Regev, A., 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* 29, 644–652.
- Greiling, T.M., Stone, B., Clark, J.I., 2009. Absence of SPARC leads to impaired lens circulation. *Exp. Eye Res.* 89, 416–425.
- Hamel, F.G., Upward, J.L., Siford, G.L., Duckworth, W.C., 2003. Inhibition of proteasome activity by selected amino acids. *Metabolism* 52, 810–814.
- Harvey, A.C., Fjeldal, P.G., Solberg, M.F., Hansen, T., Glover, K.A., 2017. Ploidy elicits a whole-genome dosage effect: growth of triploid Atlantic salmon is linked to the genetic origin of the second maternal chromosome set. *BMC Genet.* 18, 34.
- Hawse, J.R., Hejtmancik, J.F., Huang, Q., Sheets, N.L., Hosack, D.A., Lempicki, R.A., Horwitz, J., Kantorow, M., 2003. Identification and functional clustering of global gene expression differences between human age-related cataract and clear lenses. *Mol. Vision* 9, 515–537.
- Hejtmancik, J.F., 2008. Congenital cataracts and their molecular genetics. *Semin. In Cell Develop. Biol.* 19 (2), 134–149. <https://doi.org/10.1016/j.semdb.2007.10.003>.
- Hejtmancik, J.F., Riazuddin, S.A., McGreal, R., Liu, W., Cvekl, A., Shiels, A., 2015. Lens biology and Biochemistry. *Prog. Mol. Biol. Transl. Sci.* 134, 169–201.
- Iseli, C., Jongeneel, C.V., Bucher, P., 1999. ESTScan: a program for detecting, evaluating, and reconstructing potential coding regions in EST sequences. In: *Proceedings/... International Conference on Intelligent Systems for Molecular Biology; ISMB. International Conference on Intelligent Systems for Molecular Biology*, pp. 138–148.
- Ito, Y., Nabekura, T., Takeda, M., Nakao, M., Terao, M., Hori, R., Tomohiro, M., 2001. Nitric oxide participates in cataract development in selenite-treated rats. *Curr. Eye Res.* 22, 215–220.
- Iyengar, L., Wang, Q., Rasko, J.E., McAvoy, J.W., Lovicu, F.J., 2007. Duration of ERK1/2 phosphorylation induced by FGF or ocular media determines lens cell fate. *Differentiation* 75, 662–668.
- Jennen, D.G., Magkoufopoulou, C., Ketelslegers, H.B., van Herwijnen, M.H., Kleinjans, J. C., van Delft, J.H., 2010. Comparison of HepG2 and HepaRG by whole-genome gene expression analysis for the purpose of chemical hazard identification. *Toxicol. Sci.* 115, 66–79.
- Ji, Y., Rong, X., Ye, H., Zhang, K., Lu, Y., 2015. Proteomic analysis of aqueous humor proteins associated with cataract development. *Clin. Biochem.* 48, 1304–1309.
- Kakrana, A., Yang, A., Anand, D., Djordjevic, D., Ramachandruni, D., Singh, A., Huang, H., Ho, J.W.K., Lachke, S.A., 2018. iSyTE 2.0: a database for expression-based gene discovery in the eye. *Nucleic Acids Res.* 46, D875–D885.
- Ketola, H.G., 1979. Influence of dietary zinc on cataracts in rainbow trout (*Salmo gairdneri*). *J. Nutr.* 109, 965–969.
- Lenox, A.R., Bhootada, Y., Gorbatyuk, O., Fullard, R., Gorbatyuk, M., 2015. Unfolded protein response is activated in aged retinas. *Neurosci. Lett.* 609, 30–35.
- Lou, M.F., 2003. Redox regulation in the lens. *Prog. Retin. Eye Res.* 22, 657–682.

- Mansergh, F.C., Wride, M.A., Walker, V.E., Adams, S., Hunter, S.M., Evans, M.J., 2004. Gene expression changes during cataract progression in Sparc null mice: differential regulation of mouse globins in the lens. *Mol. Vision* 10, 490–511.
- Manthey, A.L., Terrell, A.M., Lachke, S.A., Polson, S.W., Duncan, M.K., 2014. Development of novel filtering criteria to analyze RNA-sequencing data obtained from the murine ocular lens during embryogenesis. *Genom. Data* 2, 369–374.
- Medvedovic, M., Tomlinson, C.R., Call, M.K., Grogg, M., Tsonis, P.A., 2006. Gene expression and discovery during lens regeneration in mouse: regulation of epithelial to mesenchymal transition and lens differentiation. *Mol. Vision* 12, 422–440.
- Nuez-Ortin, W.G., Carter, C.G., Wilson, R., Cooke, I.R., Amoroso, G., Cobcroft, J.M., Nichols, P.D., 2017. Triploid Atlantic salmon shows similar performance, fatty acid composition and proteome response to diploids during early freshwater rearing. *Comp. Biochem. Physiol. Part D Genomics Proteomics* 22, 67–77.
- Ornek, K., Karel, F., Buyukbingol, Z., 2003. May nitric oxide molecule have a role in the pathogenesis of human cataract? *Exp. Eye Res.* 76, 23–27.
- Ottonello, S., Foronia, C., Cartab, A., Petruccob, S., Maraini, G., 2000. Oxidative stress and age-related cataract. *Ophthalmologica* 214, 78–85.
- Piferrer, F., Beaumont, A., Falguiere, J.-C., Flajshans, M., Haffray, P., Colombo, L., 2009. Polyploid fish and shellfish: production, biology and applications to aquaculture for performance improvement and genetic containment. *Aquaculture* 293, 125–156.
- Posner, M., McDonald, M.S., Murray, K.L., Kiss, A.J., 2019. Why does the zebrafish cloche mutant develop lens cataract? *PLoS One* 14 (3), e0211399.
- Remø, S.C., Olsvik, P.A., Torstensen, B.E., Amlund, H., Breck, O., Waagbø, R., 2011. Susceptibility of Atlantic salmon lenses to hydrogen peroxide oxidation *ex vivo* after being fed diets with vegetable oil and methylmercury. *Exp. Eye Res.* 92, 414–424.
- Remø, S.C., Hevrøy, E.M., Olsvik, P.A., Fontanillas, R., Breck, O., Waagbø, R., 2014. Dietary histidine requirement to reduce the risk and severity of cataracts is higher than the requirement for growth in Atlantic salmon smolts, independently of the dietary lipid source. *Br. J. Nutr.* 111, 1759–1772.
- Remø, S.C., Hevrøy, E.M., Breck, O., Olsvik, P.A., Waagbø, R., 2017. Lens metabolomic profiling as a tool to understand cataractogenesis in Atlantic salmon and rainbow trout reared at optimum and high temperature. *PLoS One* 12, e0175491.
- Rhodes, J.D., Breck, O., Waagbø, R., Bjerkås, E., Sanderson, J., 2010. N-acetylhistidine, a novel osmolyte in the lens of Atlantic salmon (*Salmo salar* L.). *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 299, R1075–R1081.
- Rosenbaum, J.T., Samples, J.R., Seymour, B., Langlois, L., David, L., 1987. Chemotactic activity of lens proteins and the pathogenesis of phacolytic glaucoma. *Arch. Ophthalmol.* 105, 1582–1584.
- Sambraus, F., Fjellidal, P.G., Remø, S.C., Hevrøy, E.M., Nilsen, T.O., Thorsen, A., Hansen, T.J., Waagbø, R., 2017. Water temperature and dietary histidine affect cataract formation in Atlantic salmon (*Salmo salar* L.) diploid and triploid yearling smolt. *J. Fish. Dis.* 40, 1195–1212.
- Sanderson, L.M., de Groot, P.J., Hooiveld, G.J., Koppen, A., Kalkhoven, E., Muller, M., Kersten, S., 2008. Effect of synthetic dietary triglycerides: a novel research paradigm for nutrigenomics. *PLoS One* 3, e1681.
- Schneider, E.H., Weaver, J.D., Gaur, S.S., Tripathi, B.K., Jesaitis, A.J., Zelenka, P.S., Gao, J.L., Murphy, P.M., 2012. The leukocyte chemotactic receptor FPR1 is functionally expressed on human lens epithelial cells. *J. Biol. Chem.* 287, 40779–40792.
- Shichida, Y., Yamashita, T., 2003. Diversity of visual pigments from the viewpoint of G protein activation—comparison with other G protein-coupled receptors. *Photochem. Photobiol. Sci.* 2, 1237–1246.
- Shiels, A., Bennett, T.M., Hejtmancik, J.F., 2010. Cat-Map: putting cataract on the map. *Mol. Vision* 16, 2007–2015.
- Shiels, A., Hejtmancik, J.F., 2017. Mutations and mechanisms in congenital and age-related cataracts. *Exp. Eye Res.* 156, 95–102.
- Shiels, A., Hejtmancik, J.F., 2019. Biology of inherited cataracts and opportunities for treatment. *Annu. Rev. Vis. Sci.* 5, 123–149.
- Slingsby, C., Wistow, G.J., Clark, A.R., 2013. Evolution of crystallins for a role in the vertebrate eye lens. *Protein Sci.* 22, 367–380.
- Small, S.A., Benfey, T.J., 1987. Cell-size in triploid salmon. *J. Exp. Zool.* 241, 339–342.
- Sousounis, K., Tsonis, P.A., 2012. Patterns of gene expression in microarrays and expressed sequence tags from normal and cataractous lenses. *Hum. Genom.* 6, 14.
- Stien, L.H., Sæter, P.A., Kristiansen, T., Fjellidal, P.G., Sambraus, F., 2019. First collective report: welfare of triploid salmon in Northern Norway. *Rapport fra Havforskningen* 2019–2047.
- Stojanovic, A., Stitham, J., Hwa, J., 2004. Critical role of transmembrane segment zinc binding in the structure and function of rhodopsin. *J. Biol. Chem.* 279, 35932–35941.
- Swarup, H., 1959. Effect of triploidy on the body size, general organization and cellular structure in *Gasterosteus aculeatus* (L.). *J. Genet.* 56, 143–155.
- Taranger, G.L., Carrillo, M., Schulz, R.W., Fontaine, P., Zanuy, S., Felip, A., Weltzien, F. A., Dufour, S., Karlsen, O., Norberg, B., Andersson, E., Hansen, T., 2010. Control of puberty in farmed fish. *Gen. Comp. Endocrinol.* 165, 483–515.
- Tarazona, S., Garcia-Alcalde, F., Dopazo, J., Ferrer, A., Conesa, A., 2011. Differential expression in RNA-seq: a matter of depth. *Genome Res.* 21, 2213–2223.
- Taube, A.B., Hardenborg, E., Wetterhall, M., Artemenko, K., Hanrieder, J., Andersson, M., Alm, A., Bergquist, J., 2012. Proteins in aqueous humor from cataract patients with and without pseudoexfoliation syndrome. *Eur. J. Mass Spectrom.* 18, 531–541.
- Taylor, A., Davies, K.J.A., 1987. Protein oxidation and loss of protease activity may lead to cataract formation in the aged lens. *Free Rad. Biol. Med.* 3, 371–377.
- Taylor, J.F., Waagbø, R., Diez-Padrisa, M., Campbell, P., Walton, J., Hunter, D., Matthews, C., Migaud, H., 2015. Adult triploid Atlantic salmon (*Salmo salar*) have higher dietary histidine requirements to prevent cataract development in seawater. *Aquacult. Nutr.* 21, 18–32.
- Terrell, A.M., Anand, D., Smith, S.F., Dang, C.A., Waters, S.M., Pathania, M., Beebe, D.C., Lachke, S.A., 2015. Molecular characterization of mouse lens epithelial cell lines and their suitability to study RNA granules and cataract associated genes. *Exp. Eye Res.* 131, 42–55.
- Trøbe, C., Waagbø, R., Breck, O., Stavrum, A.K., Petersen, K., Olsvik, P.A., 2009. Genome-wide transcription analysis of histidine-related cataract in Atlantic salmon (*Salmo salar* L.). *Mol. Vision* 15, 1332–1350.
- Trøbe, C., Waagbø, R., Breck, O., Olsvik, P.A., 2010. Optimisation of gene expression analysis in Atlantic salmon lenses by refining sampling strategy and tissue storage. *Fish Physiol. Biochem.* 36, 1217–1225.
- Urbak, L., Vorum, H., 2010. Heat shock proteins in the human eye. *Int. J. Proteomics* 2010, 479571.
- Utheim, O.A., Ritland, J.S., Utheim, T.P., Espeseth, T., Lydersen, S., Rootwelt, H., Semb, S.O., Elsas, T., 2008. Apolipoprotein E genotype and risk for development of cataract and age-related macular degeneration. *Acta Ophthalmol.* 86, 401–403.
- Waagbø, R., Bjerkås, E., Sveier, H., Breck, O., Bjørnstad, E., Maage, A., 1996. Nutritional status assessed in groups of smolting Atlantic salmon, *Salmo salar* L., developing cataracts. *J. Fish. Dis.* 19, 365–373.
- Waagbø, R., Sveier, H., Breck, O., Bjørnstad, E., Maage, A., Bjerkås, E., 1998. Cataract formation in smolting Atlantic salmon, *Salmo salar*, fed low and high energy diets. *Bull. Eur. Ass. Fish Pathol.* 18, 201.
- Waagbø, R., Trøbe, C., Koppe, W., Fontanillas, R., Breck, O., 2010. Dietary histidine supplementation prevents cataract development in adult Atlantic salmon, *Salmo salar* L., in seawater. *Br. J. Nutr.* 104, 1460–1470.
- Wall, A.E., Richards, R.H., 1992. Occurrence of cataracts in triploid Atlantic salmon (*Salmo salar*) on four farms in Scotland. *Vet. Rec.* 131, 553–557.
- Weleber, R.G., Francis, P.J., Trzucek, K.M., Beattie, C., 1993. Leber congenital amaurosis. In: Pagon, R.A., Adam, M.P., Ardinger, H.H., Wallace, S.E., Amemiya, A., Bean, L.J.H., Bird, T.D., Fong, C.T., Mefford, H.C., Smith, R.J.H., Stephens, K. (Eds.), *GeneReviews*(R), Seattle (WA).
- Wang, K.Y., Spector, A., 1995. Alpha-crystallin can act as a chaperone under conditions of oxidative stress. *Invest. Ophthalm. Vis. Sci.* 36, 311–321.
- Weng, J., Luo, J., Cheng, X., Jin, C., Zhou, X., Qu, J., Tu, L., Ai, D., Li, D., Wang, J., Martin, J.F., Amendt, B.A., Liu, M., 2008. Deletion of G protein-coupled receptor 48 leads to ocular anterior segment dysgenesis (ASD) through down-regulation of Pitx2. *Proc. Natl. Acad. Sci. U. S. A.* 105, 6081–6086.
- Williams, D.L., 2006. Oxidation, antioxidants and cataract formation: a literature review. *Vet. Ophthalmol.* 9, 292–298.
- Wistow, G., Bernstein, S.L., Wyatt, M.K., Behal, A., Touchman, J.W., Bouffard, G., Smith, D., Peterson, K., 2002. Expressed sequence tag analysis of adult human lens for the NEIBank Project: over 2000 non-redundant transcripts, novel genes and splice variants. *Mol. Vision* 8, 171–184.
- Wistow, G., 2006. The NEIBank project for ocular genomics: data-mining gene expression in human and rodent eye tissues. *Prog. Retin. Eye Res.* 25, 43–77.
- Wride, M.A., Mansergh, F.C., Adams, S., Everitt, R., Minnema, S.E., Rancourt, D.E., Evans, M.J., 2003. Expression profiling and gene discovery in the mouse lens. *Mol. Vision* 9, 360–396.
- Wu, C.Y., Rolfe, P.A., Gifford, D.K., Fink, G.R., 2010. Control of transcription by cell size. *PLoS Biol.* 8, e1000523.
- Yan, Q., Clark, J.I., Sage, E.H., 2000. Expression and characterization of SPARC in human lens and in the aqueous and vitreous humors. *Exp. Eye Res.* 71, 81–90.
- Ye, J., Fang, L., Zheng, H., Zhang, Y., Chen, J., Zhang, Z., Wang, J., Li, S., Li, R., Bolund, L., Wang, J., 2006. WEGO: a web tool for plotting GO annotations. *Nucleic Acids Res.* 34, W293–W297.