



This is a post-peer-review, pre-copyedit version of an article published in European Food Research and Technology. The final authenticated version is available online at: <https://doi.org/10.1007/s00217-020-03529-5>

Document downloaded from:



1 **Effectiveness of a pomegranate peel extract (PGE) in reducing *Listeria monocytogenes in vitro***  
2 **and on fresh-cut pear, apple and melon.**

3  
4 **Belgacem I.<sup>1</sup>, Schena L.<sup>1\*</sup>, Teixidó N.<sup>2</sup>, Romeo F.V.<sup>3</sup>, Ballistreri G.<sup>3</sup>, Abadias M.<sup>2</sup>.**

5  
6 <sup>1</sup> Dipartimento di Agraria, Università Mediterranea di Reggio Calabria, Località Feo di Vito, 89122 Reggio Calabria,  
7 Italy.

8 <sup>2</sup> IRTA, XaRTA-Postharvest, Edifici Fruitcentre, Parc Científic i Tecnològic Agroalimentari de Lleida, Parc de Gardeny,  
9 Lleida, Catalonia, Spain.

10 <sup>3</sup> Consiglio per la ricerca in agricoltura e l'analisi dell'economia agraria (CREA) – Centro di Ricerca  
11 Olivicoltura, Frutticoltura e Agrumicoltura, Corso Savoia 190, 95024 Acireale, Italy.

12  
13 **Abstract**

14 Pomegranate peel extract (PGE) is a new promising natural alternative control substance with large spectrum of activity  
15 against wide range of pathogenic microorganisms. In the present study, PGE was firstly investigated as natural  
16 antimicrobial against *Listeria monocytogenes* both *in vitro* and on fresh-cut fruits. The *in vitro* results showed quick and  
17 strong bactericidal and bacteriostatic activity against 5 different strains which were almost completely inhibited by the  
18 extract. Furthermore, it significantly decreased growth rate and maximum growth of all tested strains. *In vivo* trials,  
19 confirmed a strong antibacterial activity of the extract that significantly reduced the bacterial load on fresh-cut apple,  
20 melon and pear and maintained the population at low levels throughout the storage period (7 days). PGE at 12 g/l reduced  
21 *L. monocytogenes* by 1.24, 1.89, and 0.91 log units soon after treatment and by 3.81, 1.53, and 2.99 log units, after 7 days  
22 of storage on apple, pear and melon, respectively. This high antibacterial activity could be mainly related to the high  
23 content of polyphenols (ellagitannins) in the extract. Overall, results of this study suggest a potential industrial application  
24 of PGE to reduce the growth of the pathogenic microorganisms in fresh-cut fruit and ensure a microbial safety in case of  
25 contamination.

26 **Keywords:** PGE, *Listeria monocytogenes*, antimicrobial activity, fresh-cut fruits.

27 **Introduction**

28 In recent years, the demand for healthy and ready-to-eat fresh-cut products has highly increased and, therefore, the  
29 industry is in continuous search for new and improved methods to maintain the quality and extend the shelf-life of  
30 products. Fresh-cut fruits and vegetables are minimally processed products (trimmed, peeled and/or cut) that offer to  
31 consumers high nutritional value, freshness, convenience and flavour similar to the original raw intact product [1,2].  
32 However, these products deteriorate faster than the unprocessed raw material, mainly due to the damages caused by  
33 peeling operation as well as the other minimally processing operations [3,4]. This alters the processed product and makes  
34 it more vulnerable to microbial contamination and colonization with the consequent reduction of quality and shelf life  
35 [3].

---

\* Corresponding author

E-mail addresses: [imen.belgacem@unirc.it](mailto:imen.belgacem@unirc.it) (Belgacem I.), [lschena@unirc.it](mailto:lschena@unirc.it) (Schena L.), [neus.teixido@irta.cat](mailto:neus.teixido@irta.cat) (Teixidó N.), [floravaleria.romeo@crea.gov.it](mailto:floravaleria.romeo@crea.gov.it) (Romeo F.V.), [gabriele.ballistreri@crea.gov.it](mailto:gabriele.ballistreri@crea.gov.it) (Ballistreri G.), [isabel.abadias@irta.cat](mailto:isabel.abadias@irta.cat) (Abadias M.).

36 Microbial contamination may represent a direct critical risk for human health because of the proliferation of important  
37 pathogens such as *Listeria monocytogenes* [5]. This bacteria is an important human pathogen that can contaminate fresh-  
38 cut produces in any step of the processing chain [6]. Therefore, several methods and strategies have been developed and  
39 used by the fresh-cut industries in attempt to reduce the occurrence and the risk associated to foodborne diseases.  
40 Sanitizers, including chlorine [7], organic acids [8], heat treatments [9], ultraviolet (UV) light [10], and ozone [11] have  
41 been widely applied to disinfect and reduce the initial bacterial load on fruits and vegetables. However, these methods  
42 have shown several drawbacks such as the formation of potential carcinogenic by-products from using chlorine, low  
43 efficiency in reducing the bacterial population, chemical residues, destruction of nutrients and the alteration of sensory  
44 characteristics [6,12]. This, together with the increase of the consumer awareness in food safety and healthy living, has  
45 increased the interest to safe and environmentally friendly alternative control means and mainly plant substances including  
46 essential oils and plant extracts.

47 Recently, a pomegranate peel extract (PGE) proved to be very effective in controlling fungal postharvest rots on different  
48 fruit species [13]. Experiments demonstrated a complex mechanism of action which include the induction of resistance  
49 in treated host tissues and a strong antimicrobial activity against both fungi and bacteria [14-16]. The high antimicrobial  
50 activity was associated to the high content of phenolic compounds in PGE [17]. Although PGE has never been tested  
51 against potential human bacterial pathogens, other extracts from pomegranate by-products were able to reduce the  
52 germination and growth of several pathogenic bacteria including *Listeria monocytogenes*, *L. innocua*, *Staphylococcus*  
53 *aureus*, *Escherichia coli*, *Yersinia enterocolitica*, *Pseudomonas aeruginosa*, and *Salmonella* spp. [18,19,14].  
54 Furthermore, edible coatings formulated with a pomegranate peel extract and other anti-browning agents were used to  
55 extend the shelf life of fresh-cut persimmon fruits [20].

56 The aim of the present study was to evaluate the potential use of PGE as natural antimicrobial to reduce the growth of  
57 foodborne pathogens using *L. monocytogenes* as a model pathogen *in vitro* and on fresh-cut melons, apples and pears.

## 58 **Material and Methods**

### 59 **Pomegranate peel extract (PGE) and bacterial strains**

60 All experiments were conducted using a stock solution of an aqueous pomegranate peel extract (PGE) prepared according  
61 to Romeo et al. (2015). The solution was stored, before use, at  $5\pm 1$  °C and diluted to have 3 concentrations of PGE  
62 containing 12 (PGE-12), 2.4 (PGE-2.4), and 1.2 (PGE-1.2) g/l of dry matter. Since the pH of these solutions was very  
63 low (2.7, 2.8, 3.1, respectively), PGE-12 was adjusted with phosphate buffer to increase the pH to 4.4 (aPGE-12) and  
64 included in experiments with fresh-cut fruit plugs in order to evaluate the potential impact of solution acidity on the  
65 antimicrobial activity.

66 Four strains of *L. monocytogenes* belonging to the Spanish Type Culture Collection (CECT 4031, serovar 1/2, CECT 933,  
67 serovar 3a, CECT 940, serovar 4d, CECT 4032, serovar 4b) and a strain that was previously isolated from fresh-cut lettuce  
68 (Lm 230, serovar 1/2 a,[21]) were used in the present study. Strains were grown individually in tryptone soy broth (TSB,  
69 Biokar Diagnostics, Beauvois, France) supplemented with 6 g/l of yeast extract (TSYEB). After 24 h of incubation at  
70  $37\pm 1$  °C, bacterial cells were harvested by centrifugation ( $9800 \times g$  for 10 min at 10 °C) and resuspended in a saline  
71 solution (8.5 g/l NaCl) to obtain single-strain stock suspensions. The concentration of each strain suspension was  
72 determined by plating duplicate 10-fold serial dilutions on TSA media (TSA, Biokar Diagnostics, Beauvois, France)  
73 enriched with 6 g/l of yeast extract, 2.5 g/l glucose and 2.5 g/l  $K_2HPO_4$ , TSAYE) and incubated at 37 °C for 24 h.

#### 74 ***In vitro* assays**

75 To evaluate the bactericidal activity of PGE, 50 µl of *L. monocytogenes* suspensions (10<sup>9</sup> UFC/ml) were added to 5 ml of  
76 PGE at three different concentrations (12, 2.4 and 1.2 g/l). Sterile water was used as control. For each strain and  
77 concentration three replicates were used. After 2, 5, 10 and 30 min of contact time at 20°C, bacterial suspensions of *L.*  
78 *monocytogenes* were 10-fold serially diluted in saline peptone (8.5 g L<sup>-1</sup> NaCl and 1 g L<sup>-1</sup> peptone) and plated on TSA  
79 (TSA, Biokar Diagnostics, Beauvois, France) enriched with 6 g/l of yeast extract, 2.5 g/l glucose and 2.5 g/l K<sub>2</sub>HPO<sub>4</sub>,  
80 TSAYE). After 24 h of incubation at 37 °C, the number of colony forming units was recorded and converted to CFU/ml.  
81 To evaluate the impact of PGE on the growth parameters of *L. monocytogenes*, 20 µl of bacterial suspensions containing  
82 approximately 10<sup>5</sup> CFU/ml were added to 180 µl of TSBYE to obtain final PGE concentrations of 2.4 or 1.2 g/l in a round-  
83 bottomed 96-well microplate (Greiner, Frickenhausen, Germany). TSBYE without PGE served as a control and each  
84 treatment was replicated four times. The microplate was incubated for 36 h at 37±1°C and the absorbance of suspensions  
85 was recorded every 30 min using a spectrophotometer (Epoch Microplate Spectrophotometer, Biotek-Instruments,  
86 Winooski, USA) set at λ = 700 nm. Plates were automatically agitated before measurements.

#### 87 ***In vivo* assays**

88 Experiments were performed on apples (cv. Golden Delicious) and pears (cv. Conference) obtained from local  
89 packinghouses in Lleida (Catalonia, Spain) and on melons (cv. *Cantaloupe*), purchased from a local supermarket. Fruits  
90 were preliminary washed with tap water, surface disinfected with 70% ethanol and dried at room temperature. Fruits were  
91 peeled and cut with a sterilized cork-borer to have cylindrical plugs of 1.2 cm diameter × 1.0 cm long (weighting  
92 approximately 1 g).

93 Fruit plugs were inoculated with *L. monocytogenes* by dipping in a bacterial suspension (10<sup>6</sup> CFU/mL) containing the  
94 five strains of the pathogen, for 2 min. The bacterial suspension was obtained by mixing equal volumes of the single-  
95 strain stock solutions. Inoculated fruit plugs were air dried at room temperature for 30 min and incubated overnight at  
96 5°C. Plugs of each fruit were then divided into 6 uniform groups and subjected to different treatments including PGE-12,  
97 PGE-2.4, PGE-1.2, aPGE-12, and distilled water. Other plugs did not receive any treatment. Treatments were performed  
98 by dipping the inoculated plugs for 10 min at 150 rpm. After drying for 30 min at room temperature, plugs from each  
99 treatment were further divided into two sub-groups, each consisting of 6 replicates. Sub-groups were used to determine  
100 the concentration of bacterial cells soon after the treatment or after 7 days of storage at 10±1°C. To determine the bacterial  
101 population, plugs were put in a sterile bag containing 9 ml of buffered peptone water (BPW, Oxoid, LTD, Basingstoke,  
102 Hampshire, England) and blended in a homogenizer (Minimix® 100, Interscience, France) for 120 s at 12 strokes/s. The  
103 homogenized mixtures were then serially diluted in saline peptone, plated on duplicate plates of selective Palcam agar  
104 (Biokar Diagnostics, Beauvois, France) and incubated at 37±1°C for 48 h. The bacterial concentration was expressed as  
105 log CFU/g. The reductions in bacteria were calculated by subtracting the initial mean bacteria population of the untreated  
106 samples from the bacteria population after each treatment.

#### 107 **Statistical analysis**

108 Prior analyses, all CFU g<sup>-1</sup> data were transformed to log<sub>10</sub> CFU g<sup>-1</sup>. For the bacterial growth experiment, primary models  
109 were fitted using the DMFit 3.5 Excel add-in provided by ComBase predictive modelling tool (<https://www.combase.cc>)  
110 and growth parameters (lag time, growth rate, and maximum population density) were determined using the re-  
111 parameterized Gompertz model described by Zwietering *et al.* (1990) based on the equation  $y =$   
112  $A \exp \left\{ -\exp \left( \frac{\mu_{max} e}{A} (\lambda - t) + 1 \right) \right\}$  where  $y$ ,  $\mu_m$ ,  $t$ ,  $\lambda$ , and  $A$  represent the absorbance (OD) at time  $t$ , maximum growth  
113 rate (h<sup>-1</sup>), incubation time (h), lag time (h), and asymptotic value, respectively.

114 Data were analysed using general linear model analysis with JMP®8, 2004 software (JMP®8, SAS Institute, Cary, NC,  
 115 USA). After analysis of variance (ANOVA), significant differences between treatments were determined according to  
 116 Tukey's test at a significance level of  $P < 0.05$ .

## 117 Results

### 118 *In vitro* assays

119 *In vitro* experiment showed a strong bactericidal activity of PGE. The number of viable cells (log CFU/mL) of *L.*  
 120 *monocytogenes* was always significantly reduced by the extract (Table 1). No significant differences were observed  
 121 among the 3 tested concentrations of PGE. In addition, the incubation time did not have a relevant influence on the  
 122 bactericidal activity as similar results were achieved after 2, 5, 10 and 30 min of contact. On the contrary, important  
 123 differences were observed among *L. monocytogenes* strains. Strains CECT 4031 and CECT 933 were the most sensitive  
 124 since their population was always below the detection limit for almost all tested concentrations and incubation times  
 125 (Table 1). A slightly higher tolerance was revealed for the strain CECT 4032. Strains CECT 940 and Lm230 showed the  
 126 highest rates of survival, but still their population was reduced at least by 3.3 log units after 2 minutes of incubation with  
 127 all the PGE doses.

128 **Table 1** Concentration of *L. monocytogenes* cells (log<sub>10</sub> CFU/ml) after 2, 5, 10 or 30 min of incubation in PGE solutions  
 129 at three different concentrations (1.2, 2.4 or 12.0 g/l) or in water (control). Separate statistical analyses were conducted  
 130 for each strain, contact period, and incubation time. Different letters indicate significantly different values according to  
 131 Tukey's test ( $P < 0.05$ ).

Strain	Treatment	Incubation period (min)							
		2		5		10		30	
CECT 933	Water	7.11	a	7.04	a	7.08	a	7.11	a
	PGE 1.2 g/l	<dl	b	<dl	b	<dl	b	<dl	b
	PGE 2.4 g/l	<dl	b	<dl	b	<dl	b	<dl	b
	PGE 12 g/l	<dl	b	<dl	b	<dl	b	<dl	b
CECT 940	Water	7.26	a	7.08	a	6.97	a	6.63	a
	PGE 1.2 g/l	3.59	b	3.49	b	3.32	b	3.23	b
	PGE 2.4 g/l	3.67	b	3.48	b	3.11	b	3.00	b
	PGE 12 g/l	3.61	b	3.54	b	3.43	b	3.30	b
Lm230	Water	6.93	a	6.90	a	6.93	a	6.90	a
	PGE 1.2 g/l	3.52	b	3.49	b	3.38	b	3.34	b
	PGE 2.4 g/l	3.48	b	3.41	b	3.30	b	3.28	b
	PGE 12 g/l	3.61	b	3.45	b	3.40	b	3.23	b
CECT 4032	Water	7.28	a	7.18	a	7.08	a	7.28	a
	PGE 1.2 g/l	2.28	b	1.89	b	1.37	b	1.30	b
	PGE 2.4 g/l	2.56	b	0.92	b	1.74	b	0.23	b
	PGE 12 g/l	1.56	b	<dl	b	<dl	b	<dl	b
CECT 4031	Water	6.70	a	6.85	a	6.74	a	6.65	a
	PGE 1.2 g/l	0.20	b	0.52	b	<dl	b	<dl	b
	PGE 2.4 g/l	<dl	b	<dl	b	<dl	b	<dl	b
	PGE 12 g/l	<dl	b	<dl	b	<dl	b	<dl	b

132 < dl: below detection level

133 The analysis of the growth parameters of *L. monocytogenes* in TSBYE showed a significant impact of PGE on the  
 134 maximum cell growth of all investigated strains (Table 2). Interestingly, the effect of PGE was directly correlated to its  
 135 concentration since significant differences were always revealed between the two tested concentrations. In particular,

136 PGE-2.4 reduced the maximum cell growth between 46.9% (CECT 9333) and 62.9% (CECT 4031) as compared to the  
 137 control (TSBYE without PGE). While, PGE-1.2, reductions ranged between 18.4% (strain CECT 4032) and 35.3%  
 138 (CECT 933).

139 PGE-2.4 significantly decreased also the growth rate of all strains with reductions ranging between 41.6% (CECT 4032)  
 140 and 63.9% (CECT 933) as compared to the control. Lower, but still significant reductions were also achieved with PGE  
 141 at 1.2 g/l for 4 out of 5 strains. Similarly, the duration of the lag phase was increased by PGE.

142 **Table 2** Growth kinetic parameters (lag time, growth rate, and max absorbance) of the five tested strains of *L.*  
 143 *monocytogenes* cultured in standard TSBYE (control) or in TSBYE amended with PGE at 1.2 and 2.4 g/l. For each  
 144 parameter and strain, different letters indicate statistically different values according to Tukey's test ( $P < 0.05$ ).


Strains	Medium	Lag time, $\lambda$ (h)		Growth rate, $\mu$ (Absorbance at $\lambda = 700$ nm)		Max absorbance	
Lm230	TSBYE	7.5	c	0.002229	a	0.564225	a
	TSBYE+PGE 1.2 g/l	7.9	b	0.001435	b	0.370295	b
	TSBYE+PGE 2.4 g/l	8.4	a	0.001005	c	0.242917	c
CECT 933	TSBYE	9.3	c	0.001122	a	0.491362	a
	TSBYE+PGE 1.2 g/l	11.3	b	0.000610	b	0.317862	b
	TSBYE+PGE 2.4 g/l	13.0	a	0.000405	c	0.261084	c
CECT 940	TSBYE	7.9	c	0.002180	a	0.492187	a
	TSBYE+PGE 1.2 g/l	8.3	b	0.001366	b	0.359077	b
	TSBYE+PGE 2.4 g/l	8.9	a	0.000972	c	0.234525	c
CECT 4031	TSBYE	9.0	b	0.001738	a	0.402777	a
	TSBYE+PGE 1.2 g/l	9.3	ba	0.001000	b	0.313139	b
	TSBYE+PGE 2.4 g/l	9.9	a	0.000685	c	0.149649	c
CECT 4032	TSBYE	6.7	c	0.002335	a	0.495242	a
	TSBYE+PGE 1.2 g/l	7.4	b	0.002658	a	0.403998	b
	TSBYE+PGE 2.4 g/l	8.2	a	0.001364	b	0.216037	c

#### 145 ***In vivo* trials**

##### 146 **Effect of PGE on *L. monocytogenes* population on fresh-cut apple**

147 Initial population of *L. monocytogenes* soon after the inoculation was 5.65 log CFU/g (Fig. 1). Without PGE treatment,  
 148 the population of *L. monocytogenes* greatly increased during the cold storage and reached 7.18 and 7.43 Log CFU/g after  
 149 7 days of cold storage on untreated and water treated fresh-cut apple samples, respectively.

150 PGE proved to be very effective in reducing the population of *L. monocytogenes*. Soon after treatments, significant  
 151 reductions of 1.57, 1.24, 1.21, and 0.85 log units were recorded, comparing to the untreated samples, with aPGE-12, PGE-  
 152 12, PGE-2.4, and PGE-1.2, respectively. These reductions greatly increased after 7 days of cold storage to reach 3.81,  
 153 3.31, and 2.19 log units in apple plugs treated with PGE-12, PGE-2.4, and aPGE-12, respectively. However, PGE with  
 154 the lowest concentration (PGE-1.2) did not show any significant effect in reducing the bacterium population comparing  
 155 to the controls.




156

157 **Fig. 1** Population of *L. monocytogenes* (log CFU/g) determined on fresh-cut apple plugs soon after the treatment (blue  
 158 column) and after 7 days of storage at 10°C (orange column). Inoculated plugs were dipped in different PGE solutions or  
 159 in distilled water (control). Furthermore, untreated plugs were also used as control. Bars indicate standard errors of the  
 160 means. For each assessment time, different columns with different letters indicate significant differences between  
 161 treatments according to Tukey's test ( $P < 0.05$ ).

162 **Effect PGE treatments on *L. monocytogenes* population on fresh-cut pear**

163 Soon after the inoculation, the population of *L. monocytogenes* on untreated pear plugs and on plugs dipped in water was  
 164 estimated at approximately 6.40 log CFU/g (Fig. 2). On these samples, the bacterium greatly proliferated reaching  
 165 approximately 8.50 log CFU/g, after seven days of cold storage.

166 Except of PGE-1.2, all other PGE treatments significantly reduced the populations of *L. monocytogenes*, soon after the  
 167 treatment (Fig. 2). In particular, PGE-12, PGE-2.4, and a-PGE12 reduced the bacterium by 1.89, 1.08, and 1.22 log units,  
 168 respectively. However, after 7 days of cold storage, only PGE-12 significantly reduced the growth of the bacterium with  
 169 a reduction of 1.53 log units compared to untreated fresh-cut pear.



170

171 **Fig. 2** Population of *L. monocytogenes* (log CFU/g) on fresh-cut pear plugs soon after treatments (blue column) and after  
 172 7 days of storage at 10°C (orange column). Inoculated plugs were dipped in different PGE solutions or in distilled water  
 173 (control). Furthermore, untreated plugs were also used as control. Bars indicate standard errors of the means. For each


174 assessment time, different columns with different letters indicate significant differences between treatments according to  
175 Tukey's test ( $P < 0.05$ ).

### 176 Effect of PGE on *L. monocytogenes* population on fresh-cut melon

177 Soon after the inoculation, the population of *L. monocytogenes* on untreated and water treated melon plugs was 6.9 and  
178 6.74 log CFU/ml, respectively. On both samples, the bacterium population increased during the storage at 10 °C, reaching  
179 approximately 9.0 log CFU/g (Fig. 3).

180 Soon after treatments, a significant reduction of *Listeria* population was achieved with all PGE treatments except of PGE-  
181 1.2. In particular, compared to untreated fresh-cut melons, PGE-12, PGE-2.4 and aPGE-12, reduced the bacterial  
182 population by 0.91, 0.58, and 0.85 log units, respectively.

183 After 7 days of storage, the bacterium population on melon plugs was reduced by 2.99, 1.93, and 1.88 log units with PGE-  
184 12, PGE-2.4 and aPGE-12, respectively. However, PGE at the lowest concentration (1.2 g/l) did not show any significant  
185 effect.



186

187 **Fig. 3** Population of *L. monocytogenes* (log CFU/g) on fresh-cut melon plugs soon after treatments (blue column) and  
188 after 7 days of storage at 10°C (orange column). Inoculated plugs were dipped in different PGE solutions or in distilled  
189 water (control). Furthermore, untreated plugs were also used as control. Bars indicate standard errors of the means. For  
190 each assessment time, different columns with different letters indicate significant differences between treatments  
191 according to Tukey's test ( $P < 0.05$ ).

### 192 Discussion

193 The present study represents the first investigation of PGE as natural antimicrobial to reduce and control the growth of  
194 foodborne pathogens on ready-to-eat fresh-cut fruits. Experiments were conducted using *L. monocytogenes* as a model  
195 species in light of its primarily importance as food contaminant, and future investigations will be needed to evaluate the  
196 efficacy of PGE against other foodborne microorganisms. Overall, *in vitro* and *in vivo* results showed high bactericidal  
197 and bacteriostatic effects of PGE against *L. monocytogenes*. In particular, the *in vitro* results (Table 1) revealed that  
198 regardless of the tested concentration, PGE exerted a quick and high significant inhibitory activity against all the  
199 *L. monocytogenes* tested strains by reducing their population by at least 3.3 log units after a short contact time (2 minutes).  
200 This high antibacterial activity could be explained by the composition of the extract. In fact, PGE is rich in polyphenols  
201 (ellagitannins) mainly represented by punicalagins, ellagic acid and its derivatives that have been reported to exert a



202 strong antimicrobial activity against Gram-negative and Gram-positive bacteria [22]. In any case, the absence of the outer  
203 membrane in *L. monocytogenes*, as a Gram-positive bacterium, makes it easier for the extract to alter and, therefore,  
204 causing a loss of the bacterium cellular components [23].

205 The antimicrobial activity of other extracts from pomegranate by-products against a variety of food-borne pathogens  
206 including *L. monocytogenes*, *E. coli*, and *S. aureus* has been already reported [22,19,23]. However, PGE, seems to be  
207 more effective due to its higher content in polyphenols and, being obtained with food grade solvent, it can be considered  
208 a safe and eco-friendly antimicrobial preparation [18,17]. Furthermore, PGE bactericidal activity seems to be stronger  
209 comparing to other plant extracts such as cherry pomace and plum extracts [24,25].

210 PGE also revealed strong bacteriostatic effect and its activity was significantly influenced by the concentration of the  
211 extract (Table 2). In particular, the log phases of the tested bacteria strains grown in broth media containing PGE were  
212 significantly longer. This delayed response of the growth indicates that PGE can negatively modify the growth  
213 environment making it longer for the bacteria to adjust [26]. More importantly, PGE showed high efficiency in reducing  
214 the growth rate as well as the maximum growth of *L. monocytogenes*. This effect may be attributed to the richness of the  
215 extract in tannins that may combine with proteins and cause their precipitation [22,27]. Likely tannins of the extract may  
216 combine with proteins of the bacterial membrane as well as with protein of the culture media forming complexes that lead  
217 to the lysis and death of the bacteria. Moreover, the high concentration of polyphenols of PGE causes the decrease of pH  
218 gradient around the cell membrane and the increase of its permeability, leading to cell death (Singh et al., 2018).

219 *In vivo* results confirmed a strong antibacterial activity of PGE that significantly reduced the bacterial load on fresh-cut  
220 apples, pears and melons and was able to maintain the population at low levels throughout the storage period (7 days).  
221 However, the reduction of the bacterial population in the *in vivo* experiments was overall lower as compared to *in vitro*  
222 conditions. This could be mainly explained by the presence of organic matter as well as to the presence of a solid matrix  
223 that increase the bacterial survival and decrease the contact between the treatment and the bacteria [28,29]. For the same  
224 reason, a higher concentration of PGE seems to be needed to control the bacterium in practical *in vivo* conditions as  
225 confirmed by the low efficacy of the lowest tested concentration of PGE (1.2 g/L). Interestingly, aPGE-12 (pH4.4) showed  
226 a similar efficacy as compared to normal PGE-12 (pH 2.7), both soon after treatments and after 7 days of storage at 10 °C.  
227 This result confirms that the composition of PGE, rather than its low pH, was the main determinant factor for its activity.  
228 In this context, the higher concentration of polyphenols compared to other plant extracts, make pomegranate peel extracts  
229 particularly promising for future applications especially that it already proved major beneficial effects on human health  
230 [30,31]. However, the slight lower efficacy of aPGE-12 may suggest a secondary role of pH in modulating the level of  
231 efficacy, due to the influence of pH on chemical structure and functions of polyphenols. This aspect needs to be taken  
232 into account in future applications and/or in the development of commercial formulations. On a practical point of view,  
233 PGE seem to provide higher levels of reductions of *L. monocytogenes* populations compared to other widely used  
234 alternative compounds. For instance, on fresh cut apples, PGE proved higher efficacy comparing to vanillin, citrox,  
235 hydrogen peroxide and peroxyacetic acid that are generally recognized by the scientific community as effective sanitizers  
236 of fresh-cut fruits [32]. In particular, after storage, the microbial reductions observed with PGE treated apple plugs were  
237 almost double the reduction obtained after treatment with hydrogen peroxide [32].

238 Although the impact of PGE on the sensory quality of the fruits still needs to be evaluated, available data suggest a high  
239 potential of the extract as a natural antimicrobial against foodborne pathogens. Furthermore, the antimicrobial activity of  
240 PGE and/or its spectrum of activity could be further enhanced by combining it with other alternative control means.

241 **Acknowledgements:** The authors wish to thank Marina Anguera and Cristina Solsona for their technical support.

242 **Compliance with ethical standards**

243 **Conflict of Interest:** The authors declare that they have no conflict of interest.

244 **Compliance with ethics requirements:** This article does not contain any studies with human or animal subjects.

## 245 **References**

- 246 1. Gómez-López VM, Rajkovic A, Ragaert P, Smigic N, Devlieghere F (2009) Chlorine dioxide for minimally processed  
247 produce preservation: a review. *Trends Food Sci Technol* 20 (1):17-26
- 248 2. Del Nobile M, Conte A, Scrocco C, Brescia I (2009) New strategies for minimally processed cactus pear packaging.  
249 *Innov Food Sci Emerg Technol* 10 (3):356-362
- 250 3. Prakash A, Baskaran R, Paramasivam N, Vadivel V (2018) Essential oil based nanoemulsions to improve the microbial  
251 quality of minimally processed fruits and vegetables: A review. *Food Res Int* 111:509-523
- 252 4. Rolfe RS, CHISM III GW (1987) Physiological consequences of minimally processed fruits and vegetables. *J Food*  
253 *Qual* 10 (3):157-177
- 254 5. Leverentz B, Conway WS, Janisiewicz W, Abadias M, Kurtzman CP, Camp MJ (2006) Biocontrol of the food-borne  
255 pathogens *Listeria monocytogenes* and *Salmonella enterica* serovar Poona on fresh-cut apples with naturally occurring  
256 bacterial and yeast antagonists. *Appl Environ Microbiol* 72 (2):1135-1140
- 257 6. Chaves RD, Martinez RCR, Rezende ACB, Rocha MD, Oteiza JM, de Souza Sant'Ana A (2016) Salmonella and  
258 *Listeria monocytogenes* in ready-to-eat leafy vegetables. In: *Food Hygiene and Toxicology in Ready-to-Eat Foods*.  
259 Elsevier, pp 123-149
- 260 7. Wu VC, Kim B (2007) Effect of a simple chlorine dioxide method for controlling five foodborne pathogens, yeasts and  
261 molds on blueberries. *Food Microbiol* 24 (7-8):794-800
- 262 8. Mani-Lopez E, García HS, López-Malo A (2012) Organic acids as antimicrobials to control Salmonella in meat and  
263 poultry products. *Food Res International* 45 (2):713-721
- 264 9. Bermúdez-Aguirre D, Corradini MG (2012) Inactivation kinetics of Salmonella spp. under thermal and emerging  
265 treatments: a review. *Food Research Int* 45 (2):700-712
- 266 10. Yaun BR, Sumner SS, Eifert JD, Marcy JE (2004) Inhibition of pathogens on fresh produce by ultraviolet energy. *Int*  
267 *J Food Microbiol* 90 (1):1-8
- 268 11. Wysok B, Uradziński J, Gomólka-Pawlicka M (2006) Ozone as an alternative disinfectant-a review. *Polish J Food*  
269 *Nutr Sci* 15 (1):3
- 270 12. Gil MI, Selma MV, López-Gálvez F, Allende A (2009) Fresh-cut product sanitation and wash water disinfection:  
271 problems and solutions. *Int J Food Microbiol* 134 (1-2):37-45
- 272 13. Li Destri Nicosia MG, Pangallo S, Raphael G, Romeo FV, Strano MC, Rapisarda P, Droby S, Schena L (2016) Control  
273 of postharvest fungal rots on citrus fruit and sweet cherries using a pomegranate peel extract. *Postharvest Biol Technol*  
274 114:54-61
- 275 14. Pangallo S, Li Destri Nicosia MG, Agosteo GE, Abdelfattah A, Romeo FV, Cacciola SO, Rapisarda P, Schena L  
276 (2017) Evaluation of a pomegranate peel extract as an alternative means to control olive anthracnose. *Phytopathology*  
277 107 (12):1462-1467
- 278 15. Pangallo S, Li Destri Nicosia M, Raphael G, Levin E, Ballistreri G, Cacciola S, Rapisarda P, Droby S, Schena L  
279 (2017) Elicitation of resistance responses in grapefruit and lemon fruits treated with a pomegranate peel extract. *Plant*  
280 *Pathol* 66 (4):633-640

- 281 16. Belgacem I, Pangallo S, Abdelfattah A, Romeo FV, Cacciola SO, Li Destri Nicosia MG, Ballistreri G, Schena L  
282 (2019) Transcriptomic Analysis of Orange Fruit Treated with Pomegranate Peel Extract (PGE). *Plants* 8 (4):101
- 283 17. Romeo FV, Ballistreri G, Fabroni S, Pangallo S, Li Destri Nicosia MG, Schena L, Rapisarda P (2015) Chemical  
284 characterization of different sumac and pomegranate extracts effective against *Botrytis cinerea* rots. *Molecules* 20  
285 (7):11941-11958
- 286 18. Al-Zoreky N (2009) Antimicrobial activity of pomegranate (*Punica granatum* L.) fruit peels. *Int J Food Microbiol*  
287 134 (3):244-248
- 288 19. Gullon B, Pintado ME, Pérez-Álvarez JA, Viuda-Martos M (2016) Assessment of polyphenolic profile and  
289 antibacterial activity of pomegranate peel (*Punica granatum*) flour obtained from co-product of juice extraction. *Food*  
290 *Control* 59:94-98
- 291 20. Taberner V, Sanchís E, Mateos M, Palou L, Pérez-Gago M Pectin-based edible coatings formulated with pomegranate  
292 peel extracts and other antibrowning agents to extend shelf life of fresh-cut 'Rojo Brillante' persimmon. In: VIII  
293 International Postharvest Symposium: Enhancing Supply Chain and Consumer Benefits-Ethical and Technological  
294 Issues 1194, 2016. pp 887-894
- 295 21. Abadias M, Usall J, Anguera M, Solsona C, Viñas I (2008) Microbiological quality of fresh, minimally-processed  
296 fruit and vegetables, and sprouts from retail establishments. *Int. J. Food Microbiol* 123 (1-2):121-129
- 297 22. Wu J, Goodrich KM, Eifert JD, Jahncke ML, O'Keefe SF, Welbaum GE, Neilson AP (2018) Inhibiting foodborne  
298 pathogens *Vibrio parahaemolyticus* and *Listeria monocytogenes* using extracts from traditional medicine: Chinese  
299 gallnut, pomegranate peel, Baikal skullcap root and forsythia fruit. *Open Agric* 3 (1):163-170
- 300 23. Li G, Xu Y, Wang X, Zhang B, Shi C, Zhang W, Xia X (2014) Tannin-rich fraction from pomegranate rind damages  
301 membrane of *Listeria monocytogenes*. *Foodborne Pathog Dis* 11 (4):313-319
- 302 24. Kołodziejczyk K, Sójka M, Abadias M, Viñas I, Guyot S, Baron A (2013) Polyphenol composition, antioxidant  
303 capacity, and antimicrobial activity of the extracts obtained from industrial sour cherry pomace. *Ind Crops Prod* 51:279-  
304 288
- 305 25. Sójka M, Kołodziejczyk K, Milala J, Abadias M, Viñas I, Guyot S, Baron A (2015) Composition and properties of  
306 the polyphenolic extracts obtained from industrial plum pomaces. *J Funct Foods* 12:168-178
- 307 26. Swinnen I, Bernaerts K, Dens EJ, Geeraerd AH, Van Impe J (2004) Predictive modelling of the microbial lag phase:  
308 a review. *Int J Food Microbiol* 94 (2):137-159
- 309 27. Singh B, Singh JP, Kaur A, Singh N (2019) Antimicrobial potential of pomegranate peel: a review. *Int J Food Sci*  
310 *Technol* 54 (4):959-965
- 311 28. Rodgers SL, Cash JN, Siddiq M, Ryser ET (2004) A comparison of different chemical sanitizers for inactivating  
312 *Escherichia coli* O157: H7 and *Listeria monocytogenes* in solution and on apples, lettuce, strawberries, and cantaloupe.  
313 *J Food Prot* 67 (4):721-731
- 314 29. KIM JG, Yousef AE, Chism GW (1999) Use of ozone to inactivate microorganisms on lettuce. *J Food Saf* 19 (1):17-  
315 34
- 316 30. Sorrenti V, Randazzo CL, Caggia C, Ballistreri G, Romeo FV, Fabroni S, Timpanaro N, Raffaele M, Vanella L (2019)  
317 Beneficial effects of pomegranate peel extract and probiotics on pre-adipocyte differentiation. *Front Microbiol* 10:660
- 318 31. Howell AB, D'Souza DH (2013) The pomegranate: effects on bacteria and viruses that influence human health. *Evid*  
319 *Based Complement Alternat Med* 2013:606212
- 320 32. Abadias M, Alegre I, Usall J, Torres R, Viñas I (2011) Evaluation of alternative sanitizers to chlorine disinfection for  
321 reducing foodborne pathogens in fresh-cut apple. *Postharvest Biol Technol* 59 (3):289-297