



This is a post-peer-review, pre-copyedit version of an article published in **Molecular Breeding**. The final authenticated version is available online at:
<https://doi.org/10.1007/s11032-019-1040-1>

Document downloaded from:



Multi-environment QTL analysis using an updated genetic map of a widely distributed Seri × Babax spring wheat population

Caiyun Liu¹, Mehdi Khodae², Marta S. Lopes^{3,4}, Carolina Sansaloni⁵, Susanne Dreisigacker¹, Sivakumar Sukumaran^{1,*}, Matthew Reynolds¹

Affiliations:¹Global Wheat Program, International Maize and Wheat Improvement Center, Km 45 Carretera México Veracruz, El Batán, Texcoco CP, México 56237; ²Department of Agronomy and Plant Breeding, College of Agriculture, Isfahan University of Technology (IUT), 84156-8311 Isfahan, Iran; ³Global Wheat Program, International Maize and Wheat Improvement Center (CIMMYT), Emex, 06511 Ankara, Turkey; ⁴Sustainable Field Crops Program, Institute for Food and Agricultural Research and Technology (IRTA), Lleida, Spain; ⁵Genetic resources program, International Maize and Wheat Improvement Center, Km 45 Carretera México Veracruz, El Batán, Texcoco CP, México 56237.

*Corresponding author: S.Sukumaran@cgiar.org

Tel: +521 5541 941 246

ORCIDS:

Caiyun Liu: 0000-0001-7987-703X

Mehdi Khodae:

Marta S. Lopes: 0000-0002-7698-6019

Carolina Sansaloni: 0000-0003-2675-4524

Susanne Dreisigacker: 0000-0002-3546-5989

Sivakumar Sukumaran: 0000-0003-4088-4624

Matthew Reynolds: 0000-0002-4291-4316

Keywords: 90K Wheat SNPs, DArTseq, QTL mapping, Seri × Babax, grain yield

Abbreviations:

QTL Quantitative trait locus

RIL Recombinant inbred line

SNP Single nucleotide polymorphism

PVE Phenotypic variation explained by QTL

1 **Abstract**

2 Seri/Babax spring wheat linkage mapping population was developed to minimize the confounding effect of phenology
3 in the genetic dissection of abiotic stress traits. An existing linkage map (< 500 markers) was updated with 6,470
4 polymorphic Illumina iSelect 90K array and DArTseq SNPs to a genetic map of 5576.5 cM with 1748 non-redundant
5 markers (1165 90K SNPs, 207 DArTseq SNPs, 183 AFLP, 111 DArT array, and 82 SSR) assigned to 31 linkage
6 groups. We conducted QTL mapping for yield and related traits phenotyped in seven major wheat growing areas from
7 Egypt, Sudan, Iran and India, and nine environments (heat, drought, heat plus drought, and yield potential) in Obregon,
8 Mexico. The current study confirmed QTLs from previous studies and identified novel QTLs. QTL analysis identified
9 39 (LOD: 2.5-23.6; PVE: 4.8-21.3%), 36 (LOD: 2.5-15.4; PVE: 2.9-21.4%), 30 (LOD: 2.5-13.1; PVE: 3.6-26.8%),
10 39 (LOD: 2.7-14.4; PVE: 2.6-15.9%), and 22 (LOD: 2.8-4.8; PVE: 6.8-12.9%) QTLs for grain yield, thousand-grain
11 weight, grain number, days to heading, and plant height, respectively. QTL analysis based on high-yielding and low-
12 yielding environment clusters identified 11 additional QTLs (LOD: 2.6-14.9; PVE: 2.7-19.7%). The updated map
13 thereby provides a better genome coverage (3.5-fold) especially in D genome (4-fold), higher density (1.1-fold) and a
14 good collinearity with the IWGSC RefSeq v1.0 genome, thus increased the number of detected QTLs (5-fold)
15 compared with the earlier map. This map provides a useful genomic resource for further genetic analyses of important
16 traits in this wheat population that was widely distributed around the world.

17 INTRODUCTION

18 Bread wheat (*Triticum aestivum* L.) is one of the staple food crops worldwide, contributing ~ 28% of the global grain
19 production and 20% of the calories and protein consumed by human (FAOstat 2018). With a predicted world
20 population of nine billion by 2050, the demand for wheat is expected to increase (Rosegrant and Cline 2003). As the
21 most widely cultivated crop geographically, wheat covers a large variance of climate and soil conditions. Abiotic
22 stresses such as drought, temperature, salinity, and nutrient imbalances reduce wheat yield in many environments
23 (Trethowan and Mujeeb-Kazi 2008). Improvement of grain yield in these target environments is the primary target of
24 wheat research considering climate change (Reynolds et al., 2009).

25 Grain yield (GY) is a complex quantitative trait that is strongly influenced by interacting genetic and
26 environmental factors (Quarrie et al. 2006). GY can be dissected into direct components: thousand-grain weight
27 (TGW) and grain number (GN) per unit area of land. Both TGW and especially GN are quantitatively inherited and
28 not easy to improve simultaneously through the conventional approaches due to the tradeoff between them (Yano and
29 Sasaki 1997; Griffiths et al. 2015). Quantitative trait loci (QTL) mapping in bi-parental populations has provided an
30 effective approach to dissect quantitative traits into component loci to study their relative effects on a specific trait of
31 interest (Doerge 2002). A large number of QTLs for yield and yield components have been identified to date on almost
32 all wheat chromosomes (Bennett et al. 2012; Golabadi et al. 2011; Kirigwi et al. 2007; Liu et al. 2019b; Maccaferri et
33 al. 2008; Pinto et al. 2010; Tahmasebi et al. 2016). However, the identification of QTL for yield may be confounded
34 by flowering time and plant height (PH), whereby genes of major effects have been shown to mask the identification
35 of minor effects, especially under abiotic stress where the onset of stress treatments will occur at different growth
36 stages depending on the genotypes phenological pattern (Reynolds et al. 2009). Therefore, effects of phenology and
37 PH may be restricted in the population to identify QTLs for major traits without the confounding effect of phenology
38 (Lopes et al. 2013; Pinto et al. 2010; Reynolds and Tuberosa 2008).

39 The narrow range of PH and flowering time of the Seri/Babax recombinant inbred line (RIL) population
40 makes it ideal for genetic and physiological studies. Many QTLs have been identified for agronomic and physiological
41 traits in the Seri/Babax population under water limiting and heat stress conditions over the past decade (Lopes et al.
42 2013; Mathews et al. 2008; McIntyre et al. 2010; Pinto et al. 2016; Pinto et al. 2010; Tahmasebi et al. 2016).
43 Seri/Babax population is still widely used by researchers around the world for genetic and physiological studies of
44 different traits in various environments and some of the lines have been valuable in breeding. Genetic linkage map is

45 the foundation for mapping QTLs and subsequent marker-assisted selection or map-based cloning. The existing
46 genetic map of Seri/Babax was primarily based on 475 markers comprising SSR (Simple Sequence Repeat), AFLP
47 (Amplified Fragment Length Polymorphism), and DArT (Diversity Arrays Technology) markers (Lopes et al. 2013).
48 With the rapid development of new genotyping technologies, numerous molecular makers for wheat are available
49 now. Therefore, a new linkage map is necessary for researchers to compare their results with peers. By genotyping
50 the Seri/Babax population with the wheat 90K Illumina iSelect array and DArTseq platform, the objectives of the
51 study was 1) to update the genetic linkage map of the Seri/Babax population by integrating 90K and DArTseq markers
52 to the existing linkage map; 2) to identify QTLs for yield and yield-related traits with phenotypic data from multiple
53 environments, combining studies not used earlier and from earlier published data; and 3) to compare the properties of
54 the updated genetic map with the existing linkage map.

55

56 MATERIALS AND METHODS

57 Wheat population and phenotyping

58 The Seri/Babax population comprising 156 RILs was derived from a reciprocal cross between two elite spring bread
59 wheat, Seri M82 (a released line from the “Veery” cross, KVZ/BUHO// KAL/BB) and Babax (a line derived from the
60 cross “Babax”, BOW/NAC//VEE/3/BJY/COC) (Olivares-Villegas et al. 2007). Seri M82 carries the 1BL.1RS
61 translocation and was characterized by moderate tolerance to drought conditions and high yield potential. Babax is
62 without the 1BL.1RS translocation, and is highly tolerant to severe drought (Mathews et al. 2008). Seri M82 and
63 Babax both have the same photoperiod-insensitive allele at *Ppd-D1*, and spring-type alleles for vernalization (*Vrn-B1*
64 and *Vrn-D1*) loci (Pinto et al. 2010). Hence, this population is characterized by its narrow range of plant height and
65 flowering time (ca. 17 cm, 10-15 days across environments as reported by Lopes et al. (2013)), and is ideal for genetic
66 mapping and physiological studies. The phenotyping was conducted in eight major wheat growing regions from five
67 countries around the world (**Fig. 1a**), including ten environments (Darab in Iran, Sohag in Egypt, Dongola and Wad
68 Medani in Sudan, Ludhiana and Karnal in India, and yield potential, drought stress, heat stress, and heat plus drought
69 stress in Obregon, Mexico in 2008-2009) reported in an earlier study (Lopes et al. 2013), and six new environments
70 (one from Shandaweel, Egypt in 2008-2009, and five from different stresses in Obregon, Mexico in 2004-2005 and
71 2005-2006). The sowing and harvest date, temperature, precipitation, total amount of water applied by irrigation,
72 relative humidity, and evapotranspiration during the growing season were reported in an earlier study (Lopes et al.

2013). In Mexico, the experiments were conducted at the Campo Experimental Norman E. Borlaug (CENEB), Cd. Obregon, where the environments were designated as yield potential, drought stress, heat stress, and heat plus drought stress. Yield potential (YP) was applied by normal sowing (late November) with optimum irrigation (total water supply > 700 mm); drought stress (D) was applied by normal planting (late November) with significantly reduced irrigation (total water supply < 300 mm); heat stress (H) was applied by late sowing (late February) with supplementary irrigation (total water supply > 700 mm) to avoid the effect of drought; and the combined heat plus drought stress (HD) was applied by delayed planting date (late February) with reduced irrigation (total water supply < 400 mm). By late sowing, the average maximum temperature during wheat growing cycle was above 32°C, almost 8°C higher than that in normal sowing cycles (Liu et al. 2019b; Pinto et al. 2010), a typical method for applying heat stress in Obregon that has been demonstrated to be successful in generating germplasm for heat stress environments such as ME5 (Reynolds et al. 1994). The experimental design was randomized lattice with two replications. Seeds were sown on raised beds of 2 m length with 2 rows (25 cm between rows) and 80 cm between the beds, with seed rates of 120 kg/ha. Five agronomic traits were measured: grain yield (GY; t ha⁻¹), thousand-grain weight (TGW, g), grain number (GN; estimated as: GY (t ha⁻¹)/ TGW (g) × 100 × 1000), days to heading (DTH, i.e. the number of days from emergence when 50% of the spike were emerged), and plant height (PH, cm from the soil surface to the tip of the spike without awns).

89

90 **Phenotypic data analysis**

91 The adjusted means for each genotype in individual environment were calculated using the MIXED procedure in SAS as described by Lopes et al. (2013). Cluster of environments based on yield performance was generated using the “cluster” package in R (Maechler 2019).

94 For the individual environment, the broad sense heritability (H^2) was estimated as:

$$95 \quad H^2 = \frac{\sigma_g^2}{\sigma_g^2 + \sigma_e^2/r}$$

96 where σ_g^2 and σ_e^2 were genotype and error variance, and r was the number of replications.

97 For the combined analysis of high-yielding and low-yielding clusters, the H^2 was estimated as:

$$98 \quad H^2 = \frac{\sigma_g^2}{\sigma_g^2 + \sigma_{ge}^2/l + \sigma_e^2/rl}$$

99 where σ_{ge}^2 was the genotype by environment interaction variance, and l was the number of environment.

100

101 **Genotyping, map construction and QTL mapping**

102 A genetic map with 475 markers (120 SSR, 211 AFLP, and 144 DArT) distributed over 29 linkage groups (LGs) had
103 been previously constructed for this population (Lopes et al. 2013). Recently, the Seri/Babax RIL population was
104 genotyped with the wheat 90K Illumina iSelect array and DArTseq platform (data available at
105 <https://data.cimmyt.org/dataset.xhtml?persistentId=hdl:11529/10548196>). After quality control (removing markers
106 with minor allele frequency < 5%, missing data > 15%, and segregation distortion $p < 0.05$), an additional 5386-90K
107 SNP and 609 DArTseq SNP -markers were added to construct the linkage map. A new genetic map was constructed
108 using QTL IciMapping v4.1 (Meng et al. 2015), which combined the map with the previous 475 markers and the new
109 SNP markers. The 90K and DArTseq markers were assigned to chromosomes based on the IWGSC RefSeq v1.0
110 genome. Within each chromosome, genetic distance was determined using the Kosambi mapping function (Kosambi
111 2016). Chromosomes were split into LGs and reordered when the gap between two adjacent markers were > 50 cM.

112 QTL mapping was performed using QTL IciMapping v4.1 software (Meng et al. 2015). Inclusive composite
113 interval mapping of additive (ICIM-ADD) function was selected as mapping method to detect additive QTL. The
114 LOD threshold for QTL detection was determined by using 1000 permutations at $P < 0.05$. Putative QTLs were
115 declared using a LOD of 2.5. QTLs detected across environments were considered the same if the distance between
116 the peak positions was less than 10 cM. Stable QTLs were declared when QTLs were identified in at least two
117 environments.

118

119 **RESULTS**

120 **Phenotypic variations across environments**

121 The phenotypic traits of the Seri/Babax RIL population showed a wide range of variation across environments
122 (**Supplementary Table 1**). The highest mean GY was found in Sohag (7.5 t ha^{-1}) and Shandaweel (7.5 t ha^{-1} , $H^2 =$
123 0.45) in Egypt, while the lowest mean GY was found in Obregon under heat stress in 2006 (1.1 t ha^{-1} , $H^2 = 0.79$; **Table**
124 **1**). Cluster analysis of the 16 environments for GY generated two main groups: the high-yielding group ($> 5.5 \text{ t ha}^{-1}$,
125 $H^2 = 0.35$) comprising Sohag and Shandaweel in Egypt, Ludhiana in India, Dongola in Sudan, and two yield potential
126 environments in Obregon, Mexico. The low-yielding group ($< 4.5 \text{ t ha}^{-1}$, $H^2 = 0.37$) encompassed Karnal in India,
127 Darab in Iran, Wad Medani in Sudan, and all drought, heat, and heat plus drought environments in Obregon, Mexico

128 **(Fig. 1b)**. The correlation analysis for GY was consistent with the results of cluster analysis, with Sohag and
129 Shandaweel in Egypt showing the highest correlation. Most of the environments showed positive correlations with
130 each other, while Wad Medani was negatively correlated with Obregon drought stress in 2004-2005 and 2005-2006
131 growing seasons (**Fig. 1c**).

132 Similar to GY, the highest average TGW (45.4 g, $H^2 = 0.54$) was observed in Sohag and Shandaweel in
133 Egypt, while the lowest TGW (26.6 g, $H^2 = 0.78$) was found in Obregon heat plus drought stress environment (**Table**
134 **1**). The highest GN (18558, $H^2 = 0.88$) was found in Ludhiana, India with a large range, and the lowest GN (3405, H^2
135 = 0.48) was observed in Obregon drought stress in 2005-2006. The phenology range was 6 to 18 days across
136 environments. In Darab, Iran, the Seri/Babax population had the longest time-112 days-for days to heading; while in
137 Obregon heat plus drought environment, the mean days to heading was 53 days. The Seri/Babax population had the
138 highest PH in Sohag and Shandaweel (107.5 cm), and lowest mean PH (71 cm) for the population in Obregon heat
139 plus drought condition. The largest and smallest range for PH was observed in Wad Medani (a range of 40 cm) and
140 Obregon heat stress in 2009 (a range of 15.5 cm), respectively.

141

142 **Trait associations across environments**

143 Across the 16 environments, GY was significantly positively correlated with TGW (r range: 0.29-0.77), GN (0.48-
144 0.94), DTH (0.32-0.69), and PH (0.24-0.73) in most of the environments with a few exceptions: GY was weakly
145 negatively correlated with TGW under Obregon heat stress (2005, 2006; $r = -0.03$), with GN under Obregon drought
146 stress (2005-2006; $r = -0.03$), and with DTH in Shandaweel of Egypt ($r = -0.13$; **Supplementary Fig. 1**). TGW and
147 GN was positively correlated with each other in 10 environments (r range: 0.19-0.55) except in Shandaweel of Egypt
148 ($r = -0.63$), Obregon yield potential (2008-2009; $r = -0.62$), drought (2005-2006; $r = -0.36$), and heat (2005, 2006; r
149 ranges from -0.32 to -0.46) conditions, where negative correlations were observed. TGW was positively correlated
150 with DTH (r range: 0.87-0.96) except in Shandaweel of Egypt ($r = -0.33$). TGW was positively correlated with PH (r
151 range: 0.90-0.97) except in Darab, Iran. GN was positively correlated with DTH (r range: 0.14-0.61) and PH (r range:
152 0.90-0.97) in all environments.

153

154 **Genetic linkage map construction**

155 We used 6470 markers comprising SSR (120), AFLP (211), DArT (144), 90K SNP (5386), and DArTseq SNP (609)
156 markers to construct the genetic map. Discarding redundant markers at the same loci, the final genetic map was 5576.5
157 cM with 1748 markers (82 SSR, 183 AFLP, 111 DArT, 1165 90K SNPs, and 207 DArTseq SNPs) arranged to 31 LGs
158 (**Fig. 2, Supplementary Table 2**). The total length of the updated map increased 3.5 folds than that of the previous
159 one Lopes et al. (2013). The A, B and D genomes harbored 768 (44%), 648 (37%) and 332 (19%) markers with a total
160 length of 1959.5, 1961.9, and 1655.1 cM, respectively. The current map provided a better coverage of all the 21
161 chromosomes than the previous one, in which only 11% of markers were distributed on D genome and no markers
162 were assigned to 3D. On the updated map, the longest LG was 4A (352.5 cM), followed by 6A (351.6 cM) and 1D
163 (350.6 cM). The shortest LG was 7Dc (4.7 cM), followed by 3Ab (7.2 cM). Marker density was lowest in 3D (16.2 cM
164 per marker), followed by 1Ba (14.1 cM per marker), and highest in 7Dc (0.9 cM per marker), followed by 1A (1.6 cM
165 per marker, **Table 2**). We obtained the physical position of 1059 mapped markers on the linkage map and projected
166 our linkage map to the IWGSC RefSeq v1.0 genome. Results showed there was a good collinearity between the
167 updated genetic map and the physical map (**Fig. 3**).

168 Of the 1748 mapped markers, significant segregation distortion (CHITEST, $p < 0.001$) was found in 68 (4%)
169 markers including 28 DArT, 25 AFLP, 9 SSR, and 6 DArTseq markers. Fifty-eight and ten markers showed distortion
170 with alleles biased towards parent Babax and Seri, respectively. The markers that showed segregation distortion in
171 favor of the Babax parent were mainly distributed on 1Bb (35.6-104.9 cM, 43 markers), while those in favor of the
172 Seri parent were mainly assigned to 3Aa (142-153 cM, 8 markers).

173 Clusters of markers from the same marker type on the map were frequently observed (**Fig. 2**). Of the 1748
174 mapped markers, 1270 markers (73%) were adjoined with markers from the same type. Marker intervals with a
175 distance < 0.5 cM between two adjacent markers from the same type were observed in 326, 26, 6, 4, and 1 intervals
176 for 90K, DArTseq, AFLP, DArT, and SSR markers, accounting for 34%, 29%, 8%, 15%, and 7% of their total intervals
177 (**Supplementary Fig. 2**). The frequency of clustering was primarily related to the number of markers from different
178 types.

179 We also built a linkage map only using the 90K SNPs to a map length of 5333 cM with 1222 non-redundant
180 90K markers. While there were many large gaps on the map with 90K SNPs (**Supplementary Table 3**), indicating
181 the necessity to include different types of markers. As was shown in **Fig. 2**, DArTseq, DArT, SSR, AFLP markers
182 could cover most of the large gaps.

183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209

QTLs identified in individual environments

A total of 166 QTLs with LOD scores from 2.5 to 23.7 were identified for GY, TGW, GN, DTH, and PH in the 16 environments with the updated genetic map (**Supplementary Table 4**). The number of QTLs was higher than the earlier map where 31 QTL were detected for the same traits (**Table 2**). The largest number of QTLs was identified on chromosome 4A (23), followed by 7Da (22). No QTL was identified on 1Ba, 2Da, 3Ab, 3Ba, 5Db, 7Bb, 7Db, and 7Dc (**Supplementary Fig. 3**).

QTLs for GY were identified on chromosome 1A, 1Bb, 1D, 2Db, 3Aa, 3Bb, 4A, 4B, 5A, 5Da, 6A, 6B, 6Da, 7A, and 7Da (LOD: 2.6-23.6; PVE: 4.8-21.3%; **Supplementary Table 4, Supplementary Fig. 3**). Seven stable QTLs on chromosomes 3Aa (128-134 cM), 3Bb (228-232 cM), 4A (268 cM, 280-286 cM, 304-308 cM), 6B (206-208 cM), and 7Da (124 cM) were detected in at least two environments (**Table 3**). The stable QTL on chromosome 4A (304-308 cM, **Fig. 4**) was detected across the yield potential (2005-2006, 2008-2009), drought (2008-2009), and heat plus drought (2009) conditions in Obregon, Mexico. QTLs for TGW were identified on chromosome 1A, 1D, 2B, 3Bb, 4A, 4B, 5A, 5B, 5Da, 6A, 6B, 7A, 7Ba, and 7Da (LOD: 3-15.4; PVE: 3-21.4%). Six stable QTLs on chromosomes 2B (44-46 cM), 3Bb (44-48 cM), 6A (218 cM), 7Ba (26-32 cM), and 7Da (108-116 cM, 124 cM) were identified (**Table 3**). The QTL on chromosome 7Da (108-116 cM) was detected in six environments across Karnal in India, Darab in Iran, and Obregon yield potential (2005-2006, 2008-2009) and drought (2005-2006, 2008-2009) conditions.

QTLs for GN were identified on chromosome 1A, 1Bb, 1D, 2A, 2B, 3Aa, 3Bb, 4A, 4B, 4Db, 5A, 6B, 6Db, 7A, and 7Da (LOD: 2.5-13.1; PVE: 3.6-26.9%). Three stable QTLs were identified on chromosomes 1Bb (72-80 cM), 4A (308 cM, **Fig. 4**), and 6B (214 cM) in Obregon, Mexico (**Table 3**), of which the QTL on 6B (214 cM) was a heat adaptive QTL. QTLs for DTH were located on chromosome 1Bb, 1D, 2A, 2B, 3D, 4B, 4Da, 5A, 5Da, 6B, 6Db, 7A, 7Ba, and 7Da (LOD: 2.7-14.4; PVE: 2.6-15.9%). Eight stable QTLs were identified on chromosome 2A (158-160 cM), 4B (70 cM, 186 cM), 5Da (38 cM), 6B (206 cM), 7Ba (20-22 cM), and 7Da (110-112 cM, 120-124 cM) across environments. QTLs for PH were identified on chromosome 1D, 2B, 3Aa, 4A, 4B, 4Db, 5A, 6A, 7Ba, and 7Da (LOD: 2.8-4.8; PVE: 6.9-12.9%). Four stable QTLs on chromosomes 2B (84 cM), 3Aa (166 cM), 4A (268-272 cM), and 4B (192-194 cM) were identified across environments (**Table 3**).

QTLs identified for high-yielding and low-yielding clusters

211 A QTL analysis based on the high- and low-yielding clusters identified 32 QTLs (LOD range: 2.6-14.9, PVE: 2.7-
212 19.7%) for GY, TGW, GN, DTH, and PH, of which 11 QTLs were new and absent in individual environments
213 (**Supplementary Table 5**). Six QTLs on chromosomes 1A, 1Bb, 2B, 4A, and 4Db were specific for high-yielding
214 cluster, 16 QTLs on chromosomes 1A, 1Bb, 4A, 4B, 4D, 6B, 7A, 7Ba, and 7Da were specific for low-yielding cluster,
215 and five QTLs on chromosomes 1D, 3Aa, 7Ba, and 7Da were common between the two clusters (**Fig. 5**).

216 QTLs for GY were found on chromosomes 1Bb (28 cM, new) and 4A (26 cM) in high-yielding cluster; on
217 4A (4 cM, 282 cM) and 4B (186 cM) in low-yielding cluster; and the common QTL was located on 1D (26 cM, new).
218 For TGW, the QTLs for low-yielding cluster were identified on chromosomes 1A (242 cM) and 7Ba (26 cM), and the
219 common QTL was identified on 7Da (114-116 cM). For GN, QTLs were located on chromosomes 1A (86 cM, new)
220 and 4A (2 cM, new) in high-yielding cluster, and on 1Bb (74 cM), 4A (158 cM, new; 268 cM), 4Db (90 cM), 6B (218
221 cM), and 7A (60 cM, new) in low-yielding cluster. For DTH, common QTLs were identified on 1D (30-32 cM) and
222 7Ba (20 cM); high-yielding cluster specific QTL was located on chromosome 4Db (102 cM, new); and low-yielding
223 cluster specific QTLs were located on chromosomes 4B (70 cM), 6B (206 cM), 7A (264 cM), 7Ba (20 cM), and 7Da
224 (114 cM). For PH, QTL in high and low yield cluster was identified on chromosomes 2B (64 cM, new) and 7Ba (130
225 cM), respectively. The common QTL for PH in both clusters was located on chromosome 3Aa (122 cM, new).

226

227 **Pleiotropic QTL**

228 Pleiotropic QTLs were observed in this study (**Fig. 6**). Among the 21 QTL hotspots for multiple traits, nine were
229 independent from DTH and/or PH QTL. Co-location of QTL for GY, TGW, and GN was on chromosome 1D at 194-
230 198 cM, where the favorable allele for GY and TGW was contributed by Babax, while for GN was from Seri. Co-
231 locations of QTLs for GY and TGW were observed on chromosomes 3Bb (228-236 cM) where the favorable allele
232 was contributed by Babax, and on chromosome 5A (26 cM) where the favorable allele was contributed by Seri. Co-
233 location of QTLs for GY and GN were on chromosomes 1A (120 cM) where the favorable allele was from Seri, and
234 on chromosome 1Bb (68-92 cM), 4A (56-62 cM, 280-286 cM, 304-308 cM), and 4B (32 cM) where favorable alleles
235 were contributed by Babax.

236

237 **DISCUSSION**

238 The rapid advances in genotyping technologies exponentially increase the number of markers available for genetic
239 studies. Genetic maps are very useful tools in the identification of molecular markers closely linked to QTLs of
240 interest, isolation of genes via map-based cloning, comparative mapping, and genome organization studies (Varshney
241 et al. 2007). In the present study, with the addition of the 90K SNP and DArTseq SNP markers, the number of mapped
242 markers (1748) on the genetic map of Seri/Babax increased 3.5 folds compared with the previous map of 475 markers
243 (Lopes et al. 2013). The final size of the linkage map was 5576.5 cM, almost 3.5-fold the length of the previous one
244 (Lopes et al. 2013). The current map provides a better coverage of the 21 chromosomes of wheat than the previous
245 one where linkage group 3D was absent (Lopes et al. 2013; McIntyre et al. 2010). Around 19% of markers were
246 assigned to D genome, which was higher than the earlier D genome with 11% of markers. Though the updated map
247 showed a better genome coverage, gaps with a distance > 50 cM still remained on chromosome 1B, 2D, 3A, 3B, 4D,
248 5D, 6D, 7B, and 7D, where small LGs were generated. Gaps of distances > 20 cM were observed almost on all LGs
249 and need to be filled with more markers for the non-centromeric regions. Of the 1748 mapped markers, 11 markers
250 were assigned to 3D, confirming the difficulty in identifying polymorphic markers for 3D, perhaps due to the large
251 common regions of 3D between the Seri and Babax (McIntyre et al. 2010). In addition, the lower frequency of
252 polymorphic markers in D genome compared with AB genome in wheat makes it difficult to have a good coverage of
253 D genome (Allen et al. 2011; Chao et al. 2009). The final map comprised of markers from AFLP, SSR, DArT, 90K,
254 and DArTseq, and clustering of markers from the same marker platform on the map were frequently observed. The
255 frequency of marker clustering was primarily related to the number of markers from different types. For example,
256 clustering of 90K markers was more frequent than other types, which was not surprising since the number of 90K
257 markers were more abundant than others. The occurrence of clusters are very common when different kinds of markers
258 are integrated into a single linkage map (Peleg et al. 2008; Semagn et al. 2006). Overall, the updated genetic map
259 provided a higher density (1.1-fold) map with more markers (3.5-fold) and a good collinearity with the IWGSC RefSeq
260 v1.0 genome, representing a step forward in mapping analysis and also in comparing QTLs with other studies using
261 90K genetic map.

262 With the updated integrated genetic map, the number of identified QTLs was increased by five-fold compared
263 with the previous study (Lopes et al. 2013) and 25 QTLs were common between the two studies (**Supplementary**
264 **table 6**). Since the Seri/Babax population was widely used for QTL studies (Mathews et al. 2008; McIntyre et al.
265 2010; Pinto et al. 2016; Pinto et al. 2010; Tahmasebi et al. 2016), we managed to locate all the reported QTLs on the

266 updated map and found that the QTLs were clustered on several big chromosome regions (**Supplementary table 6**).
267 For example, on chromosome 1Bb at 70-105 cM, there was a big cluster of QTLs for GY, TGW, GN, DTA, PH,
268 NDVI, CT, SPAD, harvest index, rate of senescence, and percentage of greenness lost. On chromosome 2B (24-84
269 cM), QTLs for GY, TGW, GN, DTH, PH, CT, maturity, grains per spike, leaf rolling, and spikelet compactness were
270 co-located. On chromosome 4A at 265-300 cM, QTLs for GY, GN, and PH in the present study were co-located with
271 QTLs for CT, NDVI, grain-filling rate, and water-soluble carbohydrates. On chromosome 5A (69-139 cM), QTLs for
272 GY, GN, DTH, PH, spikelet per spike, spikelet compactness, flowering time, maturity, and leaf rolling were clustered.
273 On chromosome 6B at 182-216 cM, QTLs for GY, TGW, GN, DTH, grain-filling rate, spikelet per spike, spikelet
274 compactness, grains per spike, and percentage of greenness lost at mid grain-filling stage were frequently identified.
275 Pinto et al. (2010) reported six common QTLs for drought and heat stress on chromosome 1B-a, 2B-a, 3B-b, 4A-a,
276 4B-b, and 7A-a in the Seri/Babax population in Obregon, Mexico environments, while in the present study, the QTL
277 on chromosome 1B-a can be detected in Obregon heat stress in 2006 (71-75 cM on 1Bb in the present study); the QTL
278 on chromosome 2B-a was detected in Wad Medani, Dongola, Darab, and Obregon heat stress in 2009 (39-47 cM on
279 2B); the QTL on chromosome 3B-b was detected in Obregon yield potential (2005-2006), drought (2004-2005), and
280 heat (2006) conditions (219-235 cM on 3Bb); the QTL on chromosome 4A-a was detected across Obregon heat stress
281 (2006, 2009), Ludhiana, Shandaweel, and Sohag (265-269 cM on 4A); the QTL on chromosome 4B-b was identified
282 in Obregon heat plus drought stress (2009), Darab, Dongola, and Ludhiana (189-195 cM on 4B), the QTL on 7A-a
283 was detected in Wad Medani and Karnal (265-267 cM on 7A). These results suggested that the experiments and
284 breeding conducted in drought and heat stress environments in Obregon, Mexico could be used to predict the
285 performance of the germplasm in other parts of the world. Reynolds et al. (1994) also reported that the typical heat
286 stress in Obregon was demonstrated to be successful in generating germplasm for heat stressed environments such as
287 ME5 comprising India and Sudan (Hodson and White 2007; Rajaram et al. 1993).

288 Typically, there is a trade-off between grain weight and grain number, making it difficult to improve yield
289 through increasing grain weight and grain number at the same time (Griffiths et al. 2015; Sukumaran et al. 2018a;
290 Sukumaran et al. 2018b). However, in the present study, TGW and GN were observed to be positively correlated, and
291 both TGW and GN were positively correlated with GY in most environments. Several QTLs co-located for GY and
292 GN were found on chromosome 1A (120 cM), 1Bb (68-92 cM), 4A (56-62, 280-286, 304-308 cM), and 4B (32 cM),
293 where QTLs for GY and GN shared the favorable allele from the same parent. QTLs co-located for GY and TGW

294 were on chromosome 3Bb (228-236 cM) and 5A (26 cM), where the same parent contributed the favorable allele for
295 both GY and TGW. Among the QTLs co-located for TGW and GN, most of the favorable alleles for TGW and GN
296 were contributed by different parents, while the QTL on chromosome 6B (194-200 cM) shared the favorable allele
297 from the same parent for TGW and GN, indicating it is possible to improve yield through TGW QTL complemented
298 by adding correlated GN QTL (Groos et al. 2003).

299 The stable heat adaptive QTL on chromosome 1D (84 cM) for GY could be projected to 484-493 Mb on the
300 IWGSC RefSeq v1.0 genome, where genes were located for disease resistance, heat shock family protein, and early
301 flowering. The QTL on chromosome 2Db (110 cM) for GY was previously identified for photochemical reflectance
302 index (PRI) in a CIMMYT spring wheat population (WAMI) under yield potential condition (Liu et al. 2019a). The
303 QTL on chromosome 3Bb (228-236 cM) for GY and TGW was very close to a locus associated with multiple traits
304 (GY, maturity, and chlorophyll content at vegetative state) in the temperate irrigated environments in spring wheat
305 (Sukumaran et al. 2015). The QTL on chromosome 4A (308 cM) for GY and GN could be mapped to 531-535 Mb in
306 physical map, which contains 34 genes including heat-shock protein genes. The TGW QTL on chromosome 7Ba (26-
307 32 cM) was located at 3.7-5.9 Mb on the physical map, which harbored 33 genes. The GN QTL on chromosome 1A
308 (86 cM) was associated with ground cover in winter wheat (Gao et al. 2016). The GN QTL on chromosome 6Db (8
309 cM) was co-located with a QTL for DTH in a synthetic derived RIL population (Liu et al. 2019b).

310 The Seri/Babax population was initially developed in Obregon, Mexico, so the lowest range of DTH (8 days)
311 and PH (15-23 cm) was observed in the Obregon environments in this study. When grown in international trials, longer
312 DTH and higher PH with a wider range were observed in most environments. For DTH, the QTL on chromosome 4Da
313 was previously reported for maturity in spring wheat (Sukumaran et al. 2015). The QTL on chromosome 5D (38 cM)
314 was located at a distance (about 60 Mb) from the *Vrn-D4* locus (Yoshida et al. 2010). The stable QTL for DTH on
315 chromosome 7Ba (20-22 cM) was co-located with a previously reported QTL for kernel number per spike (Gao et al.
316 2015). Projecting this QTL to the physical map, it was located at the *Vrn-B3* locus (Yan et al. 2006), which could
317 accelerate flowering and bypass the vernalization requirement of its dominant allele if present. This QTL was
318 contributed by the parent Seri, and has never been reported before by other studies using Seri/Babax. The other QTLs
319 for DTH are most likely to be associated with earliness *per se* genes that act independently of environmental signals,
320 and are usually responsible for fine-tuning flowering time (Flood and Halloran 1984). For PH, the locus on
321 chromosome 4B (192-194 cM) was associated with GY in a CIMMYT spring wheat panel (Sukumaran et al. 2013).

322 This locus can be projected to 3.5-4.7 Mb on the physical map, which was very close to the *Rht-B1* gene (Pearce et al.
323 2011). The PH QTL on chromosome 5A (72 cM) was also associated with *Cephalosporium* stripe resistance in wheat
324 (Quincke et al. 2011).

325 This study updated the genetic map of the widely used Seri/Babax RIL population with the 90K and DArTseq
326 SNPs. The updated genetic map has a good genome coverage, higher marker density, and a good collinearity with the
327 physical map, and increased the number of QTLs compared with the earlier studies. A larger number of QTLs were
328 identified with this genetic map across various environments, and some of them were stable in different environments.
329 QTLs that are unique to the current study, verify the importance of an updated genetic map for QTL detection. This
330 updated genetic map is expected to serve as a platform for further genetic analyses of important quantitative traits,
331 map-based cloning, and marker-assisted selection studies using Seri/Babax RIL population.

332

333 **AUTHOR CONTRIBUTIONS**

334 SS designed the current study, ML and MR provided the phenotypic data, CS contributed with the explanation of
335 DArTseq data, SD and MR genotyped the population, CL, SM and SS created the genetic maps and analyzed the
336 data, CL and SS wrote the manuscript. All authors reviewed and accepted the contents.

337

338 **ACKNOWLEDGEMENTS**

339 This work was implemented by CIMMYT as part of IWYP-HUB, made possible by the generous support of
340 International Wheat Yield Partnership (IWYP). Any opinions, findings, conclusion, or recommendations expressed in
341 this publication are those of the authors and do not necessarily reflect the view of donors. We also acknowledge Dr.
342 Marc Ellis for providing DArTseq data for the population. Thanks are due to the staff at Wheat Physiology lab at
343 CENEB, Cd. Obregon, México.

344

345 **CONFLICT OF INTEREST**

346 The authors declare no conflict of interest.

347

348 **Reference**

349 Allen AM, Barker GL, Berry ST, Coghill JA, Gwilliam R, Kirby S, Robinson P, Brenchley RC, D'Amore R,
350 McKenzie N (2011) Transcript-specific, single-nucleotide polymorphism discovery and linkage analysis in
351 hexaploid bread wheat (*Triticum aestivum* L.). *Plant biotechnology journal* 9 (9):1086-1099

352 Bennett D, Reynolds M, Mullan D, Izanloo A, Kuchel H, Langridge P, Schnurbusch T (2012) Detection of two
353 major grain yield QTL in bread wheat (*Triticum aestivum* L.) under heat, drought and high yield potential
354 environments. *Theoretical and Applied Genetics* 125 (7):1473-1485

355 Chao S, Zhang W, Akhunov E, Sherman J, Ma Y, Luo M-C, Dubcovsky J (2009) Analysis of gene-derived SNP
356 marker polymorphism in US wheat (*Triticum aestivum* L.) cultivars. *Molecular Breeding* 23 (1):23-33

357 Doerge RW (2002) Multifactorial genetics: Mapping and analysis of quantitative trait loci in experimental
358 populations. *Nature Reviews Genetics* 3 (1):43

359 FAOstat (2018) The Statistics Division, Food and Agriculture Organization of the United Nations.

360 Flood RG, Halloran GM (1984) Basic Development Rate in Spring Wheat1. *Agronomy Journal* 76 (2):260-264.
361 doi:10.2134/agronj1984.00021962007600020021x

362 Gao F, Liu J, Yang L, Wu X, Xiao Y, Xia X, He Z (2016) Genome-wide linkage mapping of QTL for physiological
363 traits in a Chinese wheat population using the 90K SNP array. *Euphytica* 209 (3):789-804.
364 doi:10.1007/s10681-016-1682-6

365 Gao F, Wen W, Liu J, Rasheed A, Yin G, Xia X, Wu X, He Z (2015) Genome-wide linkage mapping of QTL for
366 yield components, plant height and yield-related physiological traits in the Chinese wheat cross Zhou
367 8425B/Chinese Spring. *Frontiers in Plant Science* 6:1099

368 Golabadi M, Arzani A, Maibody SM, Tabatabaei BS, Mohammadi S (2011) Identification of microsatellite markers
369 linked with yield components under drought stress at terminal growth stages in durum wheat. *Euphytica*
370 177 (2):207-221

371 Griffiths S, Wingen L, Pietragalla J, Garcia G, Hasan A, Miralles D, Calderini DF, Ankleshwaria JB, Waite ML,
372 Simmonds J (2015) Genetic dissection of grain size and grain number trade-offs in CIMMYT wheat
373 germplasm. *PloS one* 10 (3):e0118847

374 Groos C, Robert N, Bervas E, Charmet G (2003) Genetic analysis of grain protein-content, grain yield and thousand-
375 kernel weight in bread wheat. *Theoretical and Applied Genetics* 106 (6):1032-1040

376 Hodson D, White J (2007) Use of spatial analyses for global characterization of wheat-based production systems.
377 Journal of Agricultural Science Cambridge 145 (1):115-125

378 Kirigwi F, Van Ginkel M, Brown-Guedira G, Gill B, Paulsen GM, Fritz A (2007) Markers associated with a QTL
379 for grain yield in wheat under drought. *Molecular Breeding* 20 (4):401-413

380 Kosambi DD (2016) The estimation of map distances from recombination values. In: DD Kosambi. Springer, pp
381 125-130

382 Liu C, Pinto F, Cossani CM, Sukumaran S, Reynolds PM (2019a) Spectral reflectance indices as proxies for yield
383 potential and heat stress tolerance in spring wheat: heritability estimates and marker-trait associations.
384 *Frontiers of Agricultural Science and Engineering*. doi:10.15302/j-fase-2019269

385 Liu C, Sukumaran S, Claverie E, Sansaloni C, Dreisigacker S, Reynolds M (2019b) Genetic dissection of heat and
386 drought stress QTLs in phenology-controlled synthetic-derived recombinant inbred lines in spring wheat.
387 *Molecular Breeding* 39 (3):34

388 Lopes MS, Reynolds MP, McIntyre CL, Mathews KL, Kamali MJ, Mossad M, Feltaous Y, Tahir IS, Chatrath R,
389 Ogbonnaya F (2013) QTL for yield and associated traits in the Seri/Babax population grown across several
390 environments in Mexico, in the West Asia, North Africa, and South Asia regions. *Theoretical and applied*
391 *genetics* 126 (4):971-984

392 Maccaferri M, Sanguineti MC, Corneti S, Ortega JLA, Salem MB, Bort J, DeAmbrogio E, del Moral LFG,
393 Demontis A, El-Ahmed A (2008) Quantitative trait loci for grain yield and adaptation of durum wheat
394 (*Triticum durum* Desf.) across a wide range of water availability. *Genetics* 178 (1):489-511

395 Maechler M (2019) Finding Groups in Data": Cluster Analysis Extended Rousseeuw et. R Packag version 20 6

396 Mathews KL, Malosetti M, Chapman S, McIntyre L, Reynolds M, Shorter R, van Eeuwijk F (2008) Multi-
397 environment QTL mixed models for drought stress adaptation in wheat. *Theoretical and Applied Genetics*
398 117 (7):1077-1091. doi:10.1007/s00122-008-0846-8

399 McIntyre CL, Mathews KL, Rattey A, Chapman SC, Drenth J, Ghaderi M, Reynolds M, Shorter R (2010) Molecular
400 detection of genomic regions associated with grain yield and yield-related components in an elite bread
401 wheat cross evaluated under irrigated and rainfed conditions. *Theoretical and Applied Genetics* 120
402 (3):527-541. doi:10.1007/s00122-009-1173-4

403 Meng L, Li H, Zhang L, Wang J (2015) QTL IciMapping: Integrated software for genetic linkage map construction
404 and quantitative trait locus mapping in biparental populations. *The Crop Journal* 3 (3):269-283.
405 doi:10.1016/j.cj.2015.01.001

406 Olivares-Villegas JJ, Reynolds MP, McDonald GK (2007) Drought-adaptive attributes in the Seri/Babax hexaploid
407 wheat population. *Functional Plant Biology* 34 (3):189-203

408 Pearce S, Saville R, Vaughan SP, Chandler PM, Wilhelm EP, Sparks CA, Al-Kaff N, Korolev A, Boulton MI,
409 Phillips AL, Hedden P, Nicholson P, Thomas SG (2011) Molecular Characterization of *Rht-1*
410 Dwarfing Genes in Hexaploid Wheat. *Plant Physiology* 157 (4):1820-1831. doi:10.1104/pp.111.183657

411 Peleg Z, Saranga Y, Suprunova T, Ronin Y, Röder MS, Kilian A, Korol AB, Fahima T (2008) High-density genetic
412 map of durum wheat × wild emmer wheat based on SSR and DArT markers. *Theoretical and Applied*
413 *Genetics* 117 (1):103

414 Pinto RS, Lopes MS, Collins NC, Reynolds MP (2016) Modelling and genetic dissection of staygreen under heat
415 stress. *Theor Appl Genet* 129 (11):2055-2074. doi:10.1007/s00122-016-2757-4

416 Pinto RS, Reynolds MP, Mathews KL, McIntyre CL, Olivares-Villegas J-J, Chapman SC (2010) Heat and drought
417 adaptive QTL in a wheat population designed to minimize confounding agronomic effects. *Theoretical and*
418 *Applied Genetics* 121 (6):1001-1021

419 Quarrie S, Pekic Quarrie S, Radosevic R, Rancic D, Kaminska A, Barnes J, Leverington M, Ceoloni C, Dodig D
420 (2006) Dissecting a wheat QTL for yield present in a range of environments: from the QTL to candidate
421 genes. *Journal of Experimental Botany* 57 (11):2627-2637

422 Quincke MC, Peterson CJ, Zemetra RS, Hansen JL, Chen J, Riera-Lizarazu O, Mundt CC (2011) Quantitative trait
423 loci analysis for resistance to *Cephalosporium stripe*, a vascular wilt disease of wheat. *Theoretical and*
424 *applied genetics* 122 (7):1339-1349

425 Rajaram S, Van Ginkel M, Fischer R CIMMYT's wheat breeding mega-environments (ME). In: *Proceedings of the*
426 *8th International wheat genetic symposium, 1993.*

427 Reynolds M, Balota M, Delgado M, Amani I, Fischer R (1994) Physiological and morphological traits associated
428 with spring wheat yield under hot, irrigated conditions. *Functional Plant Biology* 21 (6):717-730

429 Reynolds M, Foulkes MJ, Slafer GA, Berry P, Parry MAJ, Snape JW, Angus WJ (2009) Raising yield potential in
430 wheat. *Journal of Experimental Botany* 60 (7):1899-1918. doi:10.1093/jxb/erp016

431 Reynolds M, Tuberosa R (2008) Translational research impacting on crop productivity in drought-prone
432 environments. *Curr Opin Plant Biol* 11 (2):171-179. doi:10.1016/j.pbi.2008.02.005

433 Rosegrant MW, Cline SA (2003) Global food security: challenges and policies. *Science* 302 (5652):1917-1919.
434 doi:10.1126/science.1092958

435 Semagn K, Bjørnstad Å, Skinnes H, Marøy AG, Tarkegne Y, William M (2006) Distribution of DArT, AFLP, and
436 SSR markers in a genetic linkage map of a doubled-haploid hexaploid wheat population. *Genome* 49
437 (5):545-555

438 Sukumaran S, Dreisigacker S, Lopes M, Chavez P, Reynolds MP (2015) Genome-wide association study for grain
439 yield and related traits in an elite spring wheat population grown in temperate irrigated environments.
440 *Theoretical and Applied Genetics* 128 (2):353-363

441 Sukumaran S, Ellis MH, Lopes MS, Pinto S, Bonnett D, Dreisigacker S, Mullan D, Reynolds MP (2013) Wheat
442 mapping populations available at CIMMYT for yield potential research. *Wheat Yield Consortium* 782013

443 Sukumaran S, Lopes M, Dreisigacker S, Reynolds M (2018a) Genetic analysis of multi-environmental spring wheat
444 trials identifies genomic regions for locus-specific trade-offs for grain weight and grain number.
445 *Theoretical and Applied Genetics* 131 (4):985-998. doi:10.1007/s00122-017-3037-7

446 Sukumaran S, Reynolds MP, Sansaloni C (2018b) Genome-wide association analyses identify QTL hotspots for
447 yield and component traits in durum wheat grown under yield potential, drought, and heat stress
448 environments. *Frontiers in plant science* 9:81

449 Tahmasebi S, Heidari B, Pakniyat H, McIntyre CL (2016) Mapping QTLs associated with agronomic and
450 physiological traits under terminal drought and heat stress conditions in wheat (*Triticum aestivum* L.).
451 *Genome* 60 (1):26-45

452 Trethowan RM, Mujeeb-Kazi A (2008) Novel Germplasm Resources for Improving Environmental Stress Tolerance
453 of Hexaploid Wheat. *Crop Science* 48:1255-1265. doi:10.2135/cropsci2007.08.0477

454 Varshney RK, Mahendar T, Aggarwal RK, Börner A (2007) Genic molecular markers in plants: development and
455 applications. In: *Genomics-assisted crop improvement*. Springer, pp 13-29

456 Yan L, Fu D, Li C, Blechl A, Tranquilli G, Bonafede M, Sanchez A, Valarik M, Yasuda S, Dubcovsky J (2006) The
457 wheat and barley vernalization gene *VRN3* is an orthologue of *FT*. *Proceedings of the National Academy*
458 *of Sciences* 103 (51):19581-19586

459 Yano M, Sasaki T (1997) Genetic and molecular dissection of quantitative traits in rice. In: Sasaki T, Moore G (eds)
460 Oryza: From Molecule to Plant. Springer Netherlands, Dordrecht, pp 145-153. doi:10.1007/978-94-011-
461 5794-0_14

462 Yoshida T, Nishida H, Zhu J, Nitcher R, Distelfeld A, Akashi Y, Kato K, Dubcovsky J (2010) Vrn-D4 is a
463 vernalization gene located on the centromeric region of chromosome 5D in hexaploid wheat. Theoretical
464 and Applied Genetics 120 (3):543-552

465

466

Tables and figures

467 **Table 1.** Adjusted means and range of grain yield (GY, t ha⁻¹), thousand grain weight (TGW, g), grain number (GN), days to heading (DTH, d), and plant height
 468 (PH, cm) collected on the Seri/Babax RIL population comprising 156 entries in 16 environments from five countries (Egypt, India, Iran, Mexico, and Sudan)
 469 during 2004-2005, 2005-2006, and 2008-2009 growing seasons.

Abbreviations	Country & Location	Year	Mean (Range)				
			GY (t ha ⁻¹)	TGW (g)	GN	DTH (d)	PH (cm)
EgySha09	Egypt Shandaweel	2008-09	7.5 (6.6-8.3)	45.4 (42.9-48.4)	16808 (15026-18674)	94 (87-102)	108 (97-119)
EgySoh09	Egypt Sohag	2008-09	7.5 (4.9-9.7)	45.4 (36.1-54.8)	16697 (10754-23453)	94 (87-103)	108 (94-122)
IndKar09	India Karnal	2008-09	4.1 (2.3-5.9)	35.1 (26-47)	11697 (6570-17688)	93 (86-102)	100 (88-109)
IndLud09	India Ludhiana	2008-09	6.6 (2.4-10.4)	35.8 (29.7-44.3)	18558 (5945-29194)	91 (85-103)	99 (78-111)
IraDar09	Iran Darab	2008-09	4.5 (2.5-5.9)	34.6 (15-79.8)	13229 (6551-29540)	112 (109-114)	-
MexObrD05	Mexico Obregon_Drought	2004-05	3.6 (2.8-4.2)	35.2 (29.8-42.8)	10268 (9018-11363)	-	-
MexObrD06	Mexico Obregon_Drought	2005-06	3.2 (2.3-4.0)	34 (24.8-43.7)	3405 (2463-4556)	-	-
MexObrD09	Mexico Obregon_Drought	2008-09	3.8 (2.4-5.1)	36.8 (31.7-44.1)	10457 (5820-14396)	72 (67-75)	96 (87-103)
MexObrH05	Mexico Obregon_Heat	2005	1.8 (1.0-2.8)	33.3 (28.1-38)	5530 (2872-9027)	-	-
MexObrH06	Mexico Obregon_Heat	2006	1.2 (0.9-1.5)	28.9 (23.9-35.2)	11220 (7969-14200)	-	-
MexObrH09	Mexico Obregon_Heat	2009	3.5 (1.8-4.4)	29.2 (24.2-41.6)	12019 (6712-15035)	53 (49-57)	78 (71-86)
MexObrHD09	Mexico Obregon_Heat + Drought	2009	2.7 (1.5-3.6)	26.6 (20.2-34.6)	10061 (6024-13990)	53 (48-57)	71 (63-82)
MexObrYP06	Mexico Obregon_Yield potential	2005-06	5.5 (4.7-6.0)	43.3 (36-50.6)	12778 (10855-14976)	-	-
MexObrYP09	Mexico Obregon_Yield potential	2008-09	6.6 (4.6-8.5)	41 (35.2-49.6)	16025 (10899-20819)	78 (73-81)	102 (91-114)
SudDon09	Sudan Dongola	2008-09	6.7 (4.6-8.8)	42.7 (36.5-52.1)	15703 (10387-21292)	64 (56-73)	88 (77-103)
SudWadM09	Sudan Wad Medani	2008-09	2.7 (1.6-3.9)	36 (25.6-45)	7667 (4098-11894)	55 (48-67)	76 (46-86)

470

471 **Table 2.** Genetic length, number of markers, average marker interval between two adjacent markers, and number of
 472 QTLs identified for (GY, t ha⁻¹), thousand grain weight (TGW, g), grain number (GN), days to heading (DTH, d), and
 473 plant height (PH, cm) on the 31 linkage groups of the Seri/Babax RIL population linkage map in the present study
 474 compared with the previous genetic map (Lopes et al. 2013).

The updated map						The previous map					
LG	Chr/ Arm	Length (cM)	Marker number	Interval (cM)	QTL num ber	LG	Chr/ Arm	Length (cM)	Marker number	Interval (cM)	QTL num ber
1	1A	269.3	172	1.6	6	1	1A	119.9	21	5.7	1
2	1Ba	168.7	12	14.1		2	1B	85.0	45	1.9	
3	1Bb	249.8	59	4.2	6						
4	1D	350.6	84	4.2	9	3	1Da	132.2	30	4.4	1
						4	1Db	9.1	2	4.5	
5	2A	171.8	44	3.9	3	5	2Aa	37.0	2	18.5	
						6	2Ab	29.7	5	5.9	
						7	2Ac	0.4	3	0.1	
						8	2Ad	29.8	5	6.0	
6	2B	260.7	141	1.9	11	9	2B	99.0	43	2.3	3
7	2Da	100.6	28	3.6		10	2D	74.6	14	5.3	1
8	2Db	124.6	31	4.0	1						
9	3Aa	211.2	46	4.6	5	11	3Aa	49.9	15	3.3	1
10	3Ab	7.2	4	1.8		12	3Ab	13.6	9	1.5	
11	3Ba	52.4	14	3.7		13	3B	142.4	37	3.8	
12	3Bb	314.6	94	3.4	8						
13	3D	178.7	11	16.2	1						
14	4A	352.5	125	2.8	23	14	4A	110.9	43	2.6	5
15	4B	201.4	91	2.2	16	15	4B	70.1	18	3.9	3
16	4Da	43.2	6	7.2	1	16	4D	12.3	4	3.1	1
17	4Db	132.0	20	6.6	2						
18	5A	295.9	129	2.3	7	17	5A	76.0	22	3.5	2
19	5B	171.8	80	2.2	1	18	5B	14.5	17	0.9	1
20	5Da	139.9	20	7.0	5	19	5Da	25.5	2	12.7	1
21	5Db	26.9	14	1.9		20	5Db	13.0	3	4.3	1
22	6A	351.6	109	3.2	5	21	6Aa	76.2	25	3.0	2
						22	6Ab	44.6	11	4.1	
23	6B	299.9	114	2.6	14	23	6B	90.3	38	2.4	2
24	6Da	157.6	42	3.8	1	24	6Da	41.4	4	10.4	2
25	6Db	110.2	20	5.5	2	25	6Db	12.3	4	3.1	
26	7A	300.1	139	2.2	5	26	7A	115.2	35	3.3	
27	7Ba	173.0	28	6.2	12						
28	7Bb	69.7	15	4.7		27	7B	15.6	6	2.6	
29	7Da	166.7	30	5.6	22	28	7Da	19.0	6	3.2	
30	7Db	119.4	21	5.7		29	7Db	52.6	6	8.8	4
31	7Dc	4.7	5	0.9							
Total		5576.5	1748	3.2	166			1612	475	3.4	31

475

476 **Table 3.** Stable QTLs (identified in more than two environments) for grain yield (GY, t ha⁻¹), thousand-grain weight (TGW, g), grain number (GN), days to heading
 477 (DTH, d), and plant height (PH, cm) in the Seri/Babax RIL population comprising 156 entries. PVE: phenotypic variance explained by QTL.

Trait	L G	Ch r	Peak position (cM)	Marker interval (cM)	Left marker	Right marker	LOD	PVE (%)	Favorable parent
GY	9	3Aa	130	127-137	<i>aca-cta-13</i>	<i>1089533</i>	3.1	5.4	Seri
	12	3Bb	230	219-235	<i>wPt-1804</i>	<i>gwm301e</i>	6.5	10.7	Babax
	14	4A	268	265-269	<i>wmc048d</i>	<i>act-cag-3</i>	10	16.6	Babax
	14	4A	284	273-293	<i>gUa44</i>	<i>1089414</i>	6.3	14.8	Babax
	14	4A	308	307-311	<i>wmc048c</i>	<i>Ra_c7973_1185</i>	23.7	21.3	Babax
	23	6B	206	203-209	<i>agc-cta-4</i>	<i>agg-ctg-8</i>	5.4	8.4	Seri
	29	7Da	124	123-125	<i>D_contig12156_209</i>	<i>wsnp_Ex_c4637_8299644</i>	6.2	10.5	Seri
TGW	6	2B	44	37-47	<i>wPt-7320</i>	<i>acc-ctc-2</i>	4.7	9	Babax
	12	3Bb	44	43-53	<i>barc147</i>	<i>barc087</i>	4.1	8.1	Babax
	22	6A	218	217-219	<i>wsnp_Ku_c26784_3674771_2</i>	<i>998518</i>	2.9	6.2	Babax
	27	7Ba	28	23-33	<i>wsnp_Ex_c11658_1877308_6</i>	<i>wsnp_JD_c1285_1848292</i>	3.5	9.6	Babax
	29	7Da	114	99-123	<i>Excalibur_c22419_460</i>	<i>1068196</i>	7.7	20.1	Seri
GN	29	7Da	124	123-125	<i>D_contig12156_209</i>	<i>wsnp_Ex_c4637_8299644</i>	7.1	11.5	Seri
	3	1Bb	80	71-83	<i>act-ctc-7</i>	<i>aac-ctg-4</i>	8	13.9	Babax
	14	4A	308	307-311	<i>1072050</i>	<i>Ra_c7973_1185</i>	5.8	15.3	Babax
DTH	23	6B	214	209-225	<i>wPt-4924</i>	<i>barc0178</i>	4.1	6.9	Seri
	5	2A	160	153-167	<i>RFL_Contig5277_480</i>	<i>cf0050</i>	4.2	4.1	Babax
	15	4B	70	69-71	<i>Tdurum_contig47552_957</i>	<i>Excalibur_c65023_62</i>	2.7	4.9	Seri
	15	4B	186	183-187	<i>gwm006a</i>	<i>aac-ctc-9</i>	5.6	8.9	Seri
	20	5Da	38	27-43	<i>Kukri_c444_833</i>	<i>Excalibur_c11929_1019</i>	7.8	9.4	Babax
	23	6B	206	201-209	<i>agc-cta-4</i>	<i>aac-ctc-3</i>	7.2	6.8	Babax
	27	7Ba	20	17-25	<i>IACX198</i>	<i>wsnp_Ex_c11658_1877308_6</i>	14.1	15	Seri
	29	7Da	112	107-117	<i>Excalibur_c22419_460</i>	<i>1068196</i>	7.6	15.2	Babax
	29	7Da	124	117-125	<i>Excalibur_c22419_460</i>	<i>wsnp_Ex_c4637_8299644</i>	14.4	16	Babax
	PH	6	2B	84	81-87	<i>aca-ctg-1</i>	<i>wPt-5680</i>	4.2	8.6
9		3Aa	166	161-181	<i>1086318</i>	<i>RAC875_c21944_117</i>	3	7.6-7.8	Seri
14		4A	268	265-277	<i>wmc048d</i>	<i>wmc048c</i>	3.4-4.8	8.1-13	Seri

478

15	4B	192	191-195	<i>aag-cta-5</i>	<i>wmc048a</i>	3.2- 4.2	7.7-8.8	Seri
----	----	-----	---------	------------------	----------------	-------------	---------	------

479 **Figure 1.** Phenotyping the Seri/Babax RIL population comprising 156 entries was conducted in (a) several
480 environments, (b) clusters of high-yielding and low-yielding environments, and (c) and phenotypic correlations of
481 grain yield between the environments.

482 **Figure 2.** Genetic map and marker distribution of the Seri/Babax RIL population on 31 linkage groups.

483 **Figure 3.** Updated linkage map of the Seri/Babax population (left) projected to the physical map of IWGSC RefSeq
484 genome v1.0 (right) showing a good collinearity between the genetic map and the physical map.

485 **Figure 4.** A stable QTL on chromosome 4A detected for grain yield (GY, t ha⁻¹) and grain number (GN) under yield
486 potential (YP), drought (D), and head plus drought (HD) environments in Obregon, Mexico during 2004-2005,
487 2005-2006, and 2008-2009 growing seasons.

488 **Figure 5.** Venn diagram illustrating the common and specific QTLs for high-yielding and low-yielding environment
489 clusters of the Seri/Babax RIL population. Subscript numbers indicate the *centi morgan* (cM) position on the linkage
490 map, followed by traits. GY: grain yield (t ha⁻¹); TGW: thousand-grain weight (g); GN: grain number; DTH: days to
491 heading (d); PH: plant height (cm).

492 **Figure 6.** Venn diagram illustrating the common genomic regions for grain yield (GY, t ha⁻¹), thousand-grain weight
493 (TGW, g), and grain number (GN), days to heading (DTH, d) and plant height (PH, cm). Subscript numbers indicate
494 the *centi morgan* (cM) position on the linkage map. For the GY, TGW, and GN QTLs independent from DTH and/or
495 PH, the red up arrows indicates the favorable allele contributed by the same parent, and the black down arrows
496 indicates the favorable allele contributed by different parents.

497 **Supplementary Figure 1.** Phenotypic correlations among grain yield (GY, t ha⁻¹), thousand grain weight (TGW,
498 g), grain number (GN), days to heading (DTH, d), and plant height (PH, cm) of the Seri/Babax RIL population
499 grown in different environments.

500 **Supplementary Figure 2.** Distribution of markers between two consecutive loci along genetic distance over all
501 chromosomes.

502 **Supplementary Figure 3.** An updated genetic map of Seri/Babax RIL population with QTLs identified for grain
503 yield (GY, t ha⁻¹), thousand-grain weight (TGW, g), grain number (GN), days to heading (DTH, d), and plant height

504 (PH, cm) in different environments. Markers in red and green color are distorted markers in favor of Babax and Seri
505 alleles, respectively.

506 **Supplementary Tables**

507

508 **Supplementary Table 1.** Phenotypic data including grain yield (GY, t ha⁻¹), thousand-grain weight (TGW, g), grain
509 number (GN), days to heading (DTH, d), and plant height (PH, cm) of the Seri/Babax RIL population grown in 16
510 environments from five countries.

511 **Supplementary Table 2.** Linkage map information of the Seri/Babax population. Seri is coded as "0", Babax is coded
512 as "2", and missing data is coded as "-1".

513 **Supplementary Table 3.** Linkage map information of the Seri/Babax population using 90K SNPs. Seri is coded as
514 "0", Babax is coded as "2", and missing data is coded as "-1".

515 **Supplementary Table 4.** QTLs detected for grain yield (GY, t ha⁻¹), thousand-grain weight (TGW, g), grain number
516 (GN), days to heading (DTH, d), and plant height (PH, cm) of the the Seri/Babax RIL population in 16 individual
517 environments.

518 **Supplementary Table 5.** QTLs detected for grain yield (GY, t ha⁻¹), thousand-grain weight (TGW, g), grain number
519 (GN), days to heading (DTH, d), and plant height (PH, cm) of the Seri/Babax RIL population in high-yielding and
520 low-yielding clusters.

521 **Supplementary Table 6.** Comparison of QTLs detected in the present study with the previous studies using the
522 Seri/Babax RIL population.